
a **Lines only**


HT fits with MAD ≤ 15° (as in Fig. 7r)



N = 11 boulders · R = 5.31
Ro95 5.28 · Ro99 6.25
CLUSTERED (fail)

b **Lines + planes**

unresolved specimens as remagnetization circles



N = 11 boulders · R = 6.09
Ro95 5.28 · Ro99 6.25
CLUSTERED (fail)

c **Boulder means (lines + planes)**

Boulder	D/I (°)	lines	planes
0	295.7/+19.8	13	0
1	124.7/-34.3	11	0
1.	102.0/ -2.2	14	0
2	30.3/+25.9	11	0
3	330.4/-10.3	1	10
4	57.6/-21.7	2	5
5	345.7/+51.5	12	0
6	34.9/ +0.7	3	0
7	33.0/-36.7	9	0
8	81.2/-16.6	7	0
9	14.5/-42.3	9	0

Boulders 3 and 4 do not isolate a stable endpoint by 450–510 °C; their arcs are fitted as great circles (orange) and combined with the resolved lines (McFadden & McElhinny, 1988). Arrows: line-only mean → lines + planes mean. Both boulders move toward the northerly, shallow cluster, so R rises from 5.31 to 6.09 and the fail is strengthened.

specimen line (filled lower / open upper) · remagnetization circle (dashed: upper hem.) · boulder mean (lines) · boulder mean (lines + planes) · GAD normal / reversed (open)