Table 1. The activity of non-antibiotics (active substances and medicinal products) with and without the efflux pump inhibitor PAβN against standard Gram-negative strains.

Table body
Caption (title + notes)
Merged (semantic unit)

Strains	MIC (mg/l)				
	alen/Ostenil [a] (+PAβN) [b]	tic/Apo-Clodin [a] (+PAβN)[b]	amit/Amitriptylinum [a] (+PAβN)[b]	nic/Niglostin [a] (+PAβN)[b]	crb/Carbosin [a] (+PAβN)[b]
E. coli ATCC 25922	>800 **(200/100)**	>800 **(25)**	200 **(50)**	>800 **(100)**	>800 (>800)
E. coli NCTC 8196	>800 **(200/100)**	>800 **(25)**	200 **(50)**	>800 **(100)**	>800 (>800)
E. coli NCTC 10538	>800 **(200/100)**	>800 **(25)**	400 **(25)**	>800 **(100)**	>800 (>800)
K. pneumoniae ATCC 13883	>800 **(200)**	>800 **(25)**	200 **(50)**	>800 **(200)**	>800 (>800)
K. pneumoniae ATCC 700603	>800 (800)	>800 **(100)**	400 **(100)**	>800 (>800)	>800 (>800)
P. vulgaris ATCC 13315	>800 **(400)**	>800 (>800)	800 (800)	>800 (>800)	>800 (>800)
P. aeruginosa ATCC 27853	200 **(25)**	>800 (>800)	800 **(25)**	>800 (>800)	800 **(200)**
P. aeruginosa NCTC 6749	100 **(25)**	>800 (>800)	800 (400)	>800 (>800)	800 (400)
P. aeruginosa PAO1	100 **(25)**	>800 (>800)	800 (400)	>800 (>800)	800 (400)
S. maltophilia ATCC 13637	200 (100)	>800 **(12.5/25)**	100 (50)	>800 (800)	>800 (>800)
S. maltophilia ATCC 12714	400 (400)	>800 **(50)**	200 (100)	>800 (800)	>800 (>800)
A. baumannii ATCC 19606	>800 (800)	>800 **(400)**	200 **(50)**	>800 **(100)**	>800 (>800)
B. cepacia ATCC 25416	>800 (800)	>800 (>800)	800 (800)	>800 (>800)	800 (800)

alen, alendronate sodium; tic, ticlopidine; amit, amitriptyline; nic, nicergoline; crb, carboplatin; PAβN, Phe-Arg-β-naphthylamide. [a] When a difference between the minimum inhibitory concentration (MIC) values of the non-antibiotic active substance and medicinal product with and without PAβN was observed, both minimum inhibitory concentrations (MICs) of the active substance and its medicinal product are presented; [b] At least 4-fold decrease in the MIC of non-antibiotic active substances and medicinal products in the presence of PAβN, when compared with the MIC values of non-antibiotics without PAβN, is indicated in boldface.