

## Reduction Algorithm

Given an SAT problem with  $n$  variables and a formula, turn all  $\bar{x}_i$  in the formula into a new variable  $y_i$  in a new formula, which is not negated. Also, add  $n$  new clauses  $(x_i \vee y_i)$  for all  $i \in \{1, 2, \dots, n\}$ , where  $y_i = \bar{x}_i$ . Run the imaginary Monotone SAT code on this new formula with  $n$  as  $k$ . The answer from this is the answer for SAT. For example:

$$(x_1 \vee x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2) \wedge (\bar{x}_3 \vee x_1) \wedge (\bar{x}_1 \vee x_3)$$

Will be reduced to

$$(x_1 \vee x_2 \vee x_3) \wedge (y_1 \vee y_2) \wedge (y_3 \vee x_1) \wedge (y_1 \vee x_3) \wedge (x_1 \vee y_1) \wedge (x_2 \vee y_2) \wedge (x_3 \vee y_3), \quad k = 3$$

This reduction algorithm is polynomial because you're just going through the entire formula and creating a new, non-negated variable for each negated variable (linear, actually). Transforming the output is constant time.

## Proof of Correctness

We need to prove both of the two following claims:

1. if the answer to  $\text{MonotoneSAT}(\text{formula}', k)$  is yes, then the answer to  $\text{SAT}(\text{formula})$  is also yes.
  2. if the answer to  $\text{MonotoneSAT}(\text{formula}', k)$  is no, then the answer to  $\text{SAT}(\text{formula})$  is also no.
- 
1. Suppose  $\text{MonotoneSAT}(\text{formula}', k)$  returns yes, then  $\text{formula}'$  is satisfiable with at most  $k$  variables set to true. We added  $k = n$  new clauses. So,  $\text{MonotoneSAT}(\text{formula}', k)$  returning yes means the  $k$  new clauses are all true, meaning at least one variable in each of those  $k$  disjunctions are true; and since we have the constraint of setting at most  $k$  variables to true, exactly one variable from each of these  $k$  new clauses is true. This satisfies that for any  $i$ ,  $x_i$  and  $\bar{x}_i$  cannot both be true in the original problem. The remaining clauses are essentially the original problem. Thus,  $\text{SAT}(\text{formula})$  also returns yes.
  2. We will prove the contrapositive of the 2<sup>nd</sup> claim: if the answer to  $\text{SAT}(\text{formula})$  is yes, then the answer to  $\text{MonotoneSAT}(\text{formula}', k)$  is also yes. Since there is a one-to-one correspondence from all the variables  $a \in \text{formula}$  to all variables  $b \in \text{formula}'$  (injection from  $\text{formula}$  variables to  $\text{formula}'$  variables), we copy over  $a$ 's truth value to  $b$ 's truth value: set all  $y_i = \neg x_i$ , and  $x_i = x_i$ . The  $k$  new clauses are all obviously true, and the remaining clauses are essentially the original problem, hence also true. Thus,  $\text{MonotoneSAT}(\text{formula}', k)$  also returns yes.