



DPNU-34
July, 1976

DICTIONARY
OF
ANGULAR DISTRIBUTIONS OF PARTICLES
FROM
 e^+e^- ANNIHILATION

Yasunari TOSA

Department of physics, Nagoya University, Nagoya 464, JAPAN

Abstract

This is the summary of works done by Y.S. Tsai and the author about the polar angle distributions. They are investigated in both all pair of combinations for scalar-, pseudoscalar-, vector-, axialvector-, and tensor particles, and one ~~ap~~rticle productions with a photon.

Recent experimental developments in e^+e^- colliding beams have opened up various exciting and fascinating facts: New hadron family ψ series,¹⁾ jets in inclusive hadron production,²⁾ a narrow state at 1865 MeV/c²,³⁾ anomalous $e\mu$ events,⁴⁾ and so on.

We shall investigate polar angle distributions for all pair of combinations of S(scalar), P(pseudoscalar), V(vector), A(axialvector), T(tensor), and ~~these~~ for one particle productions with a γ -emission, which are produced by e^+e^- annihilation. The calculation is carried out by adopting the effective covariant vertices.

The importance of the angular distribution is as follows: First in ψ series, the J^{PC} -determination of P_C and χ states will play a major role for the selection of models⁵⁾ and gaining an insight into the dynamics for constituents of new hadrons. Several works on angular distributions of $\psi' + \gamma\chi$ have already been done⁶⁾ using the non-relativistic approximation. But in some cases, e.g. ϕ^{*+} , the recoil effect cannot be neglected. We take it into account by making the vertices Lorentz invariant. Second, hadron production by e^+e^- annihilation has shown that the distribution of jets is $(1 + \cos^2\theta)$ where θ is the polar angle. It exists various possibilities for the interpretation of this distribution besides a pair of spin 1/2 particles. Y.S. Tsai⁷⁾ discussed them, but some cases were omitted. We include the cases done by Tsai for completeness of our study. And last, a new narrow state at 1865 MeV/c² must show $(1 + \cos^2\theta)$ distribution if it is D^0 accompanied by D^{*0} . This is the easiest method for the elimination of other possibilities, e.g., SS' , PP' , VV' , etc.

*) The prime denotes the second radial excited state of ϕ .

It should be noted that the azimuthal angle distribution for polarized beams can be derived from our results, because the general form can be written as⁸⁾

$$\frac{d\sigma}{d\Omega} = (\sigma_T + \sigma_L) \left(1 + \frac{\sigma_T - \sigma_L}{\sigma_T + \sigma_L} \cos^2\theta + \xi^2 \frac{\sigma_T - \sigma_L}{\sigma_T + \sigma_L} \sin^2\theta \cos 2\phi \right),$$

where $\theta(\phi)$ is the polar (azimuthal) angle and ξ is the polarization parameter.

The cross section is proportional to $L^{\mu\nu} M_{\mu\nu}$ where $L^{\mu\nu}(M_{\mu\nu})$ is the tensor formed by e^+e^- (final particles). The general form of $M_{\mu\nu}$ is

$$M_{\mu\nu} = \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \right) W_1 - \left(k_\mu - \frac{(kp)}{p^2} p_\mu \right) \left(k_\nu - \frac{(kp)}{p^2} p_\nu \right) \frac{W_2}{M^2}.$$

We denote the total momentum by p_μ and k_μ is the four momentum carried by final one particle. The angular distribution takes the form of

$$\left(2W_1 - k^2 \frac{W_2}{M^2} \right) + k^2 \frac{W_2}{M^2} \cos^2\theta,$$

where \vec{k} is the momentum of a final particle.

Now we are ready to open the dictionary.

Table I. Angular Distributions for $e^+e^- \rightarrow XY$

Mode	Vertex	Structure Functions	Angular Distribution ^{a)}	
				$ \vec{k} \gg M_X, M_Y$
$e^+e^- \rightarrow \bar{N}N$	$F_1 A^{\mu\gamma} \psi + F_2 F_{\mu\sigma} \bar{\psi} \sigma^{\mu\nu} \psi$	$W_1 = 2G_M^2 p^2$ $\vec{k}^2 \frac{W_2}{M^2} = 2(G_M^2 p^2 - 4m^2 G_E^2)$	$(sG_M^2 + 4m^2 G_E^2) + (sG_M^2 - 4m^2 G_E^2) \cos^2 \theta$	$1 + \cos^2 \theta$
$e^+e^- \rightarrow \bar{S}\bar{S}, SS'$	(minimal) + $F_{\mu\nu} \partial^\mu S \partial^\nu S'$	$W_1 = 0$	$1 - \cos^2 \theta$	$1 - \cos^2 \theta$
$e^+e^- \rightarrow SP$	This is forbidden.			
$e^+e^- \rightarrow SV$	$F_{\mu\nu} V^{\mu\nu} S$	$W_1 = (kp)^2, \frac{W_2}{M^2} = p^2$	$(1 + \frac{k^2}{2M_V^2}) + \frac{k^2}{2M_V^2} \cos^2 \theta$	$1 + \cos^2 \theta$
$e^+e^- \rightarrow SA$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} F^{\mu\nu} \partial^\lambda S A^\sigma$	$W_1 = (kp)^2 - k^2 p^2, \frac{W_2}{M^2} = p^2$	$1 + \cos^2 \theta$	$1 + \cos^2 \theta$
$e^+e^- \rightarrow ST$	$F_{\mu\nu} T^{\mu\lambda} \partial_\lambda \partial^\nu S$	$W_1 = \frac{1}{2} (kp)^2 (p \cdot q)$ $\frac{W_2}{M^2} = \frac{p^2}{2q^2} (kp)^2 - \frac{2}{3} \frac{(k \cdot q)^2}{q^4} p^4$	$\{7 + (3 + 4 \frac{s}{M_t^2}) \frac{k^2}{M_s^2}\} - \{1 + (-3 + 4 \frac{s}{M_t^2}) \frac{k^2}{M_s^2}\} \times \cos^2 \theta$	$1 - \cos^2 \theta$
$e^+e^- \rightarrow \bar{p}p_{pp'}$	the same as that of $e^+e^- \rightarrow \bar{S}\bar{S}, SS'$		$1 - \cos^2 \theta$	$1 - \cos^2 \theta$
$e^+e^- \rightarrow PV$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} \partial^\mu V^\lambda F^{\sigma\rho} p_\rho$	$W_1 = (kp)^2 - k^2 p^2, \frac{W_2}{M^2} = p^2$	$1 + \cos^2 \theta$	$1 + \cos^2 \theta$
$e^+e^- \rightarrow PA$	$F_{\mu\nu} A^{\mu\nu} p$	$W_1 = (kp)^2, \frac{W_2}{M^2} = p^2 (1 - \frac{p^2}{q^2})$	$\{1 + (1 + \frac{s}{M_A^2}) \frac{k^2}{2M_p^2}\} + (1 - \frac{s}{M_A^2}) \frac{k^2}{2M_p^2} \cos^2 \theta$	$1 - \cos^2 \theta$
$e^+e^- \rightarrow PT$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} T^{\mu\delta} F^{\nu\lambda} \partial_\delta \sigma p$	$W_1 = (kp)^2 - k^2 p^2, \frac{W_2}{M^2} = p^2$	$1 + \cos^2 \theta$	$1 + \cos^2 \theta$

Table I — Continued

$e^+e^- \rightarrow V\bar{V}$	$F_1(\text{minimal}) + F_2 F_{\mu\nu} V^{\mu\nu}$	$W_1 = F_1 \frac{k^2 + q^2}{2k^2 q^2} ((k \cdot q)^2 - k^2 q^2)$ $+ F_2 ((PI_q P) + (PI_k P))$ $\frac{W_2}{M^2} = F_1 (2 \frac{k^2 + q^2}{k^2 q^2} - 8)$ $+ F_2 P^2 \frac{1}{2k^2 q^2} [k^2 + q^2 - P^2]$	$\{F_2 + (F_1 + F_2) \frac{2M_V^2}{s} - 12F_1 \frac{M_V^4}{s^2}\}$ $- \{F_2 - (F_1 + F_2) \frac{2M_V^2}{s} + 12F_1 \frac{M_V^4}{s^2}\} \cos^2 \theta$	$1 - \cos^2 \theta$ $\left(\begin{array}{l} \text{If } F_2 = 0 \\ 1 + \cos^2 \theta \end{array} \right)$
$e^+e^- \rightarrow VV'$	$F_{\mu\nu} V^{\mu\nu}, V'$			$1 - \cos^2 \theta$
$e^+e^- \rightarrow VA$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} V^{\mu\nu} V'^{\lambda\sigma}$	$W_1 = \frac{(kp)^2}{k^2} + \frac{(qp)^2}{q^2}$ $\frac{W_2}{M^2} = P^2 \frac{k^2 + q^2}{k^2 q^2}$	$(1 + \frac{k^2}{4} \frac{M_V^2 + M_A^2}{M_V^2 M_A^2}) + \frac{k^2}{4} \frac{M_t^2 + M_A^2}{M_V^2 M_A^2} \cos^2 \theta$	$1 + \cos^2 \theta$
$e^+e^- \rightarrow VT^{(b)}$	$f^{(1)} \partial_\mu V_\nu F_\lambda^{\mu\nu} T^\lambda{}_\nu$ $+ f^{(2)} \partial^\mu V_\nu T^\nu{}_\lambda T^\lambda{}_\mu$	***	$[(\frac{5}{3} + \frac{s}{2M_V^2} \frac{M_t^2}{M_t^2}) k^2 + \frac{k^2}{2M_V^2} (\frac{5}{3} + \frac{11}{6} \frac{s}{M_t^2})]$ $+ \frac{k^2}{2M_V^2} (\frac{5}{3} - \frac{11}{6} \frac{s}{M_t^2}) \cos^2 \theta$	$1 - \cos^2 \theta$
$e^+e^- \rightarrow A\bar{A}$ AA'	the same as that of $e^+e^- \rightarrow V\bar{V}$			$1 - \cos^2 \theta$
$e^+e^- \rightarrow AT$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} F^{\mu\nu} T^{\lambda\sigma} \delta_\delta A^\sigma$	$W_1 = (PI_q P) [\frac{7}{6} P^2 + \frac{2}{3} (PI_q P)]$ $\frac{W_2}{M^2} = \frac{P^2}{q^2} [-\frac{1}{2} P^2 + \frac{2}{3} \frac{(qp)^2}{q^2}]$	$(13 + 4 \frac{k^2}{M_t^2}) + (1 + 4 \frac{k^2}{M_t^2}) \cos^2 \theta$	$1 + \cos^2 \theta$

a) $s = p^2$

b) Reference 9.

Table II. Angular Distributions for $e^+e^- \rightarrow V \rightarrow \gamma X$

Mode	Vertex	Angular Distribution
$e^+e^- \rightarrow V \rightarrow S\gamma$	$F_{\mu\nu} V^{\mu\nu} S$	$1 + \cos^2\theta$
$e^+e^- \rightarrow V \rightarrow P\gamma$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} \partial^\mu V^\nu F^{\lambda\sigma} P$	$1 + \cos^2\theta$
$e^+e^- \rightarrow V \rightarrow V'\gamma$	$F_{\mu\nu} V^\mu V'^\nu$	$(3 + 2 \frac{\sqrt{s}}{2} \vec{k}) - (1 - 2 \frac{\sqrt{s}}{2} \vec{k}) \cos^2\theta$
$e^+e^- \rightarrow V \rightarrow A\gamma$	$\frac{1}{2} \epsilon_{\mu\nu\lambda\sigma} F^{\mu\nu} V^\lambda A^\sigma$	$(3 + 2 \frac{\sqrt{s}}{2} \vec{k}) - (1 - 2 \frac{\sqrt{s}}{2} \vec{k}) \cos^2\theta$
$e^+e^- \rightarrow V \rightarrow T\gamma^a)$	$f^{(1)} \partial_\mu V^\nu F^\mu{}_\sigma T^\sigma{}_\nu$ $+ f^{(2)} \partial_\mu V^\nu F^\mu{}_\nu T^\sigma{}_\sigma$	$[(13 + 6 \frac{k^2}{M_t^2}) - 4g \frac{ \vec{k} }{\sqrt{s}} (4 + 10 \frac{\sqrt{s}}{2} \vec{k}) + 2g^2 \frac{k^2}{M_t^2} (5 + 4 \frac{\sqrt{s}}{2} \vec{k})]$ $+ [(1 - 6 \frac{k^2}{M_t^2}) + 4g \frac{ \vec{k} }{\sqrt{s}} (2 + \frac{\sqrt{s}}{2} \vec{k}) - 2g^2 \frac{k^2}{M_t^2} (1 - 4 \frac{\sqrt{s}}{2} \vec{k})] \cos^2\theta$ <p>where</p> $g = f^{(2)} / f^{(1)}$

a) Reference 9.

In conclusion, we have found that the angular distributions for all pair of combinations of S, V, T, P, A produced by e^+e^- annihilation can be classified in two groups according to their behaviors in the relativistic limit. One group, which shows $(1 + \cos^2\theta)$ distribution, consists of SV, SA, PV, PT, VA, AT, while the other is composed of SS', ST, PP', PA, VV', VT, AA', which are produced with $(1 - \cos^2\theta)$. The former may also be relevant to the observed jet structure.

References.

- 1) Summary at the summer of 1975 is found in Proceedings of the 1975 International Symposium on Lepton and Photon Interactions, Stanford University (1975).
- 2) R.F. Schwitters et al., Phys. Rev. Lett. 20, 1320 (1975); G. Hanson et al., Phys. Rev. Lett. 24, 1609 (1975); G. Hanson, in Proceedings of Summer Institute on Particle Physics, SLAC Report No. 191 (November, 1975).
- 3) G. Goldhaber et al., LBL-5309/SLAC-1762 (June, 1976).
- 4) M.L. Perl et al., Phys. Rev. Lett. 35, 1489 (1975).
- 5) Some models require that P_c and χ states are all natural parity particles, in contrast to charmoniums. See Y.Tosa, O.Shito, T. Kaji, and J. Iizuka, DPNU-28 (November, 1975), and N.P. Chang and C.A. Nelson, Phys. Rev. D13, 1345 (1976).
- 6) E. Eichten et al., Phys. Rev. Lett. 34, 369 (1975); L.S. Brown and R.N. Cahn, Phys. Rev. D13, 1195 (1976); G. Karl, S. Meshkov, and J.L. Rosner, Phys. Rev. D13, 1203 (1976).
- 7) Y.S. Tsai, Phys. Rev. D12, 3533 (1975).
- 8) N. Cabibbo and R. Gatto, Phys. Rev. 124, 1577 (1961); F. Gilman, in Proceedings of Summer Institute on Particle Physics, SLAC Report No. 191 (November, 1975).
- 9) Y. Tosa, O. Shito, T. Kaji, and J. Iizuka, DPNU-28 (November, 1975), and Y. Tosa, Preprint (July, 1976).

