Structural SV(outliers, t): BVAR(Minnesota(structural, eb), SV(outliers,t), p=2)

```
>>> print(fit.plan.describe())
phase     step        block            writes
equation  equation    EQ_MEAN(0)       eq.mean, eq.struct.a
                                       (δ_i, a_i) | λ_eff,i joint Gaussian (fused contemporaneous draw)
equation  residual    EQ_MEAN(0)       —
                                       e_i = ỹ_i + Σ_{j<i} a_ij ỹ_j − g_i from the new draws
equation  vol         EQ_VOL(2)        eq.vol
                                       VolBlock.sample_eq: o → p → ν → τ, then s → h → θ_SV (atomic)
post      (none)


read sets (Step.reads of the same plan):
  equation    reads eq.coef, eq.mean, eq.struct, eq.vol, glob
  residual    reads eq.mean, eq.struct, glob.mean
  vol         reads eq.vol, glob.vol
```

Reduced-form SV: BVAR(Minnesota(reduced, eb), SV, p=2)

```
>>> print(fit.plan.describe())
phase     step        block            writes
pre       cccm        G_CCCM(16)       glob.mean.Pi
equation  contemp     EQ_CONTEMP(1)    eq.struct.a
                                       a_i | g_i, λ_eff,i: regress ỹ_i − g_i on −ỹ_{<i}
equation  residual    EQ_MEAN(0)       —
                                       e_i = ỹ_i + Σ_{j<i} a_ij ỹ_j − g_i from the new draws
equation  vol         EQ_VOL(2)        eq.vol
                                       VolBlock.sample_eq: o → p → ν → τ, then s → h → θ_SV (atomic)
post      (none)


read sets (Step.reads of the same plan):
  cccm        reads eq.struct.a, eq.vol, glob.mean.Pi
  contemp     reads eq.coef, eq.mean, eq.struct, eq.vol, glob
  residual    reads eq.mean, eq.struct, glob.mean
  vol         reads eq.vol, glob.vol
```