

Cyclic Higgs bundles and minimal surfaces in pseudo-hyperbolic spaces

(Joint Seminar on Teich and Related Topics 2022-9-19)

Xin Nie (Yau Center of Southeast University)

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;
- ▶ $\text{Isom}(\mathbb{H}^{p,q}) = \text{O}(p, q+1)$;

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;
- ▶ $\text{Isom}(\mathbb{H}^{p,q}) = \text{O}(p, q+1)$; not connected, subgroups:

$$\underbrace{\text{SO}(p, q+1)}_{\text{orientation-preserving isom.}} \stackrel{1:2}{\subset} \text{O}(p, q+1)$$

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;
- ▶ $\text{Isom}(\mathbb{H}^{p,q}) = \text{O}(p, q+1)$; not connected, subgroups:

$$\underbrace{\text{SO}_0(p, q+1)}_{\substack{\text{orient.-preserving isom.} \\ \text{acting trivially on homology}}} \stackrel{1:2}{\subset} \underbrace{\text{SO}(p, q+1)}_{\text{orientation-preserving isom.}} \stackrel{1:2}{\subset} \text{O}(p, q+1)$$

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;
- ▶ $\text{Isom}(\mathbb{H}^{p,q}) = \text{O}(p, q+1)$; not connected, subgroups:

$$\underbrace{\text{SO}_0(p, q+1)}_{\substack{\text{orient.-preserving isom.} \\ \text{acting trivially on homology}}} \stackrel{1:2}{\subset} \underbrace{\text{SO}(p, q+1)}_{\text{orientation-preserving isom.}} \stackrel{1:2}{\subset} \text{O}(p, q+1)$$

- ▶ $\mathbf{S}_{p,q} := \{\text{timelike totally geodesic } q\text{-spheres}\}$

Pseudo-hyperbolic spaces

The pseudo-hyperbolic space $\mathbb{H}^{p,q}$ (hyperboloid model):

$$\mathbb{H}^{p,q} = \{x \in \mathbb{R}^{p,q+1} \mid \langle x, x \rangle = -1\},$$

with the metric restricted from $\mathbb{R}^{p,q+1}$.

- ▶ $\mathbb{H}^{p,q}$ is a pseudo-Riemannian manifold of constant sectional curvature -1 ;
- ▶ diffeomorphic to $\mathbb{B}^p \times \mathbb{S}^q$;
- ▶ $\text{Isom}(\mathbb{H}^{p,q}) = \text{O}(p, q+1)$; **not connected, subgroups:**

$$\underbrace{\text{SO}_0(p, q+1)}_{\substack{\text{orient.-preserving isom.} \\ \text{acting trivially on homology}}} \stackrel{1:2}{\subset} \underbrace{\text{SO}(p, q+1)}_{\text{orientation-preserving isom.}} \stackrel{1:2}{\subset} \text{O}(p, q+1)$$

- ▶ $\mathbf{S}_{p,q} := \{\text{timelike totally geodesic } q\text{-spheres}\}$
 $= \text{SO}_0(p, q+1) / (\text{SO}(p) \times \text{SO}(q+1)) \cong \mathbf{S}_{q+1, p-1}$
(Riemannian symmetric space of nonpositive curvature)

Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.

Pseudo-hyperbolic spaces


Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.

Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:

Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (**adS**) space.


Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:



Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.


Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:



Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.

Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:




► $\mathrm{SO}_0(2, 2) \cong \mathrm{PSL}(2, \mathbb{R}) \times \mathrm{PSL}(2, \mathbb{R})$

Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.

Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:




- ▶ $\mathrm{SO}_0(2, 2) \cong \mathrm{PSL}(2, \mathbb{R}) \times \mathrm{PSL}(2, \mathbb{R})$
- ▶ $\mathbf{S}_{2,1} = \{\text{timelike closed geodesics}\}$
 $= \mathrm{SO}_0(2, 2) / (\mathrm{SO}(2) \times \mathrm{SO}(2)) = \mathbb{H}^2 \times \mathbb{H}^2.$

Pseudo-hyperbolic spaces

Example. $\mathbb{H}^{2,1}$ = anti-de Sitter (adS) space.

Projectivized picture in the affine plane $\mathbb{R}^{2,1} \cong \{x_4 = 1\} \subset \mathbb{R}^{2,2}$:



- ▶ $\mathrm{SO}_0(2, 2) \cong \mathrm{PSL}(2, \mathbb{R}) \times \mathrm{PSL}(2, \mathbb{R})$
- ▶ $\mathbf{S}_{2,1} = \{\text{timelike closed geodesics}\}$
 $= \mathrm{SO}_0(2, 2) / (\mathrm{SO}(2) \times \mathrm{SO}(2)) = \mathbb{H}^2 \times \mathbb{H}^2.$

All the other $\mathrm{SO}_0(p, q + 1)$'s ($p, q \geq 2$) are simple Lie groups.

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.
 $\mathbf{II} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.

$\mathbf{II} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

= submanifold whose volume is stationary w.r.t. compactly supported variations

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.

$\mathbf{II} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

= submanifold whose volume is stationary w.r.t. compactly supported variations (critical point for the volume functional).

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.

$\mathbf{II} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

= submanifold whose volume is stationary w.r.t. compactly supported variations (critical point for the volume functional).

Maximality Theorem

Let Σ be a **spacelike** minimal submanifold with **timelike** normal bundle in a pseudo-Riemannian manifold M of **negative** sectional curvature.

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.

$\mathbf{II} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

= submanifold whose volume is stationary w.r.t. compactly supported variations (critical point for the volume functional).

Maximality Theorem

Let Σ be a **spacelike** minimal submanifold with **timelike** normal bundle in a pseudo-Riemannian manifold M of **negative** sectional curvature. Then any nonzero, compactly supported, normal deformation of Σ locally decreases the volume.

Maximal surfaces in $\mathbb{H}^{2,q}$

Minimal submanifold = submanifold Σ whose 2nd f.f.

$\mathbb{I} \in C^\infty(\text{Sym}^2 T^* \Sigma \otimes \mathbb{N})$ has zero trace (w.r.t. 1st f.f.)

= submanifold whose volume is stationary w.r.t. compactly supported variations (critical point for the volume functional).

Maximality Theorem

Let Σ be a **spacelike** minimal submanifold with **timelike** normal bundle in a pseudo-Riemannian manifold M of **negative** sectional curvature. Then any nonzero, compactly supported, normal deformation of Σ locally decreases the volume.

\Rightarrow Spacelike minimal surfaces in $\mathbb{H}^{2,q}$ are called **maximal surfaces**.

Proof. 2nd variation formula of volume for minimal submanifolds

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
($\xi \in C^\infty(\Sigma, \mathbb{N})$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{vol}(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\langle \nabla_{\bullet}^{\mathbb{N}} \xi, \nabla_{\bullet}^{\mathbb{N}} \xi \rangle - \langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle - \langle R_M(\bullet, \xi) \xi, \bullet \rangle \right) d\text{vol}_g$$

- \bullet = slot for tangent vector,
- g = 1st fundamental form,
- $\nabla^{\mathbb{N}}$ = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
($\xi \in C^\infty(\Sigma, \mathbb{N})$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{vol}(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\underbrace{\langle \overbrace{\nabla_{\bullet}^{\mathbb{N}} \xi}^{\text{normal vector}}, \nabla_{\bullet}^{\mathbb{N}} \xi \rangle}_{\leq 0} - \langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle - \langle R_M(\bullet, \xi) \xi, \bullet \rangle \right) d\text{vol}_g$$

- \bullet = slot for tangent vector,
- g = 1st fundamental form,
- $\nabla^{\mathbb{N}}$ = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
($\xi \in C^\infty(\Sigma, N)$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{vol}(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\underbrace{\langle \nabla_{\bullet}^N \xi, \nabla_{\bullet}^N \xi \rangle}_{\leq 0} - \underbrace{\langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle}_{\geq 0} - \langle R_M(\bullet, \xi)\xi, \bullet \rangle \right) d\text{vol}_g$$

- \bullet = slot for tangent vector,
- g = 1st fundamental form,
- ∇^N = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
($\xi \in C^\infty(\Sigma, N)$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} vol(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\underbrace{\langle \nabla_{\bullet}^N \xi, \nabla_{\bullet}^N \xi \rangle}_{\leq 0} - \underbrace{\langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle}_{\geq 0} - \underbrace{\langle R_M(\bullet, \xi) \xi, \bullet \rangle}_{\geq 0, \text{ strict on } \{\xi \neq 0\}} \right) dvol_g$$

- \bullet = slot for tangent vector,
- g = 1st fundamental form,
- ∇^N = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
 ($\xi \in C^\infty(\Sigma, N)$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} vol(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\underbrace{\langle \nabla_{\bullet}^N \xi, \nabla_{\bullet}^N \xi \rangle}_{\leq 0} - \underbrace{\langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle}_{\geq 0} - \underbrace{\langle R_M(\bullet, \xi) \xi, \bullet \rangle}_{\geq 0, \text{ strict on } \{\xi \neq 0\}} \right) dvol_g$$

- \bullet = slot for tangent vector,
- g = 1st fundamental form,
- ∇^N = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

$$\text{because } \underbrace{K(v \wedge \xi)}_{-} = \frac{\langle R_M(v, \xi) \xi, v \rangle}{\underbrace{\langle v, v \rangle}_{+} \underbrace{\langle \xi, \xi \rangle}_{-} - \cancel{\langle v, \xi \rangle^2}}$$

Maximal surfaces in $\mathbb{H}^{2,q}$

Proof. 2nd variation formula of volume for minimal submanifolds
 ($\xi \in C^\infty(\Sigma, N)$ variational vector field):

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{vol}(\Sigma_t) = \int_{\Sigma} \text{Tr}_g \left(\underbrace{\langle \nabla_{\bullet}^N \xi, \nabla_{\bullet}^N \xi \rangle}_{\leq 0} - \underbrace{\langle A_{\xi}(\bullet), A_{\xi}(\bullet) \rangle}_{\geq 0} - \underbrace{\langle R_M(\bullet, \xi) \xi, \bullet \rangle}_{\geq 0, \text{ strict on } \{\xi \neq 0\}} \right) d\text{vol}_g$$

< 0

- = slot for tangent vector,
- g = 1st fundamental form,
- ∇^N = normal connection,
- A_{ξ} = shape operator associated to ξ ,
- R_M = curvature tensor of M .

because $\underbrace{K(v \wedge \xi)}_{-} = \frac{\langle R_M(v, \xi) \xi, v \rangle}{\underbrace{\langle v, v \rangle}_{+} \underbrace{\langle \xi, \xi \rangle}_{-} - \cancel{\langle v, \xi \rangle^2}}$




Maximal surfaces in $\mathbb{H}^{2,q}$

Spacelike surface $\Sigma \subset \mathbb{H}^{2,q} \rightsquigarrow$ **Gauss map** $\psi : \Sigma \rightarrow \mathbf{S}_{2,q}$


Maximal surfaces in $\mathbb{H}^{2,q}$

Spacelike surface $\Sigma \subset \mathbb{H}^{2,q} \rightsquigarrow$ **Gauss map** $\psi : \Sigma \rightarrow \mathbf{S}_{2,q}$



Maximal surfaces in $\mathbb{H}^{2,q}$

Spacelike surface $\Sigma \subset \mathbb{H}^{2,q} \rightsquigarrow$ **Gauss map** $\psi : \Sigma \rightarrow \mathbf{S}_{2,q}$




Theorem (analogue for Gauss map of minimal surfaces in \mathbb{R}^3)

If Σ is maximal, then ψ is a conformal minimal immersion.

Maximal surfaces in $\mathbb{H}^{2,q}$

Spacelike surface $\Sigma \subset \mathbb{H}^{2,q} \rightsquigarrow$ **Gauss map** $\psi : \Sigma \rightarrow \mathbf{S}_{2,q}$



Theorem (analogue for Gauss map of minimal surfaces in \mathbb{R}^3)


If Σ is maximal, then ψ is a conformal minimal immersion.

Example. Maximal surface in $\mathbb{H}^{2,1}$

\rightsquigarrow conformal minimal immersion $\psi : \Sigma \rightarrow \mathbf{S}_{2,1} = \mathbb{H}^2 \times \mathbb{H}^2$

Maximal surfaces in $\mathbb{H}^{2,q}$

Spacelike surface $\Sigma \subset \mathbb{H}^{2,q} \rightsquigarrow$ **Gauss map** $\psi : \Sigma \rightarrow \mathbf{S}_{2,q}$



Theorem (analogue for Gauss map of minimal surfaces in \mathbb{R}^3)

If Σ is maximal, then ψ is a conformal minimal immersion.

Example. Maximal surface in $\mathbb{H}^{2,1}$

\rightsquigarrow conformal minimal immersion $\psi : \Sigma \rightarrow \mathbf{S}_{2,1} = \mathbb{H}^2 \times \mathbb{H}^2$

\rightsquigarrow pair of harmonic maps $\psi_+, \psi_- : \Sigma \rightarrow \mathbb{H}^2$ with opposite Hopf differentials.

Applications in (higher) Teichmüller theory

- ▶ (Mess 2007, Bonsante-Schlenker 2010) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,1}$ (“globally hyperbolic spacetimes”, analogue of quasi-Fuchsian convex cores) are useful for studying pairs of Fuchsian representations $\pi_1(S_g) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ (e.g. new proof and extensions of Thurston’s Earthquake Theorem).

Applications in (higher) Teichmüller theory

- ▶ (Mess 2007, Bonsante-Schlenker 2010) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,1}$ (“globally hyperbolic spacetimes”, analogue of quasi-Fuchsian convex cores) are useful for studying pairs of Fuchsian representations $\pi_1(S_g) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ (e.g. new proof and extensions of Thurston’s Earthquake Theorem).
- ▶ (Collier-Tholozan-Toulisse 2019) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,q}$ are useful for studying maximal representations $\pi_1(S_g) \rightarrow \mathrm{SO}_0(2, q + 1)$.

Applications in (higher) Teichmüller theory

- ▶ (Mess 2007, Bonsante-Schlenker 2010) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,1}$ (“globally hyperbolic spacetimes”, analogue of quasi-Fuchsian convex cores) are useful for studying pairs of Fuchsian representations $\pi_1(S_g) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ (e.g. new proof and extensions of Thurston’s Earthquake Theorem).
- ▶ (Collier-Tholozan-Toulisse 2019) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,q}$ are useful for studying maximal representations $\pi_1(S_g) \rightarrow \mathrm{SO}_0(2, q+1)$.

What about minimal surfaces in $\mathbb{H}^{p,q}, p > 2$?

Applications in (higher) Teichmüller theory

- ▶ (Mess 2007, Bonsante-Schlenker 2010) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,1}$ (“globally hyperbolic spacetimes”, analogue of quasi-Fuchsian convex cores) are useful for studying pairs of Fuchsian representations $\pi_1(S_g) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ (e.g. new proof and extensions of Thurston’s Earthquake Theorem).
- ▶ (Collier-Tholozan-Toulisse 2019) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,q}$ are useful for studying maximal representations $\pi_1(S_g) \rightarrow \mathrm{SO}_0(2, q+1)$.

What about minimal surfaces in $\mathbb{H}^{p,q}, p > 2$?

- ▶ Not maximal (critical point of area functional, but not local maximum/minimum);

Applications in (higher) Teichmüller theory

- ▶ (Mess 2007, Bonsante-Schlenker 2010) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,1}$ (“globally hyperbolic spacetimes”, analogue of quasi-Fuchsian convex cores) are useful for studying pairs of Fuchsian representations $\pi_1(S_g) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ (e.g. new proof and extensions of Thurston’s Earthquake Theorem).
- ▶ (Collier-Tholozan-Toulisse 2019) Closed maximal surfaces in some partial quotients of $\mathbb{H}^{2,q}$ are useful for studying maximal representations $\pi_1(S_g) \rightarrow \mathrm{SO}_0(2, q+1)$.

What about minimal surfaces in $\mathbb{H}^{p,q}$, $p > 2$?

- ▶ Not maximal (critical point of area functional, but not local maximum/minimum);
- ▶ No obvious notion of Gauss map to $\mathbf{S}_{p,q}$.

Summary of results

- Introduce a special type of spacelike minimal surfaces Σ (“*A-surfaces*”) in $\mathbb{H}^{n,n}$ with n even or $\mathbb{H}^{n+1,n-1}$ with n odd, related to cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles.

Summary of results

- ▶ Introduce a special type of spacelike minimal surfaces Σ (“*A-surfaces*”) in $\mathbb{H}^{n,n}$ with n even or $\mathbb{H}^{n+1,n-1}$ with n odd, related to cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles.
- ▶ Have a good notion of Gauss map $\Sigma \rightarrow \mathbf{S}_{n,n}$ or $\mathbf{S}_{n+1,n-1}$ (recovers the minimal mapping from the Higgs bundle).

Summary of results

- ▶ Introduce a special type of spacelike minimal surfaces Σ (“*A-surfaces*”) in $\mathbb{H}^{n,n}$ with n even or $\mathbb{H}^{n+1,n-1}$ with n odd, related to cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles.
- ▶ Have a good notion of Gauss map $\Sigma \rightarrow \mathbf{S}_{n,n}$ or $\mathbf{S}_{n+1,n-1}$ (recovers the minimal mapping from the Higgs bundle).
- ▶ *Key Theorem.* Normal bundle of Σ has orthogonal splitting $N = N^+ \oplus N^-$ (spacelike/timelike parts). Under some conditions, any compactly supported deformation of Σ along N^+ (resp. N^-) increases (resp. decreases) the area.

Summary of results

- ▶ Introduce a special type of spacelike minimal surfaces Σ (“*A-surfaces*”) in $\mathbb{H}^{n,n}$ with n even or $\mathbb{H}^{n+1,n-1}$ with n odd, related to cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles.
- ▶ Have a good notion of Gauss map $\Sigma \rightarrow \mathbf{S}_{n,n}$ or $\mathbf{S}_{n+1,n-1}$ (recovers the minimal mapping from the Higgs bundle).
- ▶ *Key Theorem.* Normal bundle of Σ has orthogonal splitting $N = N^+ \oplus N^-$ (spacelike/timelike parts). Under some conditions, any compactly supported deformation of Σ along N^+ (resp. N^-) increases (resp. decreases) the area.
- ▶ *Corollary.* Infinitesimally rigid.

Summary of results

- ▶ Introduce a special type of spacelike minimal surfaces Σ (“*A-surfaces*”) in $\mathbb{H}^{n,n}$ with n even or $\mathbb{H}^{n+1,n-1}$ with n odd, related to cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles.
- ▶ Have a good notion of Gauss map $\Sigma \rightarrow \mathbf{S}_{n,n}$ or $\mathbf{S}_{n+1,n-1}$ (recovers the minimal mapping from the Higgs bundle).
- ▶ *Key Theorem.* Normal bundle of Σ has orthogonal splitting $N = N^+ \oplus N^-$ (spacelike/timelike parts). Under some conditions, any compactly supported deformation of Σ along N^+ (resp. N^-) increases (resp. decreases) the area.
- ▶ *Corollary.* Infinitesimally rigid.
- ▶ New proof of Labourie Immersion Theorem (\Rightarrow Labourie Conjecture for rank 2 groups) for $\mathrm{SO}_0(n, n+1)$ & generalization to Collier’s components.

A(ternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t ,

A(Iternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t , then $\exists!$ orthonormal moving frame $(e_1(t), \dots, e_n(t))$ such that

- ▶ $e_1 = \gamma'(t)/\|\gamma'(t)\|$ is the unit tangent;

A(Iternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t , then $\exists!$ orthonormal moving frame $(e_1(t), \dots, e_n(t))$ such that

► $e_1 = \gamma'(t)/\|\gamma'(t)\|$ is the unit tangent;


► $(e'_1(t), \dots, e'_n(t)) = (e_1(t), \dots, e_n(t)) \begin{pmatrix} 0 & -\kappa_2 & & & \\ \kappa_2 & 0 & -\kappa_3 & & \\ & \kappa_3 & 0 & \ddots & \\ & & \ddots & \ddots & -\kappa_n \\ & & & \kappa_n & 0 \end{pmatrix}.$

A(Iternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t , then $\exists!$ orthonormal moving frame $(e_1(t), \dots, e_n(t))$ such that

► $e_1 = \gamma'(t)/\|\gamma'(t)\|$ is the unit tangent;

► $(e'_1(t), \dots, e'_n(t)) = (e_1(t), \dots, e_n(t)) \begin{pmatrix} 0 & -\kappa_2 & & & \\ \kappa_2 & 0 & -\kappa_3 & & \\ & \kappa_3 & 0 & \ddots & \\ & & \ddots & \ddots & -\kappa_n \\ & & & \kappa_n & 0 \end{pmatrix}.$




A(Iternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t , then $\exists!$ orthonormal moving frame $(e_1(t), \dots, e_n(t))$ such that

► $e_1 = \gamma'(t)/\|\gamma'(t)\|$ is the unit tangent;

► $(e'_1(t), \dots, e'_n(t)) = (e_1(t), \dots, e_n(t)) \begin{pmatrix} 0 & -\kappa_2 & & & \\ \kappa_2 & 0 & -\kappa_3 & & \\ & \kappa_3 & 0 & \ddots & \\ & & \ddots & \ddots & -\kappa_n \\ & & & \kappa_n & 0 \end{pmatrix}.$




$\kappa_2, \dots, \kappa_n$ are complete intrinsic invariants of the curve
($n = 3$: curvature & torsion)

A(lternating)-surfaces

Frenet frame: Given curve $t \mapsto \gamma(t) \in \mathbb{E}^n$, if $\gamma'(t), \gamma''(t), \dots, \gamma^{(n)}(t)$ are linearly independent for all t , then $\exists!$ orthonormal moving frame $(e_1(t), \dots, e_n(t))$ such that

► $e_1 = \gamma'(t)/\|\gamma'(t)\|$ is the unit tangent;

► $(e'_1(t), \dots, e'_n(t)) = (e_1(t), \dots, e_n(t)) \begin{pmatrix} 0 & -\kappa_2 & & & \\ \kappa_2 & 0 & -\kappa_3 & & \\ & \kappa_3 & 0 & \ddots & \\ & & \ddots & \ddots & -\kappa_n \\ & & & \kappa_n & 0 \end{pmatrix}.$



$\kappa_2, \dots, \kappa_n$ are complete intrinsic invariants of the curve
($n = 3$: curvature & torsion)

Alternating-surfaces

Theorem (Chern 1970)

A(lternating)-surfaces

Theorem (Chern 1970)

For any minimal 2-sphere $S \subset \mathbb{S}^{2n}$ not in any \mathbb{S}^{2n-1} ,

A(Iternating)-surfaces

Theorem (Chern 1970)

For any minimal 2-sphere $S \subset \mathbb{S}^{2n}$ not in any \mathbb{S}^{2n-1} , $\exists!$ orthogonal splitting $\mathbb{T}\mathbb{S}^{2n}|_S = L_1 \oplus \cdots \oplus L_n$ s.t.

(a) every L_i has rank 2; $L_1 = \mathbb{T}S$;

A(lternating)-surfaces

Theorem (Chern 1970)

For any minimal 2-sphere $S \subset \mathbb{S}^{2n}$ not in any \mathbb{S}^{2n-1} , $\exists!$ orthogonal splitting $\mathbb{T}\mathbb{S}^{2n}|_S = L_1 \oplus \cdots \oplus L_n$ s.t.

- (a) every L_i has rank 2; $L_1 = \mathbb{T}S$;
- (b) the Levi-Civita connection ∇ on $\mathbb{T}\mathbb{S}^{2n}|_S$ decomposes as

$$\nabla = \begin{pmatrix} \nabla_1 & -\alpha_2^* & & & \\ \alpha_2 & \nabla_2 & -\alpha_3^* & & \\ & \alpha_3 & \nabla_3 & \ddots & \\ & & \ddots & \ddots & -\alpha_n^* \\ & & & \alpha_n & \nabla_n \end{pmatrix};$$

Alternating-surfaces

Theorem (Chern 1970)

For any minimal 2-sphere $S \subset \mathbb{S}^{2n}$ not in any \mathbb{S}^{2n-1} , $\exists!$ orthogonal splitting $T\mathbb{S}^{2n}|_S = L_1 \oplus \cdots \oplus L_n$ s.t.

- (a) every L_i has rank 2; $L_1 = TS$;
- (b) the Levi-Civita connection ∇ on $T\mathbb{S}^{2n}|_S$ decomposes as

$$\nabla = \begin{pmatrix} \nabla_1 & -\alpha_2^* & & & \\ \alpha_2 & \nabla_2 & -\alpha_3^* & & \\ & \alpha_3 & \nabla_3 & \ddots & \\ & & \ddots & \ddots & -\alpha_n^* \\ & & & \alpha_n & \nabla_n \end{pmatrix};$$

- (c) each $\alpha_i \in \Omega^1(S, \text{Hom}(L_{i-1}, L_i))$ is **conformal** (i.e. any $\alpha_i(x)v : L_{i-1}|_x \rightarrow L_i|_x$ is a conformal linear map of Euclidean planes).

A(lternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (a famous idea of Hopf);

A(lternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (**a famous idea of Hopf**); find L_3, L_4, \dots and show the conformality of $\alpha_3, \alpha_4, \dots$ inductively in the same way. \square

A(lternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (**a famous idea of Hopf**); find L_3, L_4, \dots and show the conformality of $\alpha_3, \alpha_4, \dots$ inductively in the same way. \square

For surfaces not homeomorphic to \mathbb{S}^2 :

A(lternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (**a famous idea of Hopf**); find L_3, L_4, \dots and show the conformality of $\alpha_3, \alpha_4, \dots$ inductively in the same way. \square

For surfaces not homeomorphic to \mathbb{S}^2 :

Definition

A **superminimal surface** in a Riemannian manifold M^{2n} is a minimal surface Σ with an orthogonal splitting $TM|_{\Sigma} = L_1 \oplus \dots \oplus L_n$ satisfying conditions (a), (b), (c).

A(lternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (**a famous idea of Hopf**); find L_3, L_4, \dots and show the conformality of $\alpha_3, \alpha_4, \dots$ inductively in the same way. \square

For surfaces not homeomorphic to \mathbb{S}^2 :

Definition

A **superminimal surface** in a Riemannian manifold M^{2n} is a minimal surface Σ with an orthogonal splitting $TM|_{\Sigma} = L_1 \oplus \dots \oplus L_n$ satisfying conditions (a), (b), (c).

Cond.(c) relaxed s.t. only $\alpha_2, \dots, \alpha_{n-1}$ are assumed conformal (**no assumption on α_n**) \Rightarrow **quasi-superminimal surface**.

A(Iternating)-surfaces

Idea of proof. Find L_2 as the image of the 2nd fundamental form, use the fact that **any holomorphic k -differential on \mathbb{CP}^1 vanishes** to show that α_2 is conformal (**a famous idea of Hopf**); find L_3, L_4, \dots and show the conformality of $\alpha_3, \alpha_4, \dots$ inductively in the same way. \square

For surfaces not homeomorphic to \mathbb{S}^2 :

Definition

A **superminimal surface** in a Riemannian manifold M^{2n} is a minimal surface Σ with an orthogonal splitting $TM|_{\Sigma} = L_1 \oplus \dots \oplus L_n$ satisfying conditions (a), (b), (c).

Cond.(c) relaxed s.t. only $\alpha_2, \dots, \alpha_{n-1}$ are assumed conformal (**no assumption on α_n**) \Rightarrow **quasi-superminimal surface**.

Still makes sense for *pseudo-Riemannian* M if every L_i is (**positive or negative**) definite.

A(lternating)-surfaces


Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).

A(lternating)-surfaces

Definition


When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).



A(lternating)-surfaces

Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).




$\Rightarrow (p, q) = (n, n)$ (resp. $(n + 1, n - 1)$) when n is even (resp. odd).

A(lternating)-surfaces

Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).




$\Rightarrow (p, q) = (n, n)$ (resp. $(n + 1, n - 1)$) when n is even (resp. odd).

$$\begin{array}{ccccccc} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 \\ + & & - & & + & & - \end{array} \quad \begin{array}{ccccccc} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 \oplus L_5 \\ + & & - & & + & & - & + \end{array}$$

A(lternating)-surfaces

Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).



$\Rightarrow (p, q) = (n, n)$ (resp. $(n + 1, n - 1)$) when n is even (resp. odd).


$$\begin{matrix} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 \\ + & & - & & + & & - \end{matrix} \quad \begin{matrix} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 & \oplus & L_5 \\ + & & - & & + & & - & & + \end{matrix}$$

Gauss map $\psi : \Sigma \rightarrow \mathbf{S}_{p,q}$ of an A-surface $\Sigma \subset \mathbb{H}^{p,q}$:

A(Iternating)-surfaces

Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).



$\Rightarrow (p, q) = (n, n)$ (resp. $(n + 1, n - 1)$) when n is even (resp. odd).

$$\begin{matrix} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 \\ + & & - & & + & & - \end{matrix} \quad \begin{matrix} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 & \oplus & L_5 \\ + & & - & & + & & - & & + \end{matrix}$$


Gauss map $\psi : \Sigma \rightarrow \mathbf{S}_{p,q}$ of an A-surface $\Sigma \subset \mathbb{H}^{p,q}$:

$\psi(x) :=$ totally geodesic q -sphere generated by $(L_2 \oplus L_4 \oplus \cdots)|_x$.

A(Iternating)-surfaces

Definition

When M is pseudo-Riemannian, Σ is called an **A-surface** if it is quasi-superminimal and L_i is positive (resp. negative) definite for all i odd (resp. even) (*alternating space-/time-likeness*).



$\Rightarrow (p, q) = (n, n)$ (resp. $(n + 1, n - 1)$) when n is even (resp. odd).

$$\begin{array}{ccccccccc} L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 & & L_1 & \oplus & L_2 & \oplus & L_3 & \oplus & L_4 & \oplus & L_5 \\ + & & - & & + & & - & & + & & - & & + & & - & & + \end{array}$$

Gauss map $\psi : \Sigma \rightarrow \mathbf{S}_{p,q}$ of an A-surface $\Sigma \subset \mathbb{H}^{p,q}$:

$\psi(x) :=$ totally geodesic q -sphere generated by $(L_2 \oplus L_4 \oplus \cdots)|_x$.

Infinitesimal rigidity

Generalization of Maximality Theorem:

Min-Max Theorem (N. 2022)

Given A-surface $\Sigma \subset \mathbb{H}^{p,q}$ with $T\mathbb{H}^{p,q}|_{\Sigma} = L_1 \oplus \cdots \oplus L_n$ ($L_1 = T\Sigma$).

Generalization of Maximality Theorem:

Min-Max Theorem (N. 2022)

Given A-surface $\Sigma \subset \mathbb{H}^{p,q}$ with $T\mathbb{H}^{p,q}|_{\Sigma} = L_1 \oplus \cdots \oplus L_n$ ($L_1 = T\Sigma$).

Suppose that each of L_2, \dots, L_n , viewed as a hermitian holomorphic line bundle, has *semi-negative* Chern form (no assumption on L_1).

Infinitesimal rigidity

Generalization of Maximality Theorem:

Min-Max Theorem (N. 2022)

Given A-surface $\Sigma \subset \mathbb{H}^{p,q}$ with $T\mathbb{H}^{p,q}|_{\Sigma} = L_1 \oplus \cdots \oplus L_n$ ($L_1 = T\Sigma$).
Suppose that each of L_2, \dots, L_n , viewed as a hermitian holomorphic line bundle, has *semi-negative* Chern form (no assumption on L_1).
Then any nonzero, compactly supported deformation of Σ along $N^+ := L_3 \oplus L_5 \oplus \cdots$ (resp. $N^- := L_2 \oplus L_4 \oplus \cdots$) locally increases (resp. decreases) the area.

Infinitesimal rigidity

Generalization of Maximality Theorem:

Min-Max Theorem (N. 2022)

Given A-surface $\Sigma \subset \mathbb{H}^{p,q}$ with $T\mathbb{H}^{p,q}|_{\Sigma} = L_1 \oplus \cdots \oplus L_n$ ($L_1 = T\Sigma$). Suppose that each of L_2, \dots, L_n , viewed as a hermitian holomorphic line bundle, has *semi-negative* Chern form (no assumption on L_1). Then any nonzero, compactly supported deformation of Σ along $N^+ := L_3 \oplus L_5 \oplus \cdots$ (resp. $N^- := L_2 \oplus L_4 \oplus \cdots$) locally increases (resp. decreases) the area.

Remark. Cannot expect such property from minimal surfaces in $\mathbb{H}^p \subset \mathbb{H}^{p,q}$ (unless Σ is totally geodesic or close to t.g.).

Infinitesimal rigidity

Generalization of Maximality Theorem:

Min-Max Theorem (N. 2022)

Given A-surface $\Sigma \subset \mathbb{H}^{p,q}$ with $T\mathbb{H}^{p,q}|_{\Sigma} = L_1 \oplus \cdots \oplus L_n$ ($L_1 = T\Sigma$). Suppose that each of L_2, \dots, L_n , viewed as a hermitian holomorphic line bundle, has *semi-negative* Chern form (**no assumption on L_1**). Then any nonzero, compactly supported deformation of Σ along $N^+ := L_3 \oplus L_5 \oplus \cdots$ (resp. $N^- := L_2 \oplus L_4 \oplus \cdots$) locally increases (resp. decreases) the area.

Remark. Cannot expect such property from minimal surfaces in $\mathbb{H}^p \subset \mathbb{H}^{p,q}$ (unless Σ is totally geodesic or close to t.g.).

What is different for A-surfaces: the 2nd fundamental form $\text{Sym}^2 T\Sigma = \text{Sym}^2 L_1 \rightarrow N$ takes values in L_2 , which is timelike (**A-surfaces have “timelike osculation”**).

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, N) \rightarrow C^\infty(\Sigma, N)$ ($L_\Sigma = \Delta^N + 0\text{th order part}$)

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, N) \rightarrow C^\infty(\Sigma, N)$ ($L_\Sigma = \Delta^N + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, N)$,

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

► $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, N) \rightarrow C^\infty(\Sigma, N)$ ($L_\Sigma = \Delta^N + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, N)$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_- + \xi_+ \neq 0$ with $L_\Sigma \xi = 0$.

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_- + \xi_+ \neq 0$ with $L_\Sigma \xi = 0$.
 $\Rightarrow \langle L_\Sigma \xi, \xi_+ \rangle = 0, \langle L_\Sigma \xi, \xi_- \rangle = 0$

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt^2} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_+ + \xi_- \neq 0$ with $L_\Sigma \xi = 0$.

$$\Rightarrow \langle L_\Sigma \xi, \xi_+ \rangle = 0, \langle L_\Sigma \xi, \xi_- \rangle = 0$$

$$\Rightarrow \langle L_\Sigma \xi_+, \xi_+ \rangle = -\langle L_\Sigma \xi_-, \xi_+ \rangle, \langle L_\Sigma \xi_-, \xi_- \rangle = -\langle L_\Sigma \xi_+, \xi_- \rangle$$

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, N) \rightarrow C^\infty(\Sigma, N)$ ($L_\Sigma = \Delta^N + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, N)$,

- ▶ $\frac{d}{dt} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_+ + \xi_- \neq 0$ with $L_\Sigma \xi = 0$.

$$\Rightarrow \langle L_\Sigma \xi, \xi_+ \rangle = 0, \langle L_\Sigma \xi, \xi_- \rangle = 0$$

$$\Rightarrow \langle L_\Sigma \xi_+, \xi_+ \rangle = -\langle L_\Sigma \xi_-, \xi_+ \rangle, \langle L_\Sigma \xi_-, \xi_- \rangle = -\langle L_\Sigma \xi_+, \xi_- \rangle$$

Min-Max Thm. \Rightarrow integral $<0, >0$, resp.

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, \mathbb{N}) \rightarrow C^\infty(\Sigma, \mathbb{N})$ ($L_\Sigma = \Delta^\mathbb{N} + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, \mathbb{N})$,

- ▶ $\frac{d}{dt} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle$,
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_- + \xi_+ \neq 0$ with $L_\Sigma \xi = 0$.

$$\Rightarrow \langle L_\Sigma \xi, \xi_+ \rangle = 0, \langle L_\Sigma \xi, \xi_- \rangle = 0$$

$$\Rightarrow \langle L_\Sigma \xi_+, \xi_+ \rangle = -\langle L_\Sigma \xi_-, \xi_+ \rangle, \langle L_\Sigma \xi_-, \xi_- \rangle = -\langle L_\Sigma \xi_+, \xi_- \rangle$$

Min-Max Thm. \Rightarrow integral $<0, >0$, resp. L_Σ is self-adjoint

integrals are equal because

Infinitesimal rigidity

Recall: Any minimal submanifold Σ has a **Jacobi operator**

$L_\Sigma : C^\infty(\Sigma, N) \rightarrow C^\infty(\Sigma, N)$ ($L_\Sigma = \Delta^N + 0\text{th order part}$)

s.t. \forall deformation (Σ_t) with variational vector field $\xi \in C^\infty(\Sigma, N)$,

- ▶ $\frac{d}{dt} \Big|_{t=0} \text{vol}(\Sigma_t) = - \int_\Sigma \langle L_\Sigma \xi, \xi \rangle,$
- ▶ if very Σ_t is minimal, then $L_\Sigma \xi = 0$.

So those ξ with $L_\Sigma \xi = 0$ are call **Jacobi fields** and can be viewed as the infinitesimal deformations of Σ .

Corollary of Min-Max Theorem (Infinitesimal Rigidity)

Under the same assumption, Σ does not have any nonzero compactly supported Jacobi field.

Proof. A-surface Σ NOT inf. rigid $\Rightarrow \exists \xi = \xi_- + \xi_+ \neq 0$ with $L_\Sigma \xi = 0$.

$$\Rightarrow \langle L_\Sigma \xi, \xi_+ \rangle = 0, \langle L_\Sigma \xi, \xi_- \rangle = 0$$

$$\Rightarrow \langle L_\Sigma \xi_+, \xi_+ \rangle = -\langle L_\Sigma \xi_-, \xi_+ \rangle, \langle L_\Sigma \xi_-, \xi_- \rangle = -\langle L_\Sigma \xi_+, \xi_- \rangle$$

Min-Max Thm. \Rightarrow integral $<0, >0$, resp. **contradiction!** integrals are equal because L_Σ is self-adjoint



Remark. This is an infinite-dimensional generalization of the linear-algebraic fact “if $A = \begin{pmatrix} A_+ & * \\ * & A_- \end{pmatrix}$ is symmetric and the blocks A_+ and A_- are positive/negative definite resp., then $\det(A) \neq 0$ ”

Infinitesimal rigidity

Remark. This is an infinite-dimensional generalization of the linear-algebraic fact “if $A = \begin{pmatrix} A_+ & * \\ * & A_- \end{pmatrix}$ is symmetric and the blocks A_+ and A_- are positive/negative definite resp., then $\det(A) \neq 0$ ” (saddle-type critical points are non-degenerate).

Higgs bundles

A **Higgs bundle** on a closed Riemann surface Σ is a pair (E, Φ) where

- E is a holomorphic vector bundle on Σ ;
- $\Phi \in H^0(\Sigma, K_\Sigma \otimes \text{End}(E))$ (**$\text{End}(E)$ -valued holomorphic 1-form**).

Higgs bundles

A **Higgs bundle** on a closed Riemann surface Σ is a pair (E, Φ) where

- E is a holomorphic vector bundle on Σ ;
- $\Phi \in H^0(\Sigma, K_\Sigma \otimes \text{End}(E))$ (**$\text{End}(E)$ -valued holomorphic 1-form**).

Given a complex reductive group $G_{\mathbb{C}}$ (e.g. $\text{SL}(n, \mathbb{C})$),

replace E by a holomorphic principal $G_{\mathbb{C}}$ -bundle

\rightsquigarrow **$G_{\mathbb{C}}$ -Higgs bundle**.

Higgs bundles

A **Higgs bundle** on a closed Riemann surface Σ is a pair (E, Φ) where

- E is a holomorphic vector bundle on Σ ;
- $\Phi \in H^0(\Sigma, K_\Sigma \otimes \text{End}(E))$ (**End(E)-valued holomorphic 1-form**).

Given a complex reductive group $G_{\mathbb{C}}$ (e.g. $\text{SL}(n, \mathbb{C})$),

replace E by a holomorphic principal $G_{\mathbb{C}}$ -bundle

\rightsquigarrow **$G_{\mathbb{C}}$ -Higgs bundle**.

Given a real semisimple Lie group G (e.g. $\text{SL}(n, \mathbb{R})$),

$G_{\mathbb{C}}$ -Higgs bundle ($G_{\mathbb{C}}$ = complexification of G) + some extra structure

\rightsquigarrow **G -Higgs bundle**.

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

the connection $D = \nabla^h + \Phi + \Phi^{*h}$ is flat.

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

the connection $D = \nabla^h + \Phi + \Phi^{*h}$ is flat.

(harmonic metric, Hitchin equation, Hitchin connection)

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

the connection $D = \nabla^h + \Phi + \Phi^{*h}$ is flat.

(harmonic metric, Hitchin equation, Hitchin connection)

\Rightarrow Give representation $\rho : \pi_1(\Sigma) \rightarrow \mathrm{GL}(n, \mathbb{C})$ and

ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow \mathrm{GL}(n, \mathbb{C})/\mathrm{U}(n)$.

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

the connection $D = \nabla^h + \Phi + \Phi^{*h}$ is flat.

(harmonic metric, Hitchin equation, Hitchin connection)

\Rightarrow Give representation $\rho : \pi_1(\Sigma) \rightarrow \mathrm{GL}(n, \mathbb{C})$ and

ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow \mathrm{GL}(n, \mathbb{C})/\mathrm{U}(n)$.

If (E, Φ) is G -Higgs bundle for real semisimple G

Higgs bundles

(Part of) nonabelian Hodge correspondence:

$\deg(E) = 0$ and (E, Φ) is **stable**

$\Rightarrow \exists!$ hermitian metric h on E satisfying $F_{\nabla^h} + [\Phi, \Phi^{*h}] = 0$;

the connection $D = \nabla^h + \Phi + \Phi^{*h}$ is flat.

(harmonic metric, Hitchin equation, Hitchin connection)

\Rightarrow Give representation $\rho : \pi_1(\Sigma) \rightarrow \mathrm{GL}(n, \mathbb{C})$ and

ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow \mathrm{GL}(n, \mathbb{C})/\mathrm{U}(n)$.

If (E, Φ) is G -Higgs bundle for real semisimple G

$\Rightarrow \rho$ has image in G ; f has image in G/K .

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Special case for today.

$$E = L_n^{-1} \oplus \cdots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \cdots \oplus L_n$$

$$\Phi = \left(\begin{array}{cccc|ccc|cccc} 0 & & & & & & & & & & \beta & \\ \alpha_n & 0 & & & & & & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & & & & & & \\ & & \ddots & \ddots & & & & & & & & \\ & & & \alpha_2 & 0 & & & & & & & \\ \hline & & & & \alpha_1 & 0 & & & & & & \\ \hline & & & & & \alpha_1 & 0 & & & & & \\ & & & & & & \alpha_2 & \ddots & & & & \\ & & & & & & & \ddots & \ddots & & & \\ & & & & & & & & \alpha_{n-1} & 0 & & \\ & & & & & & & & & \alpha_n & 0 & \end{array} \right)$$

is a **cyclic** $\mathrm{SO}_0(n, n+1)$ -Higgs bundle

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Special case for today.

$$E = L_n^{-1} \oplus \cdots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \cdots \oplus L_n$$

L_1, \dots, L_n :
holo. line bundle

$$\alpha_i \in H^0(\Sigma, K_\Sigma L_i)$$

$$\alpha_i \in H^0(\Sigma, K_\Sigma L_{i-1}^{-1} L_i)$$

$$i=2, \dots, n$$

$$\beta \in H^0(\Sigma, K_\Sigma L_{n-1}^{-1} L_n)$$

$$\Phi = \left(\begin{array}{cccc|cc|cccc} 0 & & & & & & & & & \beta \\ \alpha_n & 0 & & & & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & & & & \\ & & \ddots & \ddots & & & & & & \\ & & & \alpha_2 & 0 & & & & & \\ \hline & & & \alpha_1 & 0 & & & & & \\ \hline & & & \alpha_1 & 0 & & & & & \\ & & & & \alpha_2 & \ddots & & & & \\ & & & & & \ddots & \ddots & & & \\ & & & & & & \alpha_{n-1} & 0 & & \\ & & & & & & & \alpha_n & 0 & \end{array} \right)$$

is a **cyclic** $\mathrm{SO}_0(n, n+1)$ -Higgs bundle

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Special case for today.

$$E = L_n^{-1} \oplus \cdots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \cdots \oplus L_n$$

L_1, \dots, L_n :
holo. line bundle
 $\alpha_i \in H^0(\Sigma, K_\Sigma L_i)$
 $\alpha_i \in H^0(\Sigma, K_\Sigma L_{i-1}^{-1} L_i)$
 $i=2, \dots, n$
 $\beta \in H^0(\Sigma, K_\Sigma L_{n-1}^{-1} L_n)$

$$\Phi = \left(\begin{array}{cccc|cc|cccc} 0 & & & & & & & & & \beta \\ \alpha_n & 0 & & & & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & & & & \\ & & \ddots & \ddots & & & & & & \\ & & & \alpha_2 & 0 & & & & & \\ \hline & & & \alpha_1 & 0 & & & & & \\ \hline & & & \alpha_1 & 0 & & & & & \\ & & & & \alpha_2 & \ddots & & & & \\ & & & & & \ddots & \ddots & & & \\ & & & & & & \alpha_{n-1} & 0 & & \\ & & & & & & & \alpha_n & 0 & \end{array} \right)$$

is a **cyclic** $\mathrm{SO}_0(n, n+1)$ -Higgs bundle

(= “sub-cyclic” $\mathrm{SL}(2n+1, \mathbb{R})$ -Higgs bundle)

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Properties.

- (Baraglia, Collier-Li) The harmonic metric h on $E = L_n^{-1} \oplus \cdots \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus \cdots \oplus L_n$ has the form $h = h_n^{-1} \oplus \cdots \oplus h_1^{-1} \oplus 1 \oplus h_1 \oplus \cdots \oplus h_n$ for hermitian metric h_i on L_i .

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Properties.

- ▶ (Baraglia, Collier-Li) The harmonic metric h on $E = L_n^{-1} \oplus \cdots \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus \cdots \oplus L_n$ has the form $h = h_n^{-1} \oplus \cdots \oplus h_1^{-1} \oplus 1 \oplus h_1 \oplus \cdots \oplus h_n$ for hermitian metric h_i on L_i .
- ▶ The harmonic map $f : \tilde{\Sigma} \rightarrow \mathbf{S}_{n,n} = \mathbf{S}_{n+1,n-1}$ is a conformal minimal immersion.

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1$

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$
 $\Rightarrow \rho$ is in a **Collier component** $\mathbf{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$,
 $0 \leq d \leq (2g-2)n$

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$
 $\Rightarrow \rho$ is in a **Collier component** $\mathbf{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$,
 $0 \leq d \leq (2g-2)n$ (**Col**_{*d*} consists of Θ -positive Anosov
 $\mathrm{SO}_0(n, n+1)$ -representations;

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$
 $\Rightarrow \rho$ is in a **Collier component** $\mathbf{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$,
 $0 \leq d \leq (2g-2)n$ (**Col**_{*d*} consists of Θ -positive Anosov
 $\mathrm{SO}_0(n, n+1)$ -representations; $\mathbf{Col}_{(2g-2)n} = \mathbf{Hit}$;

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$
 $\Rightarrow \rho$ is in a **Collier component** $\mathbf{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$,
 $0 \leq d \leq (2g-2)n$ (**Col**_{*d*} consists of Θ -positive Anosov
 $\mathrm{SO}_0(n, n+1)$ -representations; $\mathbf{Col}_{(2g-2)n} = \mathbf{Hit}$; $n = 1 \rightsquigarrow$ the
familiar components for $\mathrm{SO}_0(2, 1) = \mathrm{PSL}(2, \mathbb{R})$).

Cyclic $\mathrm{SO}_0(n, n+1)$ -Higgs bundles

Sub-cases.

- ▶ If $L_1 = K_\Sigma^{-1}$, $\alpha_1 = 1 \in H^0(\Sigma, K_\Sigma L_1) = H^0(\Sigma, \mathcal{O})$,
 $L_2 = K_\Sigma^{-2}$, $\alpha_2 = 1 \in H^0(\Sigma, K_\Sigma L_1^{-1} L_2) = H^0(\Sigma, \mathcal{O})$,
 \dots , $L_n = K_\Sigma^{-n}$, $\alpha_n = 1 \Rightarrow \rho$ is in the **Hitchin component**
 $\mathbf{Hit}(S_g, \mathrm{SO}_0(n, n+1)) \subset \mathbf{Hit}(S_g, \mathrm{SL}(2n+1, \mathbb{R}))$.
- ▶ If moreover $n = 3 \Rightarrow \rho$ is in $\mathbf{Hit}(S_g, G'_2) \subset \mathbf{Hit}(S_g, \mathrm{SO}_0(3, 4))$.
- ▶ More generally, if $L_k = K_\Sigma^{-k}$ and $\alpha_k = 1$ for $k = 1, \dots, n-1$
 $\Rightarrow \rho$ is in a **Collier component** $\mathbf{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$,
 $0 \leq d \leq (2g-2)n$ (**Col** _{d} consists of Θ -positive Anosov
 $\mathrm{SO}_0(n, n+1)$ -representations; $\mathbf{Col}_{(2g-2)n} = \mathbf{Hit}$; $n = 1 \rightsquigarrow$ the
familiar components for $\mathrm{SO}_0(2, 1) = \mathrm{PSL}(2, \mathbb{R})$).

Theorem (N. 2022)

If $L_1 = K_\Sigma^{-1}$ and $\alpha_1 = 1$, Then $\exists!$ A-surface immersion $F : \tilde{\Sigma} \rightarrow \mathbb{H}^{n,n}$ (if n is even) or $\mathbb{H}^{n+1, n-1}$ (if n is odd) such that f is the composition of F with the Gauss map of F .

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$
$$(E, \Phi) \mapsto [\rho]$$

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$
$$(E, \Phi) \mapsto [\rho]$$

Labourie Conjecture: Ψ is injective

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$
$$(E, \Phi) \mapsto [\rho]$$

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$
$$(E, \Phi) \mapsto [\rho]$$

restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$

constructed from
holo. differentials

$$(E, \Phi) \mapsto [\rho]$$

restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$

constructed from
holo. differentials

holo. quadratic diff. part = 0 $(E, \Phi) \mapsto [\rho]$

restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R}), n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$

not fixed
constructed from holo. differentials
holo. quadratic diff. part = 0

$$(E, \Phi) \mapsto [\rho]$$

restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R})$, $n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \rightarrow \mathbf{Hit}(S_g, G)$$

not fixed

constructed from holo. differentials

holo. quadratic diff. part = 0

$(E, \Phi) \mapsto [\rho]$

holo. vector bundle on \mathcal{T}_g
restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R})$, $n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \xrightarrow{\text{MCG-equivariant}} \mathbf{Hit}(S_g, G)$$

not fixed

constructed from holo. differentials

holo. quadratic diff. part = 0

holo. vector bundle on \mathcal{T}_g

$(E, \Phi) \mapsto [\rho]$

restricts to an immersion on the **cyclic** locus

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R})$, $n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \xrightarrow{\text{MCG-equivariant}} \mathbf{Hit}(S_g, G)$$

not fixed

constructed from holo. differentials

holo. quadratic diff. part = 0

$(E, \Phi) \mapsto [\rho]$

restricts to an immersion on the **cyclic** locus (i.e. locus where only the highest holomorphic differential is nonzero).

Labourie Conjecture: Ψ is injective (already known to be surjective).

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R})$, $n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal } G\text{-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \xrightarrow{\text{MCG-equivariant}} \mathbf{Hit}(S_g, G)$$

not fixed (above the set)

constructed from holo. differentials (next to Hitchin section)

holo. quadratic diff. part = 0 (arrow from conformal G-Higgs bundles)

$(E, \Phi) \mapsto [\rho]$ (next to the arrow)

holo. vector bundle on \mathcal{T}_g (arrow from Hitchin section)

restricts to an immersion on the **cyclic** locus (i.e. locus where only the highest holomorphic differential is nonzero).

Labourie Conjecture: Ψ is injective (already known to be surjective).

Corollary (proof of Labourie Conjecture in rank 2)

If G has rank 2 (the whole domain of Ψ is cyclic), then Ψ is injective.

Labourie Immersion Theorem & Labourie Conjecture

Labourie Immersion Theorem (2017)

Given any **split** real semisimple Lie group G of rank ≥ 2 (e.g. $SL(n, \mathbb{R})$, $n \geq 3$), the holonomy map

$$\Psi : \left\{ \begin{array}{l} \text{genus } g \text{ Riemann surfaces w.} \\ \text{conformal G-Higgs bundles} \\ \text{in Hitchin section} \end{array} \right\} / \sim \xrightarrow{\text{MCG-equivariant}} \mathbf{Hit}(S_g, G)$$

not fixed (above the set)

constructed from holo. differentials (next to Hitchin section)

holo. quadratic diff. part = 0 (arrow from conformal G-Higgs bundles)

$(E, \Phi) \mapsto [\rho]$ (next to the arrow)

holo. vector bundle on \mathcal{T}_g (arrow from Hitchin section)

restricts to an immersion on the **cyclic** locus (i.e. locus where only the highest holomorphic differential is nonzero).

Labourie Conjecture: Ψ is injective (already known to be surjective).

Corollary (proof of Labourie Conjecture in rank 2)

If G has rank 2 (the whole domain of Ψ is cyclic), then Ψ is injective.

Markovic, Markovic-Sagman-Smillie, Sagman-Smillie:

disproved the conjecture for rank ≥ 3 .

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary.

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary.

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary. Equivariance:

$$f_t(\gamma.x) = \rho_t(\gamma)f_t(x)$$

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary. Equivariance:

$$f_t(\gamma.x) = \rho_t(\gamma)f_t(x) \Rightarrow \dot{f}_0(\gamma.x) = \dot{\rho}_0(\gamma)f_0(x) + \rho_0(\gamma)\dot{f}_0(x)$$

Strategy for LIT through infinitesimal rigidity

- Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary. Equivariance:

$$f_t(\gamma.x) = \rho_t(\gamma)f_t(x) \Rightarrow \dot{f}_0(\gamma.x) = \cancel{\dot{\rho}_0(\gamma)}f_0(x) + \rho_0(\gamma)\dot{f}_0(x)$$

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary. Equivariance:

$$f_t(\gamma.x) = \rho_t(\gamma)f_t(x) \Rightarrow \dot{f}_0(\gamma.x) = \cancel{\dot{p}_0(\gamma)}f_0(x) + \rho_0(\gamma)\dot{f}_0(x)$$

This means \dot{f}_0 is a Jacobi field on the quotient minimal surface, contradicting rigidity. □

Strategy for LIT through infinitesimal rigidity

- ▶ Given (E, Φ) in the domain of Ψ , consider the conformal minimal immersion $f : \tilde{\Sigma} \rightarrow G/K$.

Lemma

If the quotient closed minimal surface in $\rho(\pi_1(S_g)) \backslash (G/K)$ given by f is **infinitesimally rigid**, then $d\Psi$ is nondegenerate at (E, Φ) .

Proof sketch. If degenerate, then \exists path (E_t, Φ_t) s.t. (ρ_t) is stationary. Equivariance:

$$f_t(\gamma.x) = \rho_t(\gamma)f_t(x) \Rightarrow \dot{f}_0(\gamma.x) = \cancel{\dot{\rho}_0(\gamma)}f_0(x) + \rho_0(\gamma)\dot{f}_0(x)$$

This means \dot{f}_0 is a Jacobi field on the quotient minimal surface, contradicting rigidity. □

- ▶ However, nobody knows how to show such a rigidity result...

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \leadsto LIT for $SL(3, \mathbb{R})$;

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n + 1)$

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n + 1)$
 - (N. 2022) A-surfaces \rightsquigarrow LIT for Collier's component $\mathrm{Col}_d(S_g, \mathrm{SO}_0(n, n + 1))$ with $d \neq 0$

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n+1)$
 - (N. 2022) A-surfaces \rightsquigarrow LIT for Collier's component $\mathrm{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$ with $d \neq 0$
(\Rightarrow new proof of Labourie Conjecture for G'_2).

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n+1)$
 - (N. 2022) A-surfaces \rightsquigarrow LIT for Collier's component $\mathrm{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$ with $d \neq 0$
(\Rightarrow new proof of Labourie Conjecture for G'_2).

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n+1)$
 - (N. 2022) A-surfaces \rightsquigarrow LIT for Collier's component $\mathrm{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$ with $d \neq 0$
(\Rightarrow **new proof of Labourie Conjecture for G'_2**).

Remark. The Min-Max Theorem requires a negativity assumption on some Chern forms.

Strategy for LIT through infinitesimal rigidity

- ▶ ... nevertheless, one can replace f by some **other surfaces** with nice rigidity property and apply the same strategy:
 - (Labourie 2007, Loftin 2004) **hyperbolic affine spheres** in \mathbb{R}^3
 \rightsquigarrow LIT for $\mathrm{SL}(3, \mathbb{R})$;
 - (Labourie 2017) **cyclic surfaces** on some homogeneous space of $G_{\mathbb{C}}$
 \rightsquigarrow LIT for any rank 2 split G ;
 - (Collier-Tholozan-Toulisse 2019) **maximal surfaces** in $\mathbb{H}^{2,n}$
 \rightsquigarrow generalization of Labourie Conjecture to maximal representations in $\mathrm{SO}_0(2, n+1)$
 - (N. 2022) A-surfaces \rightsquigarrow LIT for Collier's component $\mathrm{Col}_d(S_g, \mathrm{SO}_0(n, n+1))$ with $d \neq 0$
(\Rightarrow **new proof of Labourie Conjecture for G'_2**).

Remark. The Min-Max Theorem requires a negativity assumption on some Chern forms. In the above application, this is guaranteed by:

Strategy for Labourie Immersion Theorem

Lemma (Dai-Li, N.)

Let h_1, \dots, h_n be hermitian metrics on holo. line bundles L_1, \dots, L_n over a closed Riemann surface solving the Hitchin eqn. for

$$E = L_n^{-1} \oplus \dots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \dots \oplus L_n,$$

$$\Phi = \left(\begin{array}{ccc|c|ccc} 0 & & & & & & \beta \\ \alpha_n & 0 & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & \\ & & \ddots & \ddots & & & \\ & & & \alpha_2 & 0 & & \\ \hline & & & \alpha_1 & 0 & & \\ \hline & & & \alpha_1 & 0 & & \\ & & & & \alpha_2 & \ddots & \\ & & & & & \ddots & \ddots \\ & & & & & & \alpha_{n-1} & 0 \\ & & & & & & & \alpha_n & 0 \end{array} \right).$$

Suppose $(\alpha_1) \prec (\alpha_2) \prec \dots \prec (\alpha_{n-1}) \prec \min\{(\alpha_n), (\beta)\}$. Then $\partial_{z\bar{z}}^2 h_i \leq 0$ for $i = 2, \dots, n$, with strict inequality except at the obvious zeros of $\partial_{z\bar{z}}^2 h_i$.

Strategy for Labourie Immersion Theorem

Lemma (Dai-Li, N.)

Let h_1, \dots, h_n be hermitian metrics on holo. line bundles L_1, \dots, L_n over a closed Riemann surface solving the Hitchin eqn. for

$$E = L_n^{-1} \oplus \dots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \dots \oplus L_n,$$

$$\Phi = \left(\begin{array}{ccc|c|ccc} 0 & & & & & & \beta \\ \alpha_n & 0 & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & \\ & & \ddots & \ddots & & & \\ & & & \alpha_2 & 0 & & \\ \hline & & & \alpha_1 & 0 & & \\ \hline & & & \alpha_1 & 0 & & \\ & & & & \alpha_2 & \ddots & \\ & & & & & \ddots & \\ & & & & & & \alpha_{n-1} & 0 \\ & & & & & & & \alpha_n & 0 \end{array} \right).$$

divisor of α_1

Suppose $(\alpha_1) \prec (\alpha_2) \prec \dots \prec (\alpha_{n-1}) \prec \min\{(\alpha_n), (\beta)\}$. Then $\partial_{z\bar{z}}^2 h_i \leq 0$ for $i = 2, \dots, n$, with strict inequality except at the obvious zeros of $\partial_{z\bar{z}}^2 h_i$.

Strategy for Labourie Immersion Theorem

Lemma (Dai-Li, N.)

Let h_1, \dots, h_n be hermitian metrics on holo. line bundles L_1, \dots, L_n over a closed Riemann surface solving the Hitchin eqn. for

$$E = L_n^{-1} \oplus \dots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \dots \oplus L_n,$$

$$\Phi = \left(\begin{array}{ccc|ccc|ccc} 0 & & & & & & & & \beta \\ \alpha_n & 0 & & & & & & & \beta \\ & \alpha_{n-1} & \ddots & & & & & & \\ & & \ddots & \ddots & & & & & \\ & & & \alpha_2 & 0 & & & & \\ \hline & & & & \alpha_1 & 0 & & & \\ \hline & & & & \alpha_1 & 0 & & & \\ & & & & & \alpha_2 & \ddots & & \\ & & & & & & \ddots & \ddots & \\ & & & & & & & \alpha_{n-1} & 0 \\ & & & & & & & & \alpha_n & 0 \end{array} \right).$$

< or both empty

divisor of α_1

Suppose $(\alpha_1) \prec (\alpha_2) \prec \dots \prec (\alpha_{n-1}) \prec \min\{(\alpha_n), (\beta)\}$. Then $\partial_{z\bar{z}}^2 h_i \leq 0$ for $i = 2, \dots, n$, with strict inequality except at the obvious zeros of $\partial_{z\bar{z}}^2 h_i$.

Strategy for Labourie Immersion Theorem

Lemma (Dai-Li, N.)

Let h_1, \dots, h_n be hermitian metrics on holo. line bundles L_1, \dots, L_n over a closed Riemann surface solving the Hitchin eqn. for

$$E = L_n^{-1} \oplus \dots \oplus L_2^{-1} \oplus L_1^{-1} \oplus \mathcal{O} \oplus L_1 \oplus L_2 \oplus \dots \oplus L_n,$$

holds in par. when $\alpha_1 = \dots = \alpha_{n-1} = 1$ (Collier components)

$$\Phi = \left(\begin{array}{ccc|cc} 0 & & & & \beta \\ \alpha_n & 0 & & & \beta \\ & \alpha_{n-1} & \ddots & & \\ & & \ddots & \ddots & \\ & & & \alpha_2 & 0 \\ \hline & & & \alpha_1 & 0 \\ \hline & & & \alpha_1 & 0 \\ & \alpha_2 & \ddots & & \\ & & \ddots & \ddots & \\ & & & \alpha_{n-1} & 0 \\ & & & \alpha_n & 0 \end{array} \right).$$

< or both empty

divisor of α_1

Suppose $(\alpha_1) \prec (\alpha_2) \prec \dots \prec (\alpha_{n-1}) \prec \min\{(\alpha_n), (\beta)\}$. Then $\partial_{z\bar{z}}^2 h_i \leq 0$ for $i = 2, \dots, n$, with strict inequality except at the obvious zeros of $\partial_{z\bar{z}}^2 h_i$.

Thank you for your attention!