

Hitchin representations and minimal surfaces

Nathaniel Sagman (joint work with P.Smillie and V.Marković-P.Smillie)

University of Luxembourg

December 12, 2022

- Hitchin representations and the Hitchin base

- Hitchin representations and the Hitchin base
- Minimal surfaces and Labourie's Conjecture

- Hitchin representations and the Hitchin base
- Minimal surfaces and Labourie's Conjecture
- Results

- Hitchin representations and the Hitchin base
- Minimal surfaces and Labourie's Conjecture
- Results
- Proofs

- Fix a closed oriented surface Σ_g of genus $g \geq 2$.

- Fix a closed oriented surface Σ_g of genus $g \geq 2$.
- If G is a Lie group, we set $\text{Rep}(G) = \text{Hom}(\pi_1(\Sigma_g), G) // G$.

- Fix a closed oriented surface Σ_g of genus $g \geq 2$.
- If G is a Lie group, we set $\text{Rep}(G) = \text{Hom}(\pi_1(\Sigma_g), G) // G$.
- Teichmüller space \mathbf{T}_g is 1) the moduli space of marked Riemann surfaces (S, f) on Σ_g , or 2)

{A component of discrete faithful representations} $\subset \text{Rep}(\text{PSL}(2, \mathbb{R}))$.

- Fix a closed oriented surface Σ_g of genus $g \geq 2$.
- If G is a Lie group, we set $\text{Rep}(G) = \text{Hom}(\pi_1(\Sigma_g), G) // G$.
- Teichmüller space \mathbf{T}_g is 1) the moduli space of marked Riemann surfaces (S, f) on Σ_g , or 2)

{A component of discrete faithful representations} $\subset \text{Rep}(\text{PSL}(2, \mathbb{R}))$.

- Basic facts: $\mathbf{T}_g \simeq_{C^\infty} \mathbb{R}^{6g-6}$, \mathbf{T}_g has a complex structure, the mapping class group $MCG(\Sigma_g)$ acts properly discontinuously.

- When unspecified, G is a semisimple real Lie group with no compact factors with finite center, i.e., $G = G_1 \times \cdots \times G_m$, with each G_i simple, non-compact type, and finite center.

Definition (Wienhard)

A higher Teichmüller space is a union of connected components of $\text{Rep}(G)$ consisting entirely of discrete and faithful representations.

- Let G be the adjoint form of a split real simple Lie group (ex. $G = \mathrm{PSL}(n, \mathbb{R})$).

- Let G be the adjoint form of a split real simple Lie group (ex. $G = \mathrm{PSL}(n, \mathbb{R})$).
- Up to conjugation, there exists a unique irreducible embedding $i_G : \mathrm{PSL}(2, \mathbb{R}) \rightarrow G$. This induces a map $(i_G)_* : \mathrm{Rep}(\mathrm{PSL}(2, \mathbb{R})) \rightarrow \mathrm{Rep}(G)$.

Definition

The Hitchin component $\mathrm{Hit}(G) = \{\text{Hitchin representations}\} = \{\text{Connected component of } \mathrm{Rep}(G) \text{ containing } (i_G)_*(\mathbf{T}_g)\}.$

- Let G be the adjoint form of a split real simple Lie group (ex. $G = \mathrm{PSL}(n, \mathbb{R})$).
- Up to conjugation, there exists a unique irreducible embedding $i_G : \mathrm{PSL}(2, \mathbb{R}) \rightarrow G$. This induces a map $(i_G)_* : \mathrm{Rep}(\mathrm{PSL}(2, \mathbb{R})) \rightarrow \mathrm{Rep}(G)$.

Definition

The Hitchin component $\mathrm{Hit}(G) = \{\text{Hitchin representations}\} = \{\text{Connected component of } \mathrm{Rep}(G) \text{ containing } (i_G)_*(\mathbf{T}_g)\}$.

- $\mathrm{Hit}(G)$ is homeomorphic to $\mathbb{R}^{(2g-2)\dim(G)}$ (Hitchin, 1990), the Hitchin component is a higher Teichmüller space (Labourie, Fock-Goncharov, 2006), and MCG acts properly discontinuously (Labourie, 2006).

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$

- Let S be a Riemann surface structure on Σ_g with canonical bundle \mathcal{K} .

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$

- Let S be a Riemann surface structure on Σ_g with canonical bundle \mathcal{K} .
- From now on, $\mathrm{Hit}(n) = \mathrm{Hit}(\mathrm{PSL}(n, \mathbb{R}))$.

Definition

The Hitchin base for $\mathrm{PSL}(n, \mathbb{C})$ is $B(S) = \bigoplus_{i=2}^n H^0(S, \mathcal{K}^i) \simeq_{\mathbb{C}^\infty} \mathbb{C}^{\dim G(g-1)}$.

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$

- Let S be a Riemann surface structure on Σ_g with canonical bundle \mathcal{K} .
- From now on, $\mathrm{Hit}(n) = \mathrm{Hit}(\mathrm{PSL}(n, \mathbb{R}))$.

Definition

The Hitchin base for $\mathrm{PSL}(n, \mathbb{C})$ is $B(S) = \bigoplus_{i=2}^n H^0(S, \mathcal{K}^i) \simeq_{\mathbb{C}^\infty} \mathbb{C}^{\dim G(g-1)}$.

- A section of \mathcal{K}^i is a tensor q locally of the form $q = q(z)dz^i$, $q(z)$ holomorphic.

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$

- Let S be a Riemann surface structure on Σ_g with canonical bundle \mathcal{K} .
- From now on, $\mathrm{Hit}(n) = \mathrm{Hit}(\mathrm{PSL}(n, \mathbb{R}))$.

Definition

The Hitchin base for $\mathrm{PSL}(n, \mathbb{C})$ is $B(S) = \bigoplus_{i=2}^n H^0(S, \mathcal{K}^i) \simeq_{\mathbb{C}} \mathbb{C}^{\dim G(g-1)}$.

- A section of \mathcal{K}^i is a tensor q locally of the form $q = q(z)dz^i$, $q(z)$ holomorphic.
- There is a surjective map



Hitchin section $b_S : B(S) \rightarrow \mathrm{Rep}(\mathrm{PSL}(n, \mathbb{C}))$ such that $b_S(B(S)) = \mathrm{Hit}(n)$.

More on the Hitchin base for $\mathrm{PSL}(n, \mathbb{C})$

Let

$$\mathfrak{t}^{\mathbb{C}} = \left\{ \begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{pmatrix} : a_i \in \mathbb{C}, \sum a_i = 0 \right\}, \quad (\text{Cartan subalgebra}).$$

A point in $B(S)$ is equivalent to

- a finite degree branched covering $S_B \rightarrow S$ (the cameral cover) whose deck group has a faithful representation $\eta : \mathrm{Deck}(S_B) \rightarrow S_n$ (the Weyl group).
- On S_B , a η -equivariant holomorphic 1-form with values in the trivial $\mathfrak{t}^{\mathbb{C}}$ -bundle (satisfying a condition relative to the branch points).

The action of $\alpha \in S_n$ on the $\mathfrak{t}^{\mathbb{C}}$ -bundle is

$$\alpha \cdot \begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{pmatrix} = \begin{pmatrix} a_{\alpha(1)} & & \\ & \ddots & \\ & & a_{\alpha(n)} \end{pmatrix}.$$

That is, we have a matrix of 1-forms on a cover S_B , whose entries are permuted by the Deck group action.

In one direction.

- Let

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

be the holomorphic matrix of 1-forms on S_B .

In one direction.

- Let

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

be the holomorphic matrix of 1-forms on S_B .

- Let $\sigma_2, \dots, \sigma_n$ be the elementary symmetric polynomials, acting on ϕ by

$$\sigma_2(\phi) = \sum_{j < k} \phi_j \phi_k, \sigma_3(\phi) = \sum_{j < k < \ell} \phi_j \phi_k \phi_\ell, \dots, \sigma_n(\phi) = \prod_{j=1}^n \phi_j.$$

In one direction.

- Let

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

be the holomorphic matrix of 1-forms on S_B .

- Let $\sigma_2, \dots, \sigma_n$ be the elementary symmetric polynomials, acting on ϕ by

$$\sigma_2(\phi) = \sum_{j < k} \phi_j \phi_k, \sigma_3(\phi) = \sum_{j < k < \ell} \phi_j \phi_k \phi_\ell, \dots, \sigma_n(\phi) = \prod_{j=1}^n \phi_j.$$

- $\sigma_i(\phi)$ defines a holomorphic i -differential on S_B .

More on the Hitchin base for $\mathrm{PSL}(n, \mathbb{C})$

In one direction.

- Let

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$


be the holomorphic matrix of 1-forms on S_B .

- Let $\sigma_2, \dots, \sigma_n$ be the elementary symmetric polynomials, acting on ϕ by

$$\sigma_2(\phi) = \sum_{j < k} \phi_j \phi_k, \sigma_3(\phi) = \sum_{j < k < \ell} \phi_j \phi_k \phi_\ell, \dots, \sigma_n(\phi) = \prod_{j=1}^n \phi_j.$$


- $\sigma_i(\phi)$ defines a holomorphic i -differential on S_B .
- For any $\alpha \in S_n$, $\sigma_i(\alpha \cdot \phi) = \sigma_i(\phi)$. By equivariance, each σ_i descends to a holomorphic i -differential q_i on S .

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$



- Once we've specified S_B , by (abelian) Hodge theory, ϕ is equivalent to an n -tuple of classes in $H^1(S_B, \mathbb{C})^n (= \mathrm{Rep}(\mathbb{C}^n))$ satisfying a linear relation.

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$



- Once we've specified S_B , by (abelian) Hodge theory, ϕ is equivalent to an n -tuple of classes in $H^1(S_B, \mathbb{C})^n (= \mathrm{Rep}(\mathbb{C}^n))$ satisfying a linear relation.
- The Hitchin section $b_S : B(S) \rightarrow \mathrm{Hit}(n)$ associates the data of the cover S_B and the (abelian) section ϕ to a unique Hitchin representation.

Hitchin's parametrization for $G = \mathrm{PSL}(n, \mathbb{R})$



The Hitchin section $b_S : B(S) \rightarrow \mathrm{Hit}(n)$ associates the (abelian) data of the section ϕ to a unique Hitchin representation.

Two issues with Hitchin's parametrization:

- No $\mathrm{MCG}(\Sigma_g)$ equivariance.
- In the induced complex structure, the inclusion $\mathbf{T}_g \rightarrow \mathrm{Hit}(n)$ is not holomorphic.

- For general admissible G , let K be a maximal compact subgroup. Then G/K admits a G -invariant metric ν , so that $(G/K, \nu)$ is a symmetric space of non-compact type.

- For general admissible G , let K be a maximal compact subgroup. Then G/K admits a G -invariant metric ν , so that $(G/K, \nu)$ is a symmetric space of non-compact type.
- Ex. for $G = \mathrm{PSL}(2, \mathbb{R})$, $K = \mathrm{PSO}(2, \mathbb{R})$ and $(G/K, \nu)$ hyperbolic space.

- For general admissible G , let K be a maximal compact subgroup. Then G/K admits a G -invariant metric ν , so that $(G/K, \nu)$ is a symmetric space of non-compact type.
- Ex. for $G = \mathrm{PSL}(2, \mathbb{R})$, $K = \mathrm{PSO}(2, \mathbb{R})$ and $(G/K, \nu)$ hyperbolic space.
- Given $[\rho] \in \mathrm{Rep}(G)$, we consider C^2 maps $h : \tilde{\Sigma}_g \rightarrow G/K$ that are ρ -equivariant: for all $\gamma \in \pi_1(\Sigma_g)$, $h \circ \gamma^{-1} = \rho(\gamma)h$.

- For general admissible G , let K be a maximal compact subgroup. Then G/K admits a G -invariant metric ν , so that $(G/K, \nu)$ is a symmetric space of non-compact type.
- Ex. for $G = \mathrm{PSL}(2, \mathbb{R})$, $K = \mathrm{PSO}(2, \mathbb{R})$ and $(G/K, \nu)$ hyperbolic space.
- Given $[\rho] \in \mathrm{Rep}(G)$, we consider C^2 maps $h : \tilde{\Sigma}_g \rightarrow G/K$ that are ρ -equivariant: for all $\gamma \in \pi_1(\Sigma_g)$, $h \circ \gamma^{-1} = \rho(\gamma)h$.
- Since ρ is acting by isometries on $(G/K, \nu)$, the pullback $h^* \mathrm{dvol}_\nu$ is $\pi_1(\Sigma_g)$ -invariant, and hence descends to a 2-form on Σ_g .

- For general admissible G , let K be a maximal compact subgroup. Then G/K admits a G -invariant metric ν , so that $(G/K, \nu)$ is a symmetric space of non-compact type.
- Ex. for $G = \mathrm{PSL}(2, \mathbb{R})$, $K = \mathrm{PSO}(2, \mathbb{R})$ and $(G/K, \nu)$ hyperbolic space.
- Given $[\rho] \in \mathrm{Rep}(G)$, we consider C^2 maps $h : \tilde{\Sigma}_g \rightarrow G/K$ that are ρ -equivariant: for all $\gamma \in \pi_1(\Sigma_g)$, $h \circ \gamma^{-1} = \rho(\gamma)h$.
- Since ρ is acting by isometries on $(G/K, \nu)$, the pullback $h^* \mathrm{dvol}_\nu$ is $\pi_1(\Sigma_g)$ -invariant, and hence descends to a 2-form on Σ_g .
- The area of the map is

$$\mathrm{Area}(h) = A(h) = \int_{\Sigma_g} h^* \mathrm{dvol}_\nu.$$

Definition

h is minimal if it is a critical point of A , i.e., for all smooth variations h_t through C^2 ρ -equivariant maps,

$$\left. \frac{d}{dt} \right|_{t=0} A(h_t) = 0.$$

Theorem (Labourie)

Suppose ρ lies in any known higher Teichmüller space. Then there exists an area minimizing ρ -equivariant minimal map $h : \tilde{\Sigma}_g \rightarrow (G/K, \nu)$.

Theorem (Labourie)

Suppose ρ lies in any known higher Teichmüller space. Then there exists an area minimizing ρ -equivariant minimal map $h : \tilde{\Sigma}_g \rightarrow (G/K, \nu)$.

- Given any ρ -equivariant minimal map h , there is a Riemann surface S such that $h : \tilde{S} \rightarrow (G/K, \nu)$ is weakly conformal.

Theorem (Labourie)

Suppose ρ lies in any known higher Teichmüller space. Then there exists an area minimizing ρ -equivariant minimal map $h : \tilde{\Sigma}_g \rightarrow (G/K, \nu)$.

- Given any ρ -equivariant minimal map h , there is a Riemann surface S such that $h : \tilde{S} \rightarrow (G/K, \nu)$ is weakly conformal.
- **Fact:** for $G = \mathrm{PSL}(n, \mathbb{R})$, there exists a weakly conformal minimal map $h : \tilde{S} \rightarrow (G/K, \nu)$ if and only if $h_S(\rho) = (0, q_3, \dots, q_n) \in \oplus_{i=2}^n H^0(S, \mathcal{K}^i)$, i.e., the q_2 term vanishes (or $\sigma_2(\phi) = 0$).

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.
- Define $L : \mathbf{M} \rightarrow \text{Hit}(n)$ by $L([(S, f)], q_3, \dots, q_n) = b_S(0, q_3, \dots, q_n)$.

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.
- Define $L : \mathbf{M} \rightarrow \text{Hit}(n)$ by $L([(S, f)], q_3, \dots, q_n) = b_S(0, q_3, \dots, q_n)$.
- Labourie's existence result for minimal surfaces says that L is surjective.

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.
- Define $L : \mathbf{M} \rightarrow \text{Hit}(n)$ by $L([(S, f)], q_3, \dots, q_n) = b_S(0, q_3, \dots, q_n)$.
- Labourie's existence result for minimal surfaces says that L is surjective.
- Labourie's conjecture is that L is injective.

Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow \text{PSL}(n, \mathbb{R})$ Hitchin, there exists a unique ρ -equivariant minimal surface in the symmetric space.

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.
- Define $L : \mathbf{M} \rightarrow \text{Hit}(n)$ by $L([(S, f)], q_3, \dots, q_n) = b_S(0, q_3, \dots, q_n)$.
- Labourie's existence result for minimal surfaces says that L is surjective.
- Labourie's conjecture is that L is injective.

Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow \text{PSL}(n, \mathbb{R})$ Hitchin, there exists a unique ρ -equivariant minimal surface in the symmetric space.

- At the time, known for $n = 2, 3$ (Labourie, Loftin).

The Labourie Conjecture

- Let $\mathbf{M} \rightarrow \mathbf{T}_g$ be the holomorphic vector bundle with fiber $\mathbf{M}|_{[(S,f)]} = \bigoplus_{i=3}^n H^0(S, \mathcal{K}^i)$.
- Define $L : \mathbf{M} \rightarrow \text{Hit}(n)$ by $L([(S, f)], q_3, \dots, q_n) = b_S(0, q_3, \dots, q_n)$.
- Labourie's existence result for minimal surfaces says that L is surjective.
- Labourie's conjecture is that L is injective.

Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow \text{PSL}(n, \mathbb{R})$ Hitchin, there exists a unique ρ -equivariant minimal surface in the symmetric space.

- At the time, known for $n = 2, 3$ (Labourie, Loftin).
- A positive resolution of the conjecture means that L provides a $\text{MCG}(\Sigma_g)$ -equivariant parametrization of $\text{Hit}(n)$. Note also the inclusion $\mathbf{T}_g \rightarrow \mathbf{M}$ is holomorphic.

In general, recall G/K is a symmetric space, and let ν be a G -invariant metric.

Generalized Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow G$ in any higher Teichmüller space, there exists a unique minimal surface in $(G/K, \nu)$.

Generalized Labourie Conjecture

In general, recall G/K is a symmetric space, and let ν be a G -invariant metric.

Generalized Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow G$ in any higher Teichmüller space, there exists a unique minimal surface in $(G/K, \nu)$.

The rank of G is the dimension of a maximal flat subspace of G/K . For example:
 $\text{rank}(\text{PSL}(n, \mathbb{R})) = n - 1$.

Generalized Labourie Conjecture

Recall G/K is a symmetric space, and let ν be an invariant metric.

Generalized Labourie Conjecture

Given $\rho : \pi_1(\Sigma_g) \rightarrow G$ in any higher Teichmüller space, there exists a unique minimal surface in $(G/K, \nu)$.

The rank of G is the dimension of a maximal flat subspace of G/K . For example: $\text{rank}(\text{PSL}(n, \mathbb{R})) = n - 1$.

Theorem (Labourie, Collier-Touhousse-Tholozan, Schoen, Wan)

The generalized Labourie conjecture holds for all Hitchin components and maximal components for Lie groups of rank 2.

Hitchin: $\text{PSL}(3, \mathbb{R})$, $\text{PSp}(4, \mathbb{R})$, G'_2 (Labourie), Maximal: $\text{PSL}(2, \mathbb{R})^2$ (Schoen, Wan), and Hermitian Lie groups like $\text{PSO}_0(2, n)$ (Collier-Tholozan-Touhousse). See also work of Nie on G'_2 .

One simpler example of a higher Teichmüller space is $\mathbf{T}_g^n \subset \text{Rep}(\text{PSL}(2, \mathbb{R})^n)$.

Theorem (Marković, 2021)

For all genus g large enough and $n \geq 3$, there exists product of n discrete faithful representations $\rho = (\rho_1, \dots, \rho_n) : \pi_1(\Sigma_g) \rightarrow \text{PSL}(2, \mathbb{R})^n$ with multiple minimal surfaces.

One simpler example of a higher Teichmüller space is $\mathbf{T}_g^n \subset \text{Rep}(\text{PSL}(2, \mathbb{R})^n)$.

Theorem (Marković, 2021)

For all genus g large enough and $n \geq 3$, there exists product of n discrete faithful representations $\rho = (\rho_1, \dots, \rho_n) : \pi_1(\Sigma_g) \rightarrow \text{PSL}(2, \mathbb{R})^n$ with multiple minimal surfaces.

Consequently, the generalized Labourie conjecture fails for \mathbf{T}_g^n , $n \geq 3$.

One simpler example of a higher Teichmüller space is $\mathbf{T}_g^n \subset \text{Rep}(\text{PSL}(2, \mathbb{R})^n)$.

Theorem (Marković, 2021)

For all genus g large enough and $n \geq 3$, there exists product of n discrete faithful representations $\rho = (\rho_1, \dots, \rho_n) : \pi_1(\Sigma_g) \rightarrow \text{PSL}(2, \mathbb{R})^n$ with multiple minimal surfaces.

Consequently, the generalized Labourie conjecture fails for \mathbf{T}_g^n , $n \geq 3$.

Marković-S.-Smillie, 2022

New geometric proof of the above result, works for all genus $g \geq 2$.

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and $n \geq 4$, there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with multiple minimal surfaces. That is, Labourie's conjecture fails for $n \geq 4$.

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and $n \geq 4$, there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with multiple minimal surfaces. That is, Labourie's conjecture fails for $n \geq 4$.

More generally, let G be simple split real or a product of simple split real Lie groups. We say that $\rho : \pi_1(\Sigma_g) \rightarrow G$ is Hitchin if each factor is Hitchin. Let ν be any G -invariant metric on G/K .

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and G as above with rank ≥ 3 , there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow G$ with multiple minimal surfaces in $(G/K, \nu)$.

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and $n \geq 4$, there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with multiple minimal surfaces. That is, Labourie's conjecture fails for $n \geq 4$.

More generally, let G be simple split real or a product of simple split real Lie groups. We say that $\rho : \pi_1(\Sigma_g) \rightarrow G$ is Hitchin if each factor is Hitchin. Let ν be any G -invariant metric on G/K .

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and G as above with rank ≥ 3 , there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow G$ with multiple minimal surfaces in $(G/K, \nu)$.

- The results hold for every known higher Teichmüller space such that the restriction of the Hitchin map is surjective.

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and $n \geq 4$, there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with multiple minimal surfaces. That is, Labourie's conjecture fails for $n \geq 4$.

More generally, let G be simple split real or a product of simple split real Lie groups. We say that $\rho : \pi_1(\Sigma_g) \rightarrow G$ is Hitchin if each factor is Hitchin. Let ν be any G -invariant metric on G/K .

Theorem (S.-Smillie, 2022)

For all $g \geq 3$ and G as above with rank ≥ 3 , there exists a Hitchin representation $\rho : \pi_1(\Sigma_g) \rightarrow G$ with multiple minimal surfaces in $(G/K, \nu)$.

- The results hold for every known higher Teichmüller space such that the restriction of the Hitchin map is surjective.
- We can prove the same result as well for $g = 2$. Writing up.

Corollary

For $n \geq 4$, $L : \mathbf{M} \rightarrow \text{Hit}(n)$ has no continuous section.

Corollary

For all G as above with rank at least 3, there exists a Hitchin representation with multiple area minimizing minimal surfaces.

Question

What are the fibers of $L : \mathbf{M} \rightarrow \text{Hit}(n)$?

Question

Is there a “good” parametrization of $\text{Hit}(n)$?

- Recall that h being minimal means that for all variations h_t ,

$$\frac{d}{dt}\bigg|_{t=0} A(h_t) = 0.$$

- Recall that h being minimal means that for all variations h_t ,

$$\frac{d}{dt}\big|_{t=0}A(h_t) = 0.$$

- Labourie's minimal surface is area minimizing, and hence stable: for all h_t ,

$$\frac{d^2}{dt^2}\big|_{t=0}A(h_t) \geq 0.$$

- Recall that h being minimal means that for all variations h_t ,

$$\frac{d}{dt}\bigg|_{t=0} A(h_t) = 0.$$

- Labourie's minimal surface is area minimizing, and hence stable: for all h_t ,

$$\frac{d^2}{dt^2}\bigg|_{t=0} A(h_t) \geq 0.$$

- We say that a minimal surface is unstable if there exists a variation such that

$$\frac{d^2}{dt^2}\bigg|_{t=0} A(h_t) < 0.$$

- Recall that h being minimal means that for all variations h_t ,

$$\frac{d}{dt}\bigg|_{t=0} A(h_t) = 0.$$

- Labourie's minimal surface is area minimizing, and hence stable: for all h_t ,

$$\frac{d^2}{dt^2}\bigg|_{t=0} A(h_t) \geq 0.$$

- We say that a minimal surface is unstable if there exists a variation such that

$$\frac{d^2}{dt^2}\bigg|_{t=0} A(h_t) < 0.$$

- We produce a Hitchin representation together with an unstable minimal surface. Since it must have a stable minimal surface as well, it must have at least two minimal surfaces.

Heuristic: minimal maps to buildings


Fix a Riemann surface S .

- Given $[\rho] \in \text{Rep}(\text{PSL}(n, \mathbb{C}))$ with $h_S([\rho]) = (0, q_3, \dots, q_n)$, there exists a Hitchin ray $(\rho_R)_{R \in [1, \infty)}$: a path of representations starting at ρ with $h_S(\rho_R) = (0, R^3 q_3, \dots, R^n q_n)$ and minimal maps h_R .

Heuristic: minimal maps to buildings

Fix a Riemann surface S .


- Given $[\rho] \in \text{Rep}(\text{PSL}(n, \mathbb{C}))$ with $h_S([\rho]) = (0, q_3, \dots, q_n)$, there exists a Hitchin ray $(\rho_R)_{R \in [1, \infty)}$: a path of representations starting at ρ with $h_S(\rho_R) = (0, R^3 q_3, \dots, R^n q_n)$ and minimal maps h_R .
- As $R \rightarrow \infty$ the minimal maps h_R “converge” to a minimal map to a building of rank $n - 1$.



Heuristic: minimal maps to buildings

Fix a Riemann surface S .

- Given $[\rho] \in \text{Rep}(\text{PSL}(n, \mathbb{C}))$ with $h_S([\rho]) = (0, q_3, \dots, q_n)$, there exists a Hitchin ray $(\rho_R)_{R \in [1, \infty)}$: a path of representations starting at ρ with $h_S(\rho_R) = (0, R^3 q_3, \dots, R^n q_n)$ and minimal maps h_R .
- As $R \rightarrow \infty$ the minimal maps h_R “converge” to a minimal map to a building of rank $n - 1$.



- Heuristic: minimal surfaces in rank k buildings look like minimal surfaces in \mathbb{R}^k . For $k \geq 3$, minimal surfaces in \mathbb{R}^k are often unstable.

- Starting with $(0, q_3, \dots, q_n)$ take the branched cover $\tau : S_B \rightarrow S$ and the holomorphic $\mathbb{t}^{\mathbb{C}}$ -valued 1-form

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

with $\sigma_1(\phi) := \sum_j \phi_j = 0$ and $q_2 = \sigma_2(\phi) = 0$, $q_i = \sigma_i(\phi)$ for $i > 2$.

- Starting with $(0, q_3, \dots, q_n)$ take the branched cover $\tau : S_B \rightarrow S$ and the holomorphic $\mathbb{t}^{\mathbb{C}}$ -valued 1-form

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

with $\sigma_1(\phi) := \sum_j \phi_j = 0$ and $q_2 = \sigma_2(\phi) = 0$, $q_i = \sigma_i(\phi)$ for $i > 2$.

- For each j , $\operatorname{Re}\phi_j$ is a harmonic 1-form. The cohomology class of $\operatorname{Re}\phi_j$ determines a representation $\chi_j : \pi_1(S_B) \rightarrow (\mathbb{R}, +)$.

- Starting with $(0, q_3, \dots, q_n)$ take the branched cover $\tau : S_B \rightarrow S$ and the holomorphic \mathbb{C} -valued 1-form

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

with $\sigma_1(\phi) := \sum_j \phi_j = 0$ and $q_2 = \sigma_2(\phi) = 0$, $q_i = \sigma_i(\phi)$ for $i > 2$.

- For each j , $\operatorname{Re}\phi_j$ is a harmonic 1-form. The cohomology class of $\operatorname{Re}\phi_j$ determines a representation $\chi_j : \pi_1(S_B) \rightarrow (\mathbb{R}, +)$.
- Lifting each ϕ_j to \tilde{S}_B and integrating the real part, we obtain a minimal map

$$f = (f_1, \dots, f_n) : \tilde{S}_B \rightarrow \mathbb{R}^n, \quad f_j(z) = \int_p^z \operatorname{Re}\tilde{\phi}_j$$

that is $\chi = (\chi_1, \dots, \chi_n)$ -equivariant and whose image is contained in the hyperplane $\{x = (x_1, \dots, x_n) \in \mathbb{R}^n : \sum_j x_j = 0\}$.

- Starting with $(0, q_3, \dots, q_n)$ take the branched cover $\tau : S_B \rightarrow S$ and the holomorphic \mathbb{C} -valued 1-form

$$\phi = \begin{pmatrix} \phi_1 & & \\ & \ddots & \\ & & \phi_n \end{pmatrix}$$

with $\sigma_1(\phi) := \sum_j \phi_j = 0$ and $q_2 = \sigma_2(\phi) = 0$, $q_i = \sigma_i(\phi)$ for $i > 2$.

- For each j , $\operatorname{Re}\phi_j$ is a harmonic 1-form. The cohomology class of $\operatorname{Re}\phi_j$ determines a representation $\chi_j : \pi_1(S_B) \rightarrow (\mathbb{R}, +)$.
- Lifting each ϕ_j to \tilde{S}_B and integrating the real part, we obtain a minimal map

$$f = (f_1, \dots, f_n) : \tilde{S}_B \rightarrow \mathbb{R}^n, \quad f_j(z) = \int_p^z \operatorname{Re}\tilde{\phi}_j$$

that is $\chi = (\chi_1, \dots, \chi_n)$ -equivariant and whose image is contained in the hyperplane $\{x = (x_1, \dots, x_n) \in \mathbb{R}^n : \sum_j x_j = 0\}$.

- Identifying the hyperplane with \mathbb{R}^{n-1} , we write $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$.

- Taking stock, we have a ray of representations $R \mapsto \rho_R : \pi_1(S) \rightarrow G = \mathrm{PSL}(n, \mathbb{C})$ with minimal maps h_R and an equivariant minimal map $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$.

- Taking stock, we have a ray of representations $R \mapsto \rho_R : \pi_1(S) \rightarrow G = \mathrm{PSL}(n, \mathbb{C})$ with minimal maps h_R and an equivariant minimal map $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$.
- Let $B \subset S_B$ be the branch locus of $\tau : S_B \rightarrow S$. Lift h_R to $\tilde{h}_R : \tilde{S}_B \rightarrow (G/K, \nu)$. The following is a consequence of an estimate of T. Mochizuki.

Loosely stated proposition

For “generic” Hitchin rays, the intrinsic data of \tilde{h}_R , rescaled by $1/R^2$, converges locally uniformly on $\tilde{S}_B - B$ to the intrinsic data of f . The convergence at $z \in \tilde{S}_B$ is $O(e^{-cd(z,B)R})$.

- Taking stock, we have a ray of representations $R \mapsto \rho_R : \pi_1(S) \rightarrow G = \mathrm{PSL}(n, \mathbb{C})$ with minimal maps h_R and an equivariant minimal map $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$.
- Let $B \subset S_B$ be the branch locus of $\tau : S_B \rightarrow S$. Lift h_R to $\tilde{h}_R : \tilde{S}_B \rightarrow (G/K, \nu)$. The following is a consequence of an estimate of T. Mochizuki.

Loosely stated proposition

For “generic” Hitchin rays, the intrinsic data of \tilde{h}_R , rescaled by $1/R^2$, converges locally uniformly on $\tilde{S}_B - B$ to the intrinsic data of f . The convergence at $z \in \tilde{S}_B$ is $O(e^{-cd(z,B)R})$.

- That is, away from B , the intrinsic data of the limiting minimal map to the building should look like f .

Minimal maps from the cameral cover

- Taking stock, we have a ray of representations $R \mapsto \rho_R : \pi_1(S) \rightarrow G = \mathrm{PSL}(n, \mathbb{C})$ with minimal maps h_R and an equivariant minimal map $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$.
- Let $B \subset S_B$ be the branch locus of $\tau : S_B \rightarrow S$. Lift h_R to $\tilde{h}_R : \tilde{S}_B \rightarrow (G/K, \nu)$. The following is a consequence of an estimate of T. Mochizuki.

Loosely stated proposition


For “generic” Hitchin rays, the intrinsic data of \tilde{h}_R , rescaled by $1/R^2$, converges locally uniformly on $\tilde{S}_B - B$ to the intrinsic data of f . The convergence at $z \in \tilde{S}_B$ is $O(e^{-cd(z,B)R})$.

- That is, away from B , the intrinsic data of the limiting minimal map to the building should look like f .
- **Remark:** for general G , we have a branched covering $S_B \rightarrow S$ with $\mathrm{Deck}(S_B/S)$ inside the Weyl group of the Lie algebra of G , and a minimal map to $\mathbb{R}^{\mathrm{rank} G}$.

- Every equivariant minimal map to \mathbb{R}^2 is stable. For $k \geq 3$, there exists unstable equivariant minimal maps $\tilde{S} \rightarrow \mathbb{R}^k$.


Minimal maps from the cameral cover

- Every equivariant minimal map to \mathbb{R}^2 is stable. For $k \geq 3$, there exists unstable equivariant minimal maps $\tilde{S} \rightarrow \mathbb{R}^k$.
- Simplest example: an unstable minimal map to the 3-torus $S \rightarrow T^3$, lifts to an unstable equivariant minimal map $\tilde{S} \rightarrow \mathbb{R}^3$.



Minimal maps from the cameral cover

- Every equivariant minimal map to \mathbb{R}^2 is stable. For $k \geq 3$, there exists unstable equivariant minimal maps $\tilde{S} \rightarrow \mathbb{R}^k$.
- Simplest example: an unstable minimal map to the 3-torus $S \rightarrow T^3$, lifts to an unstable equivariant minimal map $\tilde{S} \rightarrow \mathbb{R}^3$.



- **Remark:** in genus 2, every equivariant minimal surface is contained in a 2-plane, and hence stable!

The index of minimal maps

Set $G = \mathrm{PSL}(n, \mathbb{C})$, $K = \mathrm{PSU}(n, \mathbb{C})$.

- Formally, variations of a minimal map $h : \tilde{S} \rightarrow (G/K, \nu)$ are sections $\dot{h} \in \Gamma_S(h^*TG/K)$. We write " $h_t = h + t\dot{h}$."

The index of minimal maps

Set $G = \mathrm{PSL}(n, \mathbb{C})$, $K = \mathrm{PSU}(n, \mathbb{C})$.

- Formally, variations of a minimal map $h : \tilde{S} \rightarrow (G/K, \nu)$ are sections $\dot{h} \in \Gamma_S(h^*TG/K)$. We write " $h_t = h + t\dot{h}$."
- The index of h , $\mathrm{Ind}(h)$, is the maximal dimension of a subspace of $\Gamma_S(h^*TG/K)$ on which

$$\frac{d^2}{dt^2} \Big|_{t=0} A(h + t\dot{h}) < 0.$$

The index of minimal maps

Set $G = \mathrm{PSL}(n, \mathbb{C})$, $K = \mathrm{PSU}(n, \mathbb{C})$.

- Formally, variations of a minimal map $h : \tilde{S} \rightarrow (G/K, \nu)$ are sections $\dot{h} \in \Gamma_S(h^* TG/K)$. We write " $h_t = h + t\dot{h}$."
- The index of h , $\mathrm{Ind}(h)$, is the maximal dimension of a subspace of $\Gamma_S(h^* TG/K)$ on which

$$\frac{d^2}{dt^2} \Big|_{t=0} A(h + t\dot{h}) < 0.$$

- For $\tau : S_B \rightarrow S$ and $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$, $\mathrm{Ind}_\tau(f)$ is the maximal dimension of a subspace of $\Gamma_{S_B}(f^* T\mathbb{R}^{n-1})$ consisting of $\mathrm{Deck}(S_B/S)$ -invariant variations on which

$$\frac{d^2}{dt^2} \Big|_{t=0} A(f + t\dot{f}) < 0.$$

The index of minimal maps

Set $G = \mathrm{PSL}(n, \mathbb{C})$, $K = \mathrm{PSU}(n, \mathbb{C})$.

- Formally, variations of a minimal map $h : \tilde{S} \rightarrow (G/K, \nu)$ are sections $\dot{h} \in \Gamma(h^* TG/K)$. We write " $h_t = h + t\dot{h}$."
- The index of h , $\mathrm{Ind}(h)$, is the maximal dimension of a subspace of $\Gamma(h^* TG/K)$ on which

$$\frac{d^2}{dt^2} \Big|_{t=0} A(h + t\dot{h}) < 0.$$

- For $\tau : S_B \rightarrow S$ and $f : \tilde{S}_B \rightarrow \mathbb{R}^{n-1}$, $\mathrm{Ind}_\tau(f)$ is the maximal dimension of a subspace of $\Gamma(f^* T\mathbb{R}^{n-1})$ consisting of $\mathrm{Deck}(S_B/S)$ -invariant variations on which

$$\frac{d^2}{dt^2} \Big|_{t=0} A(f + t\dot{f}) < 0.$$

Theorem (S.-Smillie 2022)

$$\liminf_{R \rightarrow \infty} \mathrm{Ind}(h_R) \geq \mathrm{Ind}(f).$$

- For $n \geq 4$ and $g \geq 3$, we can choose an unstable minimal map $f : \tilde{S} \rightarrow \mathbb{R}^{n-1}$.

Disproof of the Labourie conjecture

- For $n \geq 4$ and $g \geq 3$, we can choose an unstable minimal map $f : \tilde{S} \rightarrow \mathbb{R}^{n-1}$.
- Choosing a reasonable identification $\mathbb{R}^{n-1} \rightarrow \mathbb{C}^{n-1} \rightarrow \mathfrak{t}^{\mathbb{C}}$, we turn f into a point in the Hitchin base (with trivial cameral cover).

- For $n \geq 4$ and $g \geq 3$, we can choose an unstable minimal map $f : \tilde{S} \rightarrow \mathbb{R}^{n-1}$.
- Choosing a reasonable identification $\mathbb{R}^{n-1} \rightarrow \mathbb{C}^{n-1} \rightarrow \mathfrak{t}^{\mathbb{C}}$, we turn f into a point in the Hitchin base (with trivial cameral cover).
- Using the Hitchin section, we obtain a path of Hitchin representations $\rho_R : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with minimal maps $h_R : \tilde{S} \rightarrow \mathrm{PSL}(n, \mathbb{R})/\mathrm{PSO}(n, \mathbb{R})$ such that

$$\liminf_{R \rightarrow \infty} \mathrm{Ind}(h_R) \geq \mathrm{Ind}(f).$$

- For $n \geq 4$ and $g \geq 3$, we can choose an unstable minimal map $f : \tilde{S} \rightarrow \mathbb{R}^{n-1}$.
- Choosing a reasonable identification $\mathbb{R}^{n-1} \rightarrow \mathbb{C}^{n-1} \rightarrow \mathfrak{t}^{\mathbb{C}}$, we turn f into a point in the Hitchin base (with trivial cameral cover).
- Using the Hitchin section, we obtain a path of Hitchin representations $\rho_R : \pi_1(\Sigma_g) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ with minimal maps $h_R : \tilde{S} \rightarrow \mathrm{PSL}(n, \mathbb{R})/\mathrm{PSO}(n, \mathbb{R})$ such that

$$\liminf_{R \rightarrow \infty} \mathrm{Ind}(h_R) \geq \mathrm{Ind}(f).$$

- For R sufficiently large, h_R is unstable, and hence ρ_R admits at least two minimal surfaces.

- For $\mathrm{PSL}(2, \mathbb{R})^n$, we can show $\mathrm{Ind}(h_R)$ is non-decreasing with R and $\lim_{R \rightarrow \infty} \mathrm{Ind}(h_R) = \mathrm{Ind}(f)$ (Marković-S.-Smillie).

- For $\mathrm{PSL}(2, \mathbb{R})^n$, we can show $\mathrm{Ind}(h_R)$ is non-decreasing with R and $\lim_{R \rightarrow \infty} \mathrm{Ind}(h_R) = \mathrm{Ind}(f)$ (Marković-S.-Smillie).
- For more general G , we have Hitchin rays such that $\mathrm{Ind}(h_R) > \mathrm{Ind}(f)$, although we have no example in the Hitchin component.

Question

What more can we say about $\mathrm{Ind}(h_R)$ in terms of $\mathrm{Ind}(f)$?

Thanks for listening