

Equivariant Log-concavity and Equivariant Kahler Packages

(or: Shadows of Hodge Theory)

Tao Gui

joint with Rui Xiong(熊锐)

YMSC Topology Seminar

2022/9/27

- ① Kähler packages
- ② Equivariant log-concavity
- ③ Equivariant Kähler packages

- 1 Kähler packages
- 2 Equivariant log-concavity
- 3 Equivariant Kähler packages

What is a Kähler package?

What is a Kähler package?

- X : mathematical object of "complex dimension" d

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

We expect that $(H^*, P(-, -), L)$ satisfy:

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

We expect that $(H^*, P(-, -), L)$ satisfy:

- Poincaré duality: $P(-, -)$ is non-degenerate.

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

We expect that $(H^*, P(-, -), L)$ satisfy:

- Poincaré duality: $P(-, -)$ is non-degenerate.
- Hard Lefschetz ("Théorème de Lefschetz vache"):

$$\forall k \leq d, \quad L^{d-k} : H^k(X) \xrightarrow{\sim} H^{2d-k}(X).$$

What is a Kähler package?

- X : mathematical object of "complex dimension" d
- $H^*(X) = \bigoplus_{k=0}^{2d} H^k(X)$ "cohomology" of X
- $P(-, -) : H^k(X) \times H^{2d-k}(X) \rightarrow \mathbb{R}, \forall k \leq d$ "Poincaré pairing"
- $L : H^k(X) \rightarrow H^{k+2}(X)$ "Lefschetz operator"

We expect that $(H^*, P(-, -), L)$ satisfy:

- Poincaré duality: $P(-, -)$ is non-degenerate.
- Hard Lefschetz ("Théorème de Lefschetz vache"):

$$\forall k \leq d, \quad L^{d-k} : H^k(X) \xrightarrow{\sim} H^{2d-k}(X).$$


- Hodge–Riemann relations: the bilinear form

$$\langle -, - \rangle_L = P(-, L^{d-k} -) : H^k(X) \times H^k(X) \longrightarrow \mathbb{R}$$

is $(-1)^k$ -definite on the kernel of L^{d-k+1} .

A picture to keep in mind

A picture to keep in mind



Kähler packages are both ubiquitous and fundamental

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold
(Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
 \Rightarrow Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) \Rightarrow Decomposition theorem

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) ⇒ Decomposition theorem
- Polytope (Stanley'80, Bressler-Lunts'03, Karu'04, et. al.)
⇒ McMullen g-conjecture'80, Aleksandrov-Fenchel inequality

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) ⇒ Decomposition theorem
- Polytope (Stanley'80, Bressler-Lunts'03, Karu'04, et. al.)
⇒ McMullen g-conjecture'80, Aleksandrov-Fenchel inequality
- Coxeter group (Soergel'92, Elias-Williamson'14, Williamson'16)
⇒ Kazhdan-Lusztig conjectures'79, Jantzen conjectures'79

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) ⇒ Decomposition theorem
- Polytope (Stanley'80, Bressler-Lunts'03, Karu'04, et. al.)
⇒ McMullen g-conjecture'80, Aleksandrov-Fenchel inequality
- Coxeter group (Soergel'92, Elias-Williamson'14, Williamson'16)
⇒ Kazhdan-Lusztig conjectures'79, Jantzen conjectures'79
- **Matroid (Huh and his collaborators)**
⇒ **Various combinatorial conjectures**

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) ⇒ Decomposition theorem
- Polytope (Stanley'80, Bressler-Lunts'03, Karu'04, et. al.)
⇒ McMullen g-conjecture'80, Aleksandrov-Fenchel inequality
- Coxeter group (Soergel'92, Elias-Williamson'14, Williamson'16)
⇒ Kazhdan-Lusztig conjectures'79, Jantzen conjectures'79
- Matroid (Huh and his collaborators)
⇒ Various combinatorial conjectures

Kähler packages are both ubiquitous and fundamental

- Smooth complex projective variety/ compact Kähler manifold (Poincaré'1895, Lefschetz'24, Hodge'30s, Chern'51, et. al.)
- Smooth projective variety over \mathbb{F}_q (Grothendieck'68)
⇒ Weil conjectures'49
- Singular projective variety (Goresky-MacPherson'80, Beilinson-Bernstein-Deligne'82, et. al.) ⇒ Decomposition theorem
- Polytope (Stanley'80, Bressler-Lunts'03, Karu'04, et. al.)
⇒ McMullen g-conjecture'80, Aleksandrov-Fenchel inequality
- Coxeter group (Soergel'92, Elias-Williamson'14, Williamson'16)
⇒ Kazhdan-Lusztig conjectures'79, Jantzen conjectures'79
- Matroid (Huh and his collaborators)
⇒ Various combinatorial conjectures

According to Geordie Williamson, this is "A mystery for the 21st century!"

Where and how can we find a Kähler package?

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)
- Hard Lefschetz: "Betti numbers" are unimodal in parity! (Stanley)

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)
- Hard Lefschetz: "Betti numbers" are unimodal in parity! (Stanley)
- Hodge–Riemann relations: "intersection numbers" are log-concave! (Huh)

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)
- Hard Lefschetz: "Betti numbers" are unimodal in parity! (Stanley)
- Hodge–Riemann relations: "intersection numbers" are log-concave! (Huh)

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)
- Hard Lefschetz: "Betti numbers" are unimodal in parity! (Stanley)
- Hodge–Riemann relations: "intersection numbers" are log-concave! (Huh)

Why? Essentially because

$$b^2 \geq ac \text{ if and only if } \begin{vmatrix} a & b \\ b & c \end{vmatrix} \leq 0.$$

Where and how can we find a Kähler package?

Kähler packages have some numerical shadows:

- Poincaré duality: "Betti numbers" are symmetric! (Poincaré)
- Hard Lefschetz: "Betti numbers" are unimodal in parity! (Stanley)
- Hodge–Riemann relations: "intersection numbers" are log-concave! (Huh)

Why? Essentially because

$$b^2 \geq ac \text{ if and only if } \begin{vmatrix} a & b \\ b & c \end{vmatrix} \leq 0.$$

Miracle: in some cases,
"Betti numbers" = "intersection numbers"!

- ① Kähler packages
- ② Equivariant log-concavity
- ③ Equivariant Kähler packages

Log-concavity

- A sequence b_0, b_1, b_2, \dots of non-negative real numbers is called *log-concave* if, for all i ,

$$b_i^2 \geq b_{i-1} b_{i+1}.$$

Log-concavity

- A sequence b_0, b_1, b_2, \dots of non-negative real numbers is called *log-concave* if, for all i ,

$$b_i^2 \geq b_{i-1} b_{i+1}.$$

- A polynomial $f(t) = \sum b_i t^i$ with non-negative coefficients is called log-concave if its sequence of coefficients is log-concave.

Log-concavity

- A sequence b_0, b_1, b_2, \dots of non-negative real numbers is called *log-concave* if, for all i ,

$$b_i^2 \geq b_{i-1} b_{i+1}.$$

- A polynomial $f(t) = \sum b_i t^i$ with non-negative coefficients is called log-concave if its sequence of coefficients is log-concave.
- A sequence b_0, b_1, b_2, \dots or a polynomial $f(t) = \sum b_i t^i$ is *strongly log-concave* if, for all $i \leq j \leq k \leq l$ with $i + l = j + k$, we have

$$b_j b_k \geq b_i b_l.$$

Log-concavity

- A sequence b_0, b_1, b_2, \dots of non-negative real numbers is called *log-concave* if, for all i ,

$$b_i^2 \geq b_{i-1} b_{i+1}.$$

- A polynomial $f(t) = \sum b_i t^i$ with non-negative coefficients is called log-concave if its sequence of coefficients is log-concave.
- A sequence b_0, b_1, b_2, \dots or a polynomial $f(t) = \sum b_i t^i$ is *strongly log-concave* if, for all $i \leq j \leq k \leq l$ with $i + l = j + k$, we have

$$b_j b_k \geq b_i b_l.$$

- It can be shown that a sequence or polynomial is strongly log-concave if and only if it is log-concave with *no internal zeros*.

hyperplane arrangement

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V
- Consider the complement

$$U(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$$

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V
- Consider the complement

$$U(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$$

- $b_i(\mathcal{A}) := \dim H^i(U(\mathcal{A}); \mathbb{Q})$ Betti numbers

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V
- Consider the complement

$$U(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$$

- $b_i(\mathcal{A}) := \dim H^i(U(\mathcal{A}); \mathbb{Q})$ Betti numbers
- $\pi_{\mathcal{A}}(t) := \sum_{i \geq 0} b_i(\mathcal{A}) t^i$ Poincaré polynomial

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V
- Consider the complement

$$U(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$$

- $b_i(\mathcal{A}) := \dim H^i(U(\mathcal{A}); \mathbb{Q})$ Betti numbers
- $\pi_{\mathcal{A}}(t) := \sum_{i \geq 0} b_i(\mathcal{A}) t^i$ Poincaré polynomial

hyperplane arrangement

- V : finite dimensional vector space over \mathbb{C}
- \mathcal{A} : finite collection of central hyperplanes in V
- Consider the complement

$$U(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$$

- $b_i(\mathcal{A}) := \dim H^i(U(\mathcal{A}); \mathbb{Q})$ Betti numbers
- $\pi_{\mathcal{A}}(t) := \sum_{i \geq 0} b_i(\mathcal{A}) t^i$ Poincaré polynomial

Example: Ordered configuration space / Coxeter arrangement

Let $\mathcal{A}_n := \{H_{ij}\}$, where $H_{ij} := \{z \in \mathbb{C}^n \mid z_i = z_j\} \subset \mathbb{C}^n$. Then

$$U(\mathcal{A}_n) = \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_i \neq z_j, \forall i \neq j\} =: \text{Conf}(n, \mathbb{C}),$$

and (exercise) $\pi_{\mathcal{A}_n}(t) = \prod_{k=1}^{n-1} (1 + kt)$.

Thm (Hoggar conj '74 [Huh, 2012])

For any \mathcal{A} , $\pi_{\mathcal{A}}(t)$ is strongly log-concave.

Thm (Hoggar conj '74 [Huh, 2012])

For any \mathcal{A} , $\pi_{\mathcal{A}}(t)$ is strongly log-concave.

Any arrangement \mathcal{A} determines a matroid $M_{\mathcal{A}}$.

Given a matroid M , one can define the Orlik-Solomon algebra $\text{OS}^*(M)$, and there is a canonical isomorphism

$$\text{OS}^*(M_{\mathcal{A}}) \cong H^*(U(\mathcal{A}); \mathbb{Z}).$$

Let

$$\pi_M(t) := \sum_{i \geq 0} t^i \dim \text{OS}^i(M).$$

Thm (Hoggar conj '74 [Huh, 2012])

For any \mathcal{A} , $\pi_{\mathcal{A}}(t)$ is strongly log-concave.

Any arrangement \mathcal{A} determines a matroid $M_{\mathcal{A}}$.

Given a matroid M , one can define the Orlik-Solomon algebra $\text{OS}^*(M)$, and there is a canonical isomorphism

$$\text{OS}^*(M_{\mathcal{A}}) \cong H^*(U(\mathcal{A}); \mathbb{Z}).$$

Let

$$\pi_M(t) := \sum_{i \geq 0} t^i \dim \text{OS}^i(M).$$

Thm (Heron-Rota-Welsh conj '70s [AHK, 2018])

For any matroid M , $\pi_M(t)$ is strongly log-concave.

Equivariant log-concavity

Equivariant log-concavity

- Let G be a finite group, and let

$$V = \bigoplus_{i \geq 0} V^i$$

be a graded representation of G with V^i finite dimensional.

Equivariant log-concavity

- Let G be a finite group, and let

$$V = \bigoplus_{i \geq 0} V^i$$

be a graded representation of G with V^i finite dimensional.

- We say that V is *(weakly) log-concave* if, for all i .

$V^i \otimes V^i$ contains a subrepresentation isomorphic to $V^{i-1} \otimes V^{i+1}$.

Equivariant log-concavity

- Let G be a finite group, and let

$$V = \bigoplus_{i \geq 0} V^i$$

be a graded representation of G with V^i finite dimensional.

- We say that V is (*weakly*) *log-concave* if, for all i .

$V^i \otimes V^i$ contains a subrepresentation isomorphic to $V^{i-1} \otimes V^{i+1}$.

- V is *strongly equivariantly log-concave* if, for all $i \leq j \leq k \leq l$ with $i + l = j + k$,

$V^j \otimes V^k$ contains a subrepresentation isomorphic to $V^i \otimes V^l$.

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,
- \mathcal{A} : finite collection of hyperplanes in V .

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,
- \mathcal{A} : finite collection of hyperplanes in V .
- G acts linearly on V , preserving $\mathcal{A} \rightsquigarrow$ action of G on $U(\mathcal{A})$.

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,
- \mathcal{A} : finite collection of hyperplanes in V .
- G acts linearly on V , preserving $\mathcal{A} \rightsquigarrow$ action of G on $U(\mathcal{A})$.

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,
- \mathcal{A} : finite collection of hyperplanes in V .
- G acts linearly on V , preserving $\mathcal{A} \rightsquigarrow$ action of G on $U(\mathcal{A})$.

Example

The symmetric group S_n acts on \mathbb{C}^n preserving \mathcal{A}_n . this induces an action on $U(\mathcal{A}_n) = \text{Conf}(n, \mathbb{C})$ by permuting labels of the points.

Equivariant log-concavity conjecture

- V : finite dimensional vector space over \mathbb{C} ,
- \mathcal{A} : finite collection of hyperplanes in V .
- G acts linearly on V , preserving $\mathcal{A} \rightsquigarrow$ action of G on $U(\mathcal{A})$.

Example

The symmetric group S_n acts on \mathbb{C}^n preserving \mathcal{A}_n . this induces an action on $U(\mathcal{A}_n) = \text{Conf}(n, \mathbb{C})$ by permuting labels of the points.

Conjecture (Matherne-Miyata-Proudfoot-Ramos, 2021)

The cohomology ring $H^*(U(\mathcal{A}); \mathbb{Q})$ is a strongly equivariantly log-concave graded representation of G .

More generally, if M is a matroid with an action of G , the Orlik-Solomon algebra $OS^*(M)$ is a strongly equivariantly log-concave graded representation of G .

Flag variety

Flag variety

- \mathcal{F}_n : (full) flag variety over \mathbb{C} , parametrizing all complete flags in a n -dimensional \mathbb{C} -vector space V

$$\{0\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_n = V$$

Flag variety

- \mathcal{F}_n : (full) flag variety over \mathbb{C} , parametrizing all complete flags in a n -dimensional \mathbb{C} -vector space V

$$\{0\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_n = V$$

- $\mathcal{F}_n \cong \mathrm{GL}(n, \mathbb{C})/B_n \Rightarrow \dim \mathcal{F}_n = n(n-1)/2$

Flag variety

- \mathcal{F}_n : (full) flag variety over \mathbb{C} , parametrizing all complete flags in a n -dimensional \mathbb{C} -vector space V

$$\{0\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_n = V$$

- $\mathcal{F}_n \cong \mathrm{GL}(n, \mathbb{C})/B_n \Rightarrow \dim \mathcal{F}_n = n(n-1)/2$

Flag variety

- \mathcal{F}_n : (full) flag variety over \mathbb{C} , parametrizing all complete flags in a n -dimensional \mathbb{C} -vector space V

$$\{0\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_n = V$$

- $\mathcal{F}_n \cong \mathrm{GL}(n, \mathbb{C})/B_n \Rightarrow \dim \mathcal{F}_n = n(n-1)/2$

Theorem (Borel, 1935) "Borel's picture"

As a graded ring,

$$H^*(\mathcal{F}_n, \mathbb{Z}) \cong \mathbb{Z}[t_1, \dots, t_n] / (\sigma_1, \dots, \sigma_n),$$

where the t_j 's are of degree 2 and the σ_i 's are the elementary symmetric polynomials in the variables t_j 's.

Equivariant log-concavity conjecture on flag variety

As a corollary, $P_{\mathcal{F}_n}(q) := \sum_i b_{2i} q^i = \prod_{k=0}^{n-1} (1 + q + \cdots + q^k)$, which is a symmetric, unimodal, and log-concave polynomial.

Equivariant log-concavity conjecture on flag variety

As a corollary, $P_{\mathcal{F}_n}(q) := \sum_i b_{2i} q^i = \prod_{k=0}^{n-1} (1 + q + \cdots + q^k)$, which is a symmetric, unimodal, and log-concave polynomial.

Conj (G., Conjecture 1 in arXiv:2205.05408)

For all integer $n \geq 1$, the cohomology ring of the flag variety is equivariantly log-concave as graded representation of S_n .

Equivariant log-concavity conjecture on flag variety

As a corollary, $P_{\mathcal{F}_n}(q) := \sum_i b_{2i} q^i = \prod_{k=0}^{n-1} (1 + q + \cdots + q^k)$, which is a symmetric, unimodal, and log-concave polynomial.

Conj (G., Conjecture 1 in arXiv:2205.05408)

For all integer $n \geq 1$, the cohomology ring of the flag variety is equivariantly log-concave as graded representation of S_n .

In terms of Kronecker coefficients, it is equivalent to

Conj (G., Conjecture 5 in arXiv:2205.05408)

For all natural number n and $1 \leq i \leq n(n-1)/2 - 1$,

$$\sum_{\lambda \vdash n} \sum_{\mu \vdash n} b_{\lambda, i-1} b_{\mu, i+1} g_{\lambda\mu}^\nu \leq \sum_{\lambda \vdash n} \sum_{\mu \vdash n} b_{\lambda, i} b_{\mu, i} g_{\lambda\mu}^\nu$$

hold for all $\nu \vdash n$, where $g_{\lambda\mu}^\nu$ are the Kronecker coefficients, and $b_{\lambda, i}$ are the "fake degrees".

Using the representation stability theory, we have the following

Theorem (G.)

Above conjectures hold for degree ≤ 3 and co-degree ≤ 3 , i.e., $\forall n$:

$$\begin{aligned} (H^{2*1})^{\otimes 2} &\supseteq H^{2*0} \otimes H^{2*2}, \quad \left(H^{2*\left(\binom{n}{2}-1\right)}\right)^{\otimes 2} \supseteq H^{2*\binom{n}{2}} \otimes H^{2*\left(\binom{n}{2}-2\right)}, \\ (H^{2*2})^{\otimes 2} &\supseteq H^{2*1} \otimes H^{2*3}, \quad \left(H^{2*\left(\binom{n}{2}-2\right)}\right)^{\otimes 2} \supseteq H^{2*\left(\binom{n}{2}-1\right)} \otimes H^{2*\left(\binom{n}{2}-3\right)}, \\ (H^{2*3})^{\otimes 2} &\supseteq H^{2*2} \otimes H^{2*4}, \quad \left(H^{2*\left(\binom{n}{2}-3\right)}\right)^{\otimes 2} \supseteq H^{2*\left(\binom{n}{2}-2\right)} \otimes H^{2*\left(\binom{n}{2}-4\right)}. \end{aligned}$$

Using the representation stability theory, we have the following

Theorem (G.)

Above conjectures hold for degree ≤ 3 and co-degree ≤ 3 , i.e., $\forall n$:

$$\begin{aligned} (H^{2*1})^{\otimes 2} &\supseteq H^{2*0} \otimes H^{2*2}, \left(H^{2*\left(\binom{n}{2}-1\right)}\right)^{\otimes 2} \supseteq H^{2*\binom{n}{2}} \otimes H^{2*\left(\binom{n}{2}-2\right)}, \\ (H^{2*2})^{\otimes 2} &\supseteq H^{2*1} \otimes H^{2*3}, \left(H^{2*\left(\binom{n}{2}-2\right)}\right)^{\otimes 2} \supseteq H^{2*\left(\binom{n}{2}-1\right)} \otimes H^{2*\left(\binom{n}{2}-3\right)}, \\ (H^{2*3})^{\otimes 2} &\supseteq H^{2*2} \otimes H^{2*4}, \left(H^{2*\left(\binom{n}{2}-3\right)}\right)^{\otimes 2} \supseteq H^{2*\left(\binom{n}{2}-2\right)} \otimes H^{2*\left(\binom{n}{2}-4\right)}. \end{aligned}$$

The numerical evidence suggests the stronger conjecture

Conj (G., Conjecture 7 in arXiv:2205.05408)

For all natural number n , $\nu \vdash n$, and $1 \leq i \leq n(n-1)/2 - 1$, the "difference" sequence $d_{\nu,i}$ is symmetric and unimodal, where

$$d_{\nu,i} := \sum_{\lambda \vdash n} \sum_{\mu \vdash n} b_{\lambda,i} b_{\mu,i} g_{\lambda\mu}^{\nu} - \sum_{\lambda \vdash n} \sum_{\mu \vdash n} b_{\lambda,i-1} b_{\mu,i+1} g_{\lambda\mu}^{\nu}.$$

Why equivariant?

Why equivariant?

- Our feeling: establishing equivariant log-concavity might be simpler than establishing log-concavity!

Why equivariant?

- Our feeling: establishing equivariant log-concavity might be simpler than establishing log-concavity!
- Example: a general endomorphism on a vector space is an isomorphism, however establishing a specific one might be very difficult.

Why equivariant?

- Our feeling: establishing equivariant log-concavity might be simpler than establishing log-concavity!
- Example: a general endomorphism on a vector space is an isomorphism, however establishing a specific one might be very difficult.
- But *symmetry reduces choices!*

Why equivariant?

- Our feeling: establishing equivariant log-concavity might be simpler than establishing log-concavity!
- Example: a general endomorphism on a vector space is an isomorphism, however establishing a specific one might be very difficult.
- But *symmetry reduces choices!*
- Our idea: *lift the "1-dimensional" sequence to a "2-dimensional" array, and establishing the (equivariant) hard Lefschetz theorem for each "anti-diagonal", which implies the strongly (equivariant) log-concavity of the original sequence.*

An illustration

$$b_0, b_1, \dots, b_k, \dots$$

$$\downarrow$$

$$\forall k:$$

$$b_0 b_k \leq b_1 b_{k-1} \leq \dots \geq b_{k-1} b_1 \geq b_k b_0$$

$$\Updownarrow$$

$$H^* = \bigoplus_k H^k$$

$$\downarrow$$

$$\forall k:$$

$$H^0 \otimes H^k \hookrightarrow H^1 \otimes H^{k-1} \hookrightarrow \dots \twoheadrightarrow H^{k-1} \otimes H^1 \twoheadrightarrow H^k \otimes H^0$$

$$\Updownarrow$$


Illustration

$$b_0, b_1, \dots, b_k, \dots$$



$\forall k:$

$$b_0 b_k \leq b_1 b_{k-1} \leq \dots \leq b_{k-1} b_1 \geq b_k b_0$$




$$H^* = \bigoplus_k H^k$$



$\forall k:$

$$H^0 \otimes H^k \hookrightarrow H^1 \otimes H^{k-1} \hookrightarrow \dots \hookrightarrow H^{k-1} \otimes H^1 \rightarrow H^k \otimes H^0$$



- ① Kähler packages
- ② Equivariant log-concavity
- ③ Equivariant Kähler packages

Equivariant Kähler packages

In attacking the above conjectures, we discover two new Kähler packages that are *equivariant* and have *no geometric origin*. The equivariant log-concavity hints at our discoveries for these structures.

Polynomial rings

Polynomial rings

- Consider $S_n \curvearrowright B(S^1)^n = (\mathbb{CP}^\infty)^n$, the classifying space
 $\leadsto S_n \curvearrowright H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$

Polynomial rings

- Consider $S_n \curvearrowright B(S^1)^n = (\mathbb{CP}^\infty)^n$, the classifying space
 $\leadsto S_n \curvearrowright H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$
- Betti numbers $b_{2i} = \binom{n+i-1}{i}$, log-concave for fixed n .

Polynomial rings

- Consider $S_n \curvearrowright B(S^1)^n = (\mathbb{CP}^\infty)^n$, the classifying space
 $\leadsto S_n \curvearrowright H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$
- Betti numbers $b_{2i} = \binom{n+i-1}{i}$, log-concave for fixed n .
- Fix a pair (n, m) , consider the graded \mathbb{R} -vector space

$$H_{n,m} = \bigoplus_{i=0}^m H_{n,m}^{-m+2i}, \text{ with } H_{n,m}^{-m+2i} := D^i \otimes R^{m-i},$$

where $D = \mathbb{R}[d_1, \dots, d_n]$, $R = \mathbb{R}[x_1, \dots, x_n]$

Polynomial rings

- Consider $S_n \curvearrowright B(S^1)^n = (\mathbb{CP}^\infty)^n$, the classifying space
 $\leadsto S_n \curvearrowright H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$
- Betti numbers $b_{2i} = \binom{n+i-1}{i}$, log-concave for fixed n .
- Fix a pair (n, m) , consider the graded \mathbb{R} -vector space

$$H_{n,m} = \bigoplus_{i=0}^m H_{n,m}^{-m+2i}, \text{ with } H_{n,m}^{-m+2i} := D^i \otimes R^{m-i},$$

where $D = \mathbb{R}[d_1, \dots, d_n]$, $R = \mathbb{R}[x_1, \dots, x_n]$

- Define a pairing $\langle -, - \rangle$ on $H_{n,m}$ by interpreting the d_i as differential operators $\frac{\partial}{\partial x_i}$ “homology-cohomology pairing”

Polynomial rings

- Consider $S_n \curvearrowright B(S^1)^n = (\mathbb{CP}^\infty)^n$, the classifying space
 $\leadsto S_n \curvearrowright H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$
- Betti numbers $b_{2i} = \binom{n+i-1}{i}$, log-concave for fixed n .
- Fix a pair (n, m) , consider the graded \mathbb{R} -vector space

$$H_{n,m} = \bigoplus_{i=0}^m H_{n,m}^{-m+2i}, \text{ with } H_{n,m}^{-m+2i} := D^i \otimes R^{m-i},$$

where $D = \mathbb{R}[d_1, \dots, d_n]$, $R = \mathbb{R}[x_1, \dots, x_n]$

- Define a pairing $\langle -, - \rangle$ on $H_{n,m}$ by interpreting the d_i as differential operators $\frac{\partial}{\partial x_i}$ “homology-cohomology pairing”
- Define $L : H_{n,m}^i \longrightarrow H_{n,m}^{i+2}$ to be the linear map

$$L := \sum_{i=1}^n d_i \otimes \frac{\partial}{\partial x_i},$$

where d_i acts by multiplication.

Theorem (G.-Xiong)

For any pair of natural numbers (n, m) , we have

- (a) (PD) The bilinear pairing $\langle -, - \rangle : H_{n,m} \times H_{n,m} \longrightarrow \mathbb{R}$ is S_n -equivariant and graded non-degenerate;
- (b) (HL) $L : H_{n,m}^i \longrightarrow H_{n,m}^{i+2}$ is S_n -equivariant and satisfies the hard Lefschetz theorem;
- (c) (HR) For all $0 \leq i \leq m/2$, the bilinear form

$$(a, b)_L^{-m+2i} = \langle a, L^{m-2i} b \rangle : H_{n,m}^{-m+2i} \times H_{n,m}^{-m+2i} \longrightarrow \mathbb{R}$$

is S_n -equivariant and $(-1)^i$ -definite on $\ker (L^{m-2i+1})$.

Theorem (G.-Xiong)

For any pair of natural numbers (n, m) , we have

- (a) (PD) The bilinear pairing $\langle -, - \rangle : H_{n,m} \times H_{n,m} \longrightarrow \mathbb{R}$ is S_n -equivariant and graded non-degenerate;
- (b) (HL) $L : H_{n,m}^i \longrightarrow H_{n,m}^{i+2}$ is S_n -equivariant and satisfies the hard Lefschetz theorem;
- (c) (HR) For all $0 \leq i \leq m/2$, the bilinear form

$$(a, b)_L^{-m+2i} = \langle a, L^{m-2i} b \rangle : H_{n,m}^{-m+2i} \times H_{n,m}^{-m+2i} \longrightarrow \mathbb{R}$$

is S_n -equivariant and $(-1)^i$ -definite on $\ker(L^{m-2i+1})$.

Corollary

$H^*((\mathbb{CP}^\infty)^n, \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$ is strongly equivariantly log-concave and the inclusions of S_n -representations are given by the operator

$$L = \sum_{i=1}^n t_i \otimes \frac{\partial}{\partial t_i}.$$

Exterior algebras

Exterior algebras

- $S_n \curvearrowright (S^1)^n \leadsto S_n \curvearrowright H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}}[\alpha_1, \dots, \alpha_n]$

Exterior algebras

- $S_n \curvearrowright (S^1)^n \leadsto S_n \curvearrowright H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}}[\alpha_1, \dots, \alpha_n]$
- Betti numbers $b_{2i} = \binom{n}{i}$, log-concave for fixed n

Exterior algebras

- $S_n \curvearrowright (S^1)^n \leadsto S_n \curvearrowright H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}}[\alpha_1, \dots, \alpha_n]$
- Betti numbers $b_{2i} = \binom{n}{i}$, log-concave for fixed n
- Fix a pair (n, m) with $m \leq 2n$, consider

$$H'_{n,m} = \bigoplus_{i=0}^m (H'_{n,m})^{-m+2i}, \text{ with } (H'_{n,m})^{-m+2i} := \Lambda^i \otimes (\Lambda^*)^{m-i},$$

where $\Lambda = \Lambda(V) = \Lambda_{\mathbb{R}}[\theta_1, \dots, \theta_n]$, $\Lambda^* = \Lambda(V^*)$

Exterior algebras

- $S_n \curvearrowright (S^1)^n \rightsquigarrow S_n \curvearrowright H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}}[\alpha_1, \dots, \alpha_n]$
- Betti numbers $b_{2i} = \binom{n}{i}$, log-concave for fixed n
- Fix a pair (n, m) with $m \leq 2n$, consider

$$H'_{n,m} = \bigoplus_{i=0}^m (H'_{n,m})^{-m+2i}, \text{ with } (H'_{n,m})^{-m+2i} := \Lambda^i \otimes (\Lambda^*)^{m-i},$$

where $\Lambda = \Lambda(V) = \Lambda_{\mathbb{R}}[\theta_1, \dots, \theta_n]$, $\Lambda^* = \Lambda(V^*)$

- Define a pairing $\langle -, - \rangle$ on $H'_{n,m}$ by the natural duality between V and V^* “homology-cohomology pairing”

Exterior algebras

- $S_n \curvearrowright (S^1)^n \rightsquigarrow S_n \curvearrowright H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}} [\alpha_1, \dots, \alpha_n]$
- Betti numbers $b_{2i} = \binom{n}{i}$, log-concave for fixed n
- Fix a pair (n, m) with $m \leq 2n$, consider

$$H'_{n,m} = \bigoplus_{i=0}^m (H'_{n,m})^{-m+2i}, \text{ with } (H'_{n,m})^{-m+2i} := \Lambda^i \otimes (\Lambda^*)^{m-i},$$

where $\Lambda = \Lambda(V) = \Lambda_{\mathbb{R}} [\theta_1, \dots, \theta_n]$, $\Lambda^* = \Lambda(V^*)$

- Define a pairing $\langle -, - \rangle$ on $H'_{n,m}$ by the natural duality between V and V^* “homology-cohomology pairing”
- Define

$$L : (H'_{n,m})^i \longrightarrow (H'_{n,m})^{i+2}$$

to be the linear map

$$L := \sum_{k=1}^n e_{\theta_k} \otimes i_{\theta_k}.$$

Theorem (G.-Xiong)

For any pair of natural numbers (n, m) with $m \leq 2n$, we have S_n -equivariant PD, HL and HR for $H'_{n,m}$.

Theorem (G.-Xiong)

For any pair of natural numbers (n, m) with $m \leq 2n$, we have S_n -equivariant PD, HL and HR for $H'_{n,m}$.

Corollary

$H^*((S^1)^n, \mathbb{R}) = \Lambda_{\mathbb{R}}[\alpha_1, \dots, \alpha_n]$ is strongly equivariantly log-concave and the inclusions of S_n -representations are given by the operator

$$L = \sum_{k=1}^n e_{\alpha_k} \otimes i_{\alpha_k^*}.$$

where e_{α_k} and $i_{\alpha_k^*}$ are the exterior product and interior product, respectively, and α_k^* is the linear functional on $V = \langle \alpha_1, \dots, \alpha_n \rangle$

satisfying $\alpha_k^*(\alpha_l) = \delta_{kl} = \begin{cases} 0, & \text{if } k \neq l, \\ 1, & \text{if } k = l. \end{cases}$

Some example of "Betti numbers"...

Some example of "Betti numbers"...

$\mathcal{R}[x_1, x_2]:$

4	4					
3	3	6				
2	2	4	6			
1	1	2	3	4
	1	2	3	4		

$\mathcal{R}[x_1, x_2, x_3]:$

10	10						
6	6	18					
3	3	9	18				
1	1	3	6	10	
	1	3	6	10			

$\wedge[\omega_1, \omega_2]:$

1	1	2	1
2	2	4	2
1	1	2	1
	1	2	1

$\wedge[\omega_1, \omega_2, \omega_3]:$

1	1	3	3	1
3	3	9	9	3
3	3	9	9	3
1	1	3	3	1
	1	3	3	1

$\wedge[\omega_1, \omega_2, \omega_3, \omega_4]:$

1	1	4	6	4	1
4	4	16	24	16	4
6	6	24	36	24	6
4	4	16	24	16	4
1	1	4	6	4	1
	1	4	6	4	1

An "exotic" example

For the exterior algebra, we find that "usual" gradings on tensor products satisfy PD and HL but NOT satisfy HR.

An "exotic" example

For the exterior algebra, we find that "usual" gradings on tensor products satisfy PD and HL but NOT satisfy HR.

- Fix any n , consider the graded \mathbb{R} -vector space

$$H_n = \bigoplus_{i=0}^{2n} H_n^{-n+i}, \text{ with } H_n^{-n+i} := \bigoplus_{j+k=i} \Lambda^j \otimes (\Lambda^*)^k,$$

i.e., consider the "usual" gradings (up to a degree shift).

An "exotic" example

For the exterior algebra, we find that "usual" gradings on tensor products satisfy PD and HL but NOT satisfy HR.

- Fix any n , consider the graded \mathbb{R} -vector space

$$H_n = \bigoplus_{i=0}^{2n} H_n^{-n+i}, \text{ with } H_n^{-n+i} := \bigoplus_{j+k=i} \Lambda^j \otimes (\Lambda^*)^k,$$

i.e., consider the "usual" gradings (up to a degree shift).

- Define a pairing on H_n by the multiplication map

$$H_n^{-n+i} \otimes H_n^{n-i} \rightarrow H_n^n \cong \mathbb{R}.$$

An "exotic" example

For the exterior algebra, we find that "usual" gradings on tensor products satisfy PD and HL but NOT satisfy HR.

- Fix any n , consider the graded \mathbb{R} -vector space

$$H_n = \bigoplus_{i=0}^{2n} H_n^{-n+i}, \text{ with } H_n^{-n+i} := \bigoplus_{j+k=i} \Lambda^j \otimes (\Lambda^*)^k,$$

i.e., consider the "usual" gradings (up to a degree shift).

- Define a pairing on H_n by the multiplication map

$$H_n^{-n+i} \otimes H_n^{n-i} \rightarrow H_n^n \cong \mathbb{R}.$$

- We define

$$L : H_n^i \longrightarrow H_n^{i+2}$$

to be the linear map

$$L := \sum_{k=1}^n e_{\theta_k} \otimes e_{\xi_k}.$$

Theorem

For any natural numbers n , we have

- (a) (PD) The bilinear pairing $\langle -, - \rangle : H_n \times H_n \longrightarrow \mathbb{R}$ is S_n -equivariant and graded non-degenerate;
- (b) (HL) [Kim-Rhoades, 2020; G.-Xiong] $L : H_n^i \longrightarrow H_n^{i+2}$ is S_n -equivariant and satisfies the hard Lefschetz theorem;
- (c) [G.-Xiong] No HR, which can be seen by the signature!

Theorem

For any natural numbers n , we have

- (a) (PD) The bilinear pairing $\langle -, - \rangle : H_n \times H_n \longrightarrow \mathbb{R}$ is S_n -equivariant and graded non-degenerate;
- (b) (HL) [Kim-Rhoades, 2020; G.-Xiong] $L : H_n^i \longrightarrow H_n^{i+2}$ is S_n -equivariant and satisfies the hard Lefschetz theorem;
- (c) [G.-Xiong] No HR, which can be seen by the signature!

There is also an S_n -equivariant \mathfrak{sl}_2 -action on the "usual" gradings of the tensor product of the polynomial ring. But neither Poincaré duality nor hard Lefschetz holds because the "usual" gradings are infinite-dimensional.

Features of constructions

Features of constructions

- The constructions are purely algebraic and have no "geometric origin".

Features of constructions

- The constructions are purely algebraic and have no "geometric origin".
- The Poincaré pairing $\langle -, - \rangle$, the Lefschetz operator L , and the Lefschetz form $(a, b)_L^{-m+2i}$ are all S_n -equivariant, which is rare in Lefschetz theory.

Features of constructions

- The constructions are purely algebraic and have no "geometric origin".
- The Poincaré pairing $\langle -, - \rangle$, the Lefschetz operator L , and the Lefschetz form $(a, b)_L^{-m+2i}$ are all S_n -equivariant, which is rare in Lefschetz theory.
- The adjoint ("down") operator f of the Lefschetz operator L can be written explicitly. In other algebraic/combinatorial setting, direct constructions of f seems difficult.

Features of constructions

- The constructions are purely algebraic and have no "geometric origin".
- The Poincaré pairing $\langle -, - \rangle$, the Lefschetz operator L , and the Lefschetz form $(a, b)_L^{-m+2i}$ are all S_n -equivariant, which is rare in Lefschetz theory.
- The adjoint ("down") operator f of the Lefschetz operator L can be written explicitly. In other algebraic/combinatorial setting, direct constructions of f seems difficult.
- The proof of hard Lefschetz and Hodge–Riemann relations is simple/natural and it takes advantage of the geometry of the product of projective spaces.

Questions

I would like to finish with some questions:

Questions

I would like to finish with some questions:

- Is there any geometric interpretation of these constructions?

Questions

I would like to finish with some questions:

- Is there any geometric interpretation of these constructions?
- Do these equivariant Kähler packages, especially the Hodge–Riemann relations, have some implications for the diagonal coinvariant ring and the fermionic diagonal coinvariant ring?

Questions

I would like to finish with some questions:

- Is there any geometric interpretation of these constructions?
- Do these equivariant Kähler packages, especially the Hodge–Riemann relations, have some implications for the diagonal coinvariant ring and the fermionic diagonal coinvariant ring?
- It would be interesting to know whether our approach could shed some new light on other (equivariant) log-concavity questions and conjectures.

Questions

I would like to finish with some questions:

- Is there any geometric interpretation of these constructions?
- Do these equivariant Kähler packages, especially the Hodge–Riemann relations, have some implications for the diagonal coinvariant ring and the fermionic diagonal coinvariant ring?
- It would be interesting to know whether our approach could shed some new light on other (equivariant) log-concavity questions and conjectures.
- It is interesting to see whether similar (equivariant) Hodge-theoretic structures exist in the settings of other (equivariant) log-concavity conjectures.