

# Hyperbolic Geometry and Quantum Invariants

Tian Yang  
Texas A&M University

Qinghua University, December 21, 2021

## Part I: Overview

Two very different approaches to 3-manifolds:

Two very different approaches to 3-manifolds:

- ▶ Hyperbolic geometry

Two very different approaches to 3-manifolds:

- ▶ Hyperbolic geometry

Thurston, Perelman, Agol ...

Two very different approaches to 3-manifolds:

- ▶ Hyperbolic geometry

Thurston, Perelman, Agol ...

- ▶ Quantum invariants

Two very different approaches to 3-manifolds:

- ▶ Hyperbolic geometry

Thurston, Perelman, Agol ...

- ▶ Quantum invariants

Jones, Witten, Reshetikhin, Turaev ...

$K = \text{knot in } S^3$





$K = \text{knot in } S^3$



► Hyperbolic geometry

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the **volume**,

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the **volume**, the **Chern-Simons invariant**

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#)

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.



$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.
  - ▶ [Colored Jones polynomials](#)

$K = \text{knot in } S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.
  - ▶ **Colored Jones polynomials**  $J_n(K, t) \in \mathbb{Z}[t, t^{-1}]$ ,

$K$  = knot in  $S^3$



- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
  - ▶ Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.
  - ▶ **Colored Jones polynomials**  $J_n(K, t) \in \mathbb{Z}[t, t^{-1}]$ ,  $n \in \mathbb{N}$ .

$K$  = knot in  $S^3$



► Hyperbolic geometry

- Complete metric of constant curvature  $-1$  on  $S^3 \setminus K$ .
- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.

► Quantum invariants

- Constructed using representation theory of quantum groups.
- **Colored Jones polynomials**  $J_n(K, t) \in \mathbb{Z}[t, t^{-1}]$ ,  $n \in \mathbb{N}$ .

$$J_n(\text{trefoil}, t) = \sum_{j=0}^{n-1} \prod_{k=1}^j \left( t^{\frac{n-k}{2}} - t^{-\frac{n-k}{2}} \right) \left( t^{\frac{n+k}{2}} - t^{-\frac{n+k}{2}} \right)$$

## ► Volume Conjecture (Kashaev '97)

- ▶ **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot,

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement.

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$



- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$

- There are knots with same volume, but different colored Jones polynomials.

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$

- There are knots with same volume, but different colored Jones polynomials.



$5_2$  knot



$(-2, 3, 7)$ -pretzel knot

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$

- **Asymptotic Expansion Conjecture** (Witten '89, Gukov '05)

$$J_n(K, e^{\frac{2\pi i}{n}}) = \frac{n^{\frac{3}{2}}}{\sqrt{\text{Tor}(S^3 \setminus K)}} e^{\frac{n}{2\pi} (\text{Vol}(S^3 \setminus K) + i \text{CS}(S^3 \setminus K))} \left(1 + O\left(\frac{1}{n}\right)\right)$$

as  $n \rightarrow \infty$ ,

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$

- **Asymptotic Expansion Conjecture** (Witten '89, Gukov '05)

$$J_n(K, e^{\frac{2\pi i}{n}}) = \frac{n^{\frac{3}{2}}}{\sqrt{\text{Tor}(S^3 \setminus K)}} e^{\frac{n}{2\pi} (\text{Vol}(S^3 \setminus K) + i \text{CS}(S^3 \setminus K))} \left(1 + O\left(\frac{1}{n}\right)\right)$$

as  $n \rightarrow \infty$ , where  $\text{CS}(S^3 \setminus K)$  is the Chern-Simons invariant and  $\text{Tor}(S^3 \setminus K)$  is the adjoint twisted Reidemeister torsion of the hyperbolic metric on the knot complement.

$M$  = 3-manifold

$M$  = 3-manifold (closed, oriented ...)

$M$  = 3-manifold (closed, oriented ...)

- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $M$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.

$M$  = 3-manifold (closed, oriented ...)

- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $M$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.



$M$  = 3-manifold (closed, oriented ...)

- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $M$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.
  - ▶ [Reshetikhin-Turaev invariants](#)

$M$  = 3-manifold (closed, oriented ...)

- ▶ Hyperbolic geometry

- ▶ Complete metric of constant curvature  $-1$  on  $M$ .
- ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.

- ▶ Quantum invariants

- ▶ Constructed using representation theory of quantum groups.
- ▶ [Reshetikhin-Turaev invariants](#)  $\text{RT}_r(M, q) \in \mathbb{C}$ ,

$M$  = 3-manifold (closed, oriented ...)

► Hyperbolic geometry

- Complete metric of constant curvature  $-1$  on  $M$ .
- Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.

► Quantum invariants

- Constructed using representation theory of quantum groups.
- [Reshetikhin-Turaev invariants](#)  $\text{RT}_r(M, q) \in \mathbb{C}$ ,  $r \in \mathbb{N}$ .

$M$  = 3-manifold (closed, oriented ...)

- ▶ Hyperbolic geometry
  - ▶ Complete metric of constant curvature  $-1$  on  $M$ .
  - ▶ Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- ▶ Quantum invariants
  - ▶ Constructed using representation theory of quantum groups.
  - ▶ [Reshetikhin-Turaev invariants](#)  $\text{RT}_r(M, q) \in \mathbb{C}$ ,  $r \in \mathbb{N}$ .
  - ▶ Mathematical realization of Witten's invariants

$M$  = 3-manifold (closed, oriented ...)

► Hyperbolic geometry

- Complete metric of constant curvature  $-1$  on  $M$ .
- Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.

► Quantum invariants

- Constructed using representation theory of quantum groups.
- [Reshetikhin-Turaev invariants](#)  $\text{RT}_r(M, q) \in \mathbb{C}$ ,  $r \in \mathbb{N}$ .
- Mathematical realization of Witten's invariants (at  $q = e^{\frac{\pi i}{r}}$ ).

► Conjecture (Chen-Y. '15)

- **Conjecture** (Chen-Y. '15)  $M$  closed oriented hyperbolic,

- **Conjecture** (Chen-Y. '15)  $M$  closed oriented hyperbolic, then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\text{RT}_r(M, e^{\frac{2\pi i}{r}})| = \text{Vol}(M),$$

where  $\text{Vol}(M)$  is the volume of the hyperbolic metric on  $M$ .



► **Conjecture** (Chen-Y. '15)  $M$  closed oriented hyperbolic, then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\text{RT}_r(M, e^{\frac{2\pi i}{r}})| = \text{Vol}(M),$$

where  $\text{Vol}(M)$  is the volume of the hyperbolic metric on  $M$ .  
Namely,

$$|\text{RT}_r(M, e^{\frac{2\pi i}{r}})| \asymp e^{\frac{r}{4\pi} \text{Vol}(M)} \quad \text{as } r \rightarrow \infty, \text{ odd.}$$

- **Conjecture** (Chen-Y. '15)  $M$  closed oriented hyperbolic, then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\text{RT}_r(M, e^{\frac{2\pi i}{r}})| = \text{Vol}(M),$$

where  $\text{Vol}(M)$  is the volume of the hyperbolic metric on  $M$ .  
Namely,

$$|\text{RT}_r(M, e^{\frac{2\pi i}{r}})| \asymp e^{\frac{r}{4\pi} \text{Vol}(M)} \quad \text{as } r \rightarrow \infty, \text{ odd.}$$

- **Conjecture** (Ohtsuki '16, Gang-Romo-Yamazaki '16)

$$\text{RT}_r(M, e^{\frac{2\pi i}{r}}) = \frac{1}{\sqrt{\text{Tor}(M)}} e^{\frac{r}{4\pi} (\text{Vol}(M) + i\text{CS}(M))} \left(1 + O\left(\frac{1}{r}\right)\right)$$

as  $r \rightarrow \infty$ , odd, where  $\text{CS}(M)$  is the Chern-Simons invariant and  $\text{Tor}(M)$  is the adjoint twisted Reidemeister torsion of the hyperbolic metric on  $M$ .

- **Volume Conjecture** (Kashaev '97)  $K$  hyperbolic knot, then

$$\lim_{n \rightarrow \infty} \frac{2\pi}{n} \ln |J_n(K, e^{\frac{2\pi i}{n}})| = \text{Vol}(S^3 \setminus K),$$

where  $\text{Vol}(S^3 \setminus K)$  is the volume of the hyperbolic metric on the knot complement. Namely,

$$|J_n(K, e^{\frac{2\pi i}{n}})| \asymp e^{\frac{n}{2\pi} \text{Vol}(S^3 \setminus K)} \quad \text{as } n \rightarrow \infty.$$

- **Asymptotic Expansion Conjecture** (Witten '89, Gukov '05)

$$J_n(K, e^{\frac{2\pi i}{n}}) = \frac{n^{\frac{3}{2}}}{\sqrt{\text{Tor}(S^3 \setminus K)}} e^{\frac{n}{2\pi} (\text{Vol}(S^3 \setminus K) + i \text{CS}(S^3 \setminus K))} \left(1 + O\left(\frac{1}{n}\right)\right)$$

as  $n \rightarrow \infty$ , where  $\text{CS}(S^3 \setminus K)$  is the Chern-Simons invariant and  $\text{Tor}(S^3 \setminus K)$  is the adjoint twisted Reidemeister torsion of the hyperbolic metric on the knot complement.

## Part II: Asymptotics of relative Reshetikhin-Turaev invariants

## Part II: Asymptotics of relative Reshetikhin-Turaev invariants

Joint work with Ka Ho Wong

$(M, L) = (3\text{-manifold, link in } M)$

$(M, L) = (3\text{-manifold, link in } M)$

- ▶ Hyperbolic geometry
  - ▶ Hyperbolic cone metric

$(M, L) = (3\text{-manifold, link in } M)$

- ▶ Hyperbolic geometry
  - ▶ Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).



$(M, L) = (3\text{-manifold, link in } M)$

- ▶ Hyperbolic geometry
  - ▶ Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
  - ▶ cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .  
cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .  
cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .  
cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the [volume](#), the [Chern-Simons invariant](#) and the [twisted Reidemeister torsion](#) of this metric.
- Quantum invariants
  - [Relative Reshetikhin-Turaev invariants](#)

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $RT_r(M, L, n) \in \mathbb{C}$ ,

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $RT_r(M, L, n) \in \mathbb{C}$ ,  
 $r \in \mathbb{N}$ ,

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $\text{RT}_r(M, L, \mathbf{n}) \in \mathbb{C}$ ,  
 $r \in \mathbb{N}$ ,  $\mathbf{n} = (n_1, \dots, n_{|L|})$ ,



$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $\text{RT}_r(M, L, \mathbf{n}) \in \mathbb{C}$ ,  
 $r \in \mathbb{N}$ ,  $\mathbf{n} = (n_1, \dots, n_{|L|})$ ,  $n_k \in \{0, \dots, r-2\}$ .

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $\text{RT}_r(M, L, \mathbf{n}) \in \mathbb{C}$ ,  
 $r \in \mathbb{N}$ ,  $\mathbf{n} = (n_1, \dots, n_{|L|})$ ,  $n_k \in \{0, \dots, r-2\}$ .
  - $M = S^3 \implies \text{RT}_r(M, L, \mathbf{n}) = J_n(L, q^2)$ .

$(M, L) = (3\text{-manifold, link in } M)$

► Hyperbolic geometry

- Hyperbolic cone metric (metric on  $M$  with constant curvature  $-1$  on  $M \setminus L$  and singular locus  $L$ ).
- cone angles  $= 0 \iff$  complete hyperbolic metric on  $M \setminus L$ .
- cone angles  $= 2\pi \iff$  complete hyperbolic metric on  $M$ .



- Invariants such as the **volume**, the **Chern-Simons invariant** and the **twisted Reidemeister torsion** of this metric.
- Quantum invariants
  - **Relative Reshetikhin-Turaev invariants**  $\text{RT}_r(M, L, n) \in \mathbb{C}$ ,  
 $r \in \mathbb{N}$ ,  $n = (n_1, \dots, n_{|L|})$ ,  $n_k \in \{0, \dots, r-2\}$ .
  - $M = S^3 \implies \text{RT}_r(M, L, n) = J_n(L, q^2)$ .
  - $n = 0 \implies \text{RT}_r(M, L, n) = \text{RT}_r(M)$ .

- ▶  $M$  = obtained from surgery along  $L' \subset S^3$ ,

- $M$  = obtained from surgery along  $L' \subset S^3$ ,  $L \subset S^3 \setminus L' \subset M$ .

- $M$  = obtained from surgery along  $L' \subset S^3$ ,  $L \subset S^3 \setminus L' \subset M$ .

$$(M, L) = \text{Diagram of a knot } L' \text{ with a red circle } L \text{ attached.}$$

- $M$  = obtained from surgery along  $L' \subset S^3$ ,  $L \subset S^3 \setminus L' \subset M$ .

$$(M, L) = \text{diagram of } L \text{ and } L' \text{ in } S^3$$

- **Definition** (Reshetikhin-Turaev '91, Lickorish '93)

$$RT_r(M, L, n) = c \cdot \left\langle \text{diagram of } L \text{ and } L' \text{ with markers } e_n \text{ and } \omega_r \right\rangle,$$

- $M$  = obtained from surgery along  $L' \subset S^3$ ,  $L \subset S^3 \setminus L' \subset M$ .

$$(M, L) = \text{diagram of } L \text{ and } L' \text{ in } S^3$$

- **Definition** (Reshetikhin-Turaev '91, Lickorish '93)

$$RT_r(M, L, n) = c \cdot \left\langle \text{diagram of } L \text{ and } L' \text{ with } e_n \text{ and } \omega_r \right\rangle,$$

where

- $\omega_r$  = Kirby coloring,  $e_n$  =  $n$ -th Jones-Wenzl projector.



- $M$  = obtained from surgery along  $L' \subset S^3$ ,  $L \subset S^3 \setminus L' \subset M$ .

$$(M, L) = \text{diagram of } L' \text{ and } L \text{ in } S^3$$

- **Definition** (Reshetikhin-Turaev '91, Lickorish '93)

$$RT_r(M, L, n) = c \cdot \left\langle \text{diagram of } L' \text{ and } L \text{ with } e_n \text{ and } \omega_r \right\rangle,$$

where

- $\omega_r$  = Kirby coloring,  $e_n$  =  $n$ -th Jones-Wenzl projector.
- $\langle \rangle$  = Kauffman bracket given by

$$(1) \quad \left\langle \text{crossing} \right\rangle = q^{\frac{1}{2}} \left\langle \text{two circles} \right\rangle + q^{-\frac{1}{2}} \left\langle \text{two circles with twist} \right\rangle,$$

$$(2) \quad \left\langle \text{circle} \sqcup D \right\rangle = (-q - q^{-1}) \langle D \rangle.$$

► Conjecture 1 (Wong-Y. '20)

- ▶ **Conjecture 1** (Wong-Y. '20) ( $M, L$ ).

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)}), \quad n_k^{(r)} \in \{0, \dots, r-2\},$$

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)})$ ,  $n_k^{(r)} \in \{0, \dots, r-2\}$ , such that

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)})$ ,  $n_k^{(r)} \in \{0, \dots, r-2\}$ , such that

(1)  $\theta = (\theta_1, \dots, \theta_{|L|})$ ,  $\theta_k = \lim_{r \rightarrow \infty} \left| \frac{4\pi n_k^{(r)}}{r} - 2\pi \right|$  exists,

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)})$ ,  $n_k^{(r)} \in \{0, \dots, r-2\}$ , such that

- (1)  $\theta = (\theta_1, \dots, \theta_{|L|})$ ,  $\theta_k = \lim_{r \rightarrow \infty} \left| \frac{4\pi n_k^{(r)}}{r} - 2\pi \right|$  exists,
- (2)  $M_{L_\theta} = M$  with a hyperbolic cone metric with cone angles  $\theta$  along  $L$  exists.

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)})$ ,  $n_k^{(r)} \in \{0, \dots, r-2\}$ , such that

(1)  $\theta = (\theta_1, \dots, \theta_{|L|})$ ,  $\theta_k = \lim_{r \rightarrow \infty} \left| \frac{4\pi n_k^{(r)}}{r} - 2\pi \right|$  exists,

(2)  $M_{L_\theta} = M$  with a hyperbolic cone metric with cone angles  $\theta$  along  $L$  exists.

Then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\mathbf{RT}_r(M, L, \mathbf{n}^{(r)})| = \text{Vol}(M_{L_\theta}).$$



► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)})$ ,  $n_k^{(r)} \in \{0, \dots, r-2\}$ , such that

- (1)  $\theta = (\theta_1, \dots, \theta_{|L|})$ ,  $\theta_k = \lim_{r \rightarrow \infty} \left| \frac{4\pi n_k^{(r)}}{r} - 2\pi \right|$  exists,
- (2)  $M_{L_\theta} = M$  with a hyperbolic cone metric with cone angles  $\theta$  along  $L$  exists.

Then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\mathbf{RT}_r(M, L, \mathbf{n}^{(r)})| = \text{Vol}(M_{L_\theta}).$$

► **Conjecture 2** (Wong-Y. '21)

$$\mathbf{RT}_r(M, L, \mathbf{n}^{(r)}) = \frac{e^{\frac{1}{2} \sum 1_k}}{\sqrt{\text{Tor}(M_{L_\theta})}} e^{\frac{r}{4\pi} (\text{Vol}(M_{L_\theta}) + i\text{CS}(M_{L_\theta}))} \left(1 + O\left(\frac{1}{r}\right)\right)$$

as  $r \rightarrow \infty$ , odd,

► **Conjecture 1** (Wong-Y. '20)  $(M, L)$ .

$$\mathbf{n}^{(r)} = (n_1^{(r)}, \dots, n_{|L|}^{(r)}), \quad n_k^{(r)} \in \{0, \dots, r-2\}, \text{ such that}$$

(1)  $\theta = (\theta_1, \dots, \theta_{|L|})$ ,  $\theta_k = \lim_{r \rightarrow \infty} \left| \frac{4\pi n_k^{(r)}}{r} - 2\pi \right|$  exists,

(2)  $M_{L_\theta} = M$  with a hyperbolic cone metric with cone angles  $\theta$  along  $L$  exists.

Then

$$\lim_{\substack{r \rightarrow \infty \\ \text{odd}}} \frac{4\pi}{r} \ln |\text{RT}_r(M, L, n^{(r)})| = \text{Vol}(M_{L_\theta}).$$

► Conjecture 2 (Wong-Y. '21)

$$\mathrm{RT}_r(M, L, \mathbf{n}^{(r)}) = \frac{e^{\frac{1}{2} \sum l_k}}{\sqrt{\mathrm{Tor}(M_{L_\theta})}} e^{\frac{r}{4\pi} (\mathrm{Vol}(M_{L_\theta}) + i \mathrm{CS}(M_{L_\theta}))} \left(1 + O\left(\frac{1}{r}\right)\right)$$

as  $r \rightarrow \infty$ , odd, where  $l_k$  is the length of the  $k$ -th component of  $L$  in  $M_{L_\theta}$ .

- ▶ **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that

► **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that

(1)  $M \setminus L \cong$  a fundamental shadow link complement,

► **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that

- (1)  $M \setminus L \cong$  a fundamental shadow link complement,
- (2)  $M$  is obtained from the **fslc** by an integral surgery,

► **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that


- (1)  $M \setminus L \cong$  a fundamental shadow link complement,
- (2)  $M$  is obtained from the **fslc** by an integral surgery,
- (3) cone angles  $\theta$  are sufficiently small.

- ▶ **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that
  - (1)  $M \setminus L \cong$  a **fundamental shadow link complement**,
  - (2)  $M$  is obtained from the **fslc** by an integral surgery,
  - (3) cone angles  $\theta$  are sufficiently small.
  
- ▶ (Costantino-Thurston '08) Every closed oriented 3-manifold is obtained from some fundamental shadow link complement by an integral surgery.

- ▶ **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that
  - (1)  $M \setminus L \cong$  a **fundamental shadow link complement**,
  - (2)  $M$  is obtained from the **fslc** by an integral surgery,
  - (3) cone angles  $\theta$  are sufficiently small.
- ▶ (Costantino-Thurston '08) Every closed oriented 3-manifold is obtained from some fundamental shadow link complement by an integral surgery.
- ▶ If can push  $\theta$  from small to  $2\pi$ , then solve the Volume Conjecture for **all** closed oriented hyperbolic 3-manifolds!




- ▶ **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that
  - (1)  $M \setminus L \cong$  a **fundamental shadow link complement**,
  - (2)  $M$  is obtained from the **fslc** by an integral surgery,
  - (3) cone angles  $\theta$  are sufficiently small.
- ▶ (Costantino-Thurston '08) Every closed oriented 3-manifold is obtained from some fundamental shadow link complement by an integral surgery.
- ▶ If can push  $\theta$  from small to  $2\pi$ , then solve the Volume Conjecture for **all** closed oriented hyperbolic 3-manifolds!
- ▶ **Theorem 2** (Wong-Y. '21) Conjectures 1 & 2 are true for pairs  $(M, L)$  such that

$$M \setminus L \cong S^3 \setminus \text{[link]}$$


and  $M$  is obtained by integral surgery,

- ▶ **Theorem 1** (Wong-Y. '20,'21) Conjecture 1 & 2 are true for pairs  $(M, L)$  such that
  - (1)  $M \setminus L \cong$  a **fundamental shadow link complement**,
  - (2)  $M$  is obtained from the **fslc** by an integral surgery,
  - (3) cone angles  $\theta$  are sufficiently small.
- ▶ (Costantino-Thurston '08) Every closed oriented 3-manifold is obtained from some fundamental shadow link complement by an integral surgery.
- ▶ If can push  $\theta$  from small to  $2\pi$ , then solve the Volume Conjecture for **all** closed oriented hyperbolic 3-manifolds!
- ▶ **Theorem 2** (Wong-Y. '21) Conjectures 1 & 2 are true for pairs  $(M, L)$  such that

$$M \setminus L \cong S^3 \setminus \text{[link]}$$


and  $M$  is obtained by integral surgery, for all  $\theta \in [0, 2\pi]$ .

## Part III: An explicit formula of adjoint twisted Reidemeister torsion

Joint work with Ka Ho Wong


## Fundamental shadow link complement

## Fundamental shadow link complement

- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .


## Fundamental shadow link complement

- Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .



## Fundamental shadow link complement


- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .




- ▶ Glue along triangles to get a handlebody.

## Fundamental shadow link complement

- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .




- ▶ Glue along triangles to get a handlebody.






## Fundamental shadow link complement

- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .




- ▶ Glue along triangles to get a handlebody.




- ▶  $DH = \text{double of handlebody}$ .

## Fundamental shadow link complement

- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .




- ▶ Glue along triangles to get a handlebody.




- ▶  $DH =$  double of handlebody.  $L =$  union of edges.

## Fundamental shadow link complement

- ▶ Take truncated tetrahedra  $\Delta_1, \dots, \Delta_n$ .



- ▶ Glue along triangles to get a handlebody.



- ▶  $DH$  = double of handlebody.  $L$  = union of edges.
- ▶  $DH \setminus L$  = fundamental shadow link complement.

► Gram matrix function: For  $\mathbf{u} = (u_1, \dots, u_6) \in \mathbb{C}^6$ ,


► **Gram matrix function:** For  $\mathbf{u} = (u_1, \dots, u_6) \in \mathbb{C}^6$ ,

$$G(\mathbf{u}) = \begin{bmatrix} 1 & -\cosh u_1 & -\cosh u_2 & -\cosh u_6 \\ -\cosh u_1 & 1 & -\cosh u_3 & -\cosh u_5 \\ -\cosh u_2 & -\cosh u_3 & 1 & -\cosh u_4 \\ -\cosh u_6 & -\cosh u_5 & -\cosh u_4 & 1 \end{bmatrix}.$$

- **Gram matrix function:** For  $u = (u_1, \dots, u_6) \in \mathbb{C}^6$ ,

$$G(u) = \begin{bmatrix} 1 & -\cosh u_1 & -\cosh u_2 & -\cosh u_6 \\ -\cosh u_1 & 1 & -\cosh u_3 & -\cosh u_5 \\ -\cosh u_2 & -\cosh u_3 & 1 & -\cosh u_4 \\ -\cosh u_6 & -\cosh u_5 & -\cosh u_4 & 1 \end{bmatrix}.$$


- If  $u =$  dihedral angles of a hyperbolic tetrahedron  $\times i$ ,



- **Gram matrix function:** For  $u = (u_1, \dots, u_6) \in \mathbb{C}^6$ ,

$$G(u) = \begin{bmatrix} 1 & -\cosh u_1 & -\cosh u_2 & -\cosh u_6 \\ -\cosh u_1 & 1 & -\cosh u_3 & -\cosh u_5 \\ -\cosh u_2 & -\cosh u_3 & 1 & -\cosh u_4 \\ -\cosh u_6 & -\cosh u_5 & -\cosh u_4 & 1 \end{bmatrix}.$$

- If  $u =$  dihedral angles of a hyperbolic tetrahedron  $\times i$ ,




then  $G(u)$  coincides with Gram matrix of the tetrahedron.

- Gram matrix of tetrahedron:

$$G = \begin{bmatrix} 1 & -\cos \theta_1 & -\cos \theta_2 & -\cos \theta_6 \\ -\cos \theta_1 & 1 & -\cos \theta_3 & -\cos \theta_5 \\ -\cos \theta_2 & -\cos \theta_3 & 1 & -\cos \theta_4 \\ -\cos \theta_6 & -\cos \theta_5 & -\cos \theta_4 & 1 \end{bmatrix}.$$

- If  $u =$  dihedral angles of a hyperbolic tetrahedron  $\times i$ ,



then  $G(u)$  coincides with Gram matrix of the tetrahedron.



- ▶ **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery.

- ▶ **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\mathrm{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{u_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component and  $\mathbf{u}_k = (u_{k_1}, \dots, u_{k_6})$  are the holonomy of the meridians intersecting the tetrahedron  $\Delta_k$ ;

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component and  $\mathbf{u}_k = (u_{k_1}, \dots, u_{k_6})$  are the holonomy of the meridians intersecting the tetrahedron  $\Delta_k$ ;  $\mu_i$  is the holonomy of the curve on the  $i$ -th component along which you do the surgery,

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component and  $\mathbf{u}_k = (u_{k_1}, \dots, u_{k_6})$  are the holonomy of the meridians intersecting the tetrahedron  $\Delta_k$ ;  $\mu_i$  is the holonomy of the curve on the  $i$ -th component along which you do the surgery,  $\gamma_i$  is the holonomy of the core curve of the solid torus filled in,

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component and  $\mathbf{u}_k = (u_{k_1}, \dots, u_{k_6})$  are the holonomy of the meridians intersecting the tetrahedron  $\Delta_k$ ;  $\mu_i$  is the holonomy of the curve on the  $i$ -th component along which you do the surgery,  $\gamma_i$  is the holonomy of the core curve of the solid torus filled in, and  $G$  is the Gram matrix function.



- **Conjecture** (Ohtsuki '16, Gang-Romo-Yamazaki '16)

$$\mathrm{RT}_r(M) = \frac{1}{\sqrt{\mathrm{Tor}(M)}} e^{\frac{r}{4\pi}(\mathrm{Vol}(M) + i\mathrm{CS}(M))} \left(1 + O\left(\frac{1}{r}\right)\right)$$

as  $r \rightarrow \infty$ , odd.

- **Conjecture 2** (Wong-Y. '21)

$$\mathrm{RT}_r(M, L, n^{(r)}) = \frac{e^{\frac{1}{2} \sum l_k}}{\sqrt{\mathrm{Tor}(M_{L_\theta})}} e^{\frac{r}{4\pi}(\mathrm{Vol}(M_{L_\theta}) + i\mathrm{CS}(M_{L_\theta}))} \left(1 + O\left(\frac{1}{r}\right)\right)$$

as  $r \rightarrow \infty$ , odd.

- **Theorem 3** (Wong-Y. '21)  $M$  obtained from a fundamental shadow link complement  $(DH, L)$  by a hyperbolic surgery. Then the Reidemeister torsion twisted by the adjoint action of the holonomy representation

$$\text{Tor}(M) = \det \left( \frac{\partial \mu_i}{\partial u_j} \right) \cdot \prod_{i=1}^{|L|} \frac{1}{\sinh^2 \frac{\gamma_i}{2}} \cdot \prod_{k=1}^n \sqrt{\det G \left( \frac{\mathbf{u}_k}{2} \right)},$$

where  $u_i$  is the holonomy of the meridian of the  $i$ -th component and  $\mathbf{u}_k = (u_{k_1}, \dots, u_{k_6})$  are the holonomy of the meridians intersecting the tetrahedron  $\Delta_k$ ;  $\mu_i$  is the holonomy of the curve on the  $i$ -th component along which you do the surgery,  $\gamma_i$  is the holonomy of the core curve of the solid torus filled in, and  $G$  is the Gram matrix function.

Thank You !