

Dentate Gyrus

Yile YING

April 29, 2018

1 Connection




Figure 1: “Despite the fact that mossy cells receive mossy fiber inputs from granule cells, projecting back to the granule cells, spatial representation between the two cell types appears to be independent (shown by dotted lines). Hypothetically, mossy cells may receive strong spatial information from CA3c pyramidal cells and/or semilunar granule cells, both of which also receive entorhinal inputs. EC, entorhinal cortex; GC, granule cell; MC, mossy cell; SGC, semilunar granule cell; IN, interneuron; PC, pyramidal cell; GCL, granule cell layer. GC, SGC, IN and MC are all parts of a dentate gyrus.” [1]

1.1 Input

- From Entorhinal Cortex (EC) layer II: “multiple sensory inputs, including vestibular, olfactory, visual, auditory, and somatosensory from the perirhinal and lateral entorhinal cortex”, and inputs from grid cells in the MEC [2].
- From CA3c to DG mossy cells and interneurons: spatial information [1].
- From CA3 and CA1 as shown in Figure 2: interneuron inhibitory input.




Figure 2: “Boundary-crossing interneurons with cell bodies in CA3 and CA1 constitute a numerically significant and diverse population that relays patterns of activity generated within the CA regions back to granule cells” [3].

1.2 Output

To CA3c pyramidal neuron: sparse encoded spatial representation from granule cells (GC) and multiple spatial representations from mossy cells (MC) [4].

1.3 Main Components

- Granule cells (GC): receive strong excitatory input from EC layer II (e.g. head direction, grid information) via the perforant path; send strong outputs to CA3c pyramidal neurons via the mossy fibers (not mossy cells), then to CA3b and CA1. Granule cells have low firing rate (peak firing rates around 6Hz [5], some could go to 47Hz, and the mean firing rates range from 0.17 to 1.12 Hz, the average mean firing rates are 0.5Hz [6]), sparse spatial coding. GoodSmith et al. (2017) reported only 9% are responsive in a given maze, with most of granule cells only have a single place field [5]. Senzai and Buzsaki (2017) reported 6.6% only has two or more place fields [7]. There are about 1.2×10^6 granule cells in total in a rat dentate gyrus [8].
- Mossy cells: mediate an intrinsic excitation loop of DG neurons, receiving strong input “from a relatively small number of granule cells and providing highly distributed excitatory output to a large number of granule cells”; may also receive spatial input from CA3c neurons [1]. Mossy cells have higher firing rates (about 12Hz?). It is reported 88% are active in a given environment, and mostly have multiple place fields [5]. Senzai and Buzsaki (2017) reported 58% has multiple firing field. The number of mossy cells is about one forth of the granule cells [5].
- Semilunar granule cells: receive projections from EC, and project to mossy cells [9].
- Interneurons: GABA neurons (basket cell) abundant in the dentate gyrus and have high concentration in the mossy cell hilus. [4, 8]. The ratio between basket cells and granule cells is on average around 1:150 [8].

2 Function

- Pattern separation: by sparse coding in granule cells, sparse connections between mossy fibers and CA3, and changes in the position of firings in mossy cells.
- Neuronogenesis: A very recent study showed that there could actually be no neuronogenesis in adults’ DGs [10]. The neuronogenesis is believed to help with new memory formation and pattern separation (old DG cells help with pattern completion) [11].

- Object-place conjunctive encoding: The rats with DG lesions “did not differ from controls in recognition for a change within object feature configuration and object recognition tasks. In contrast, there was a deficit for the dDG lesioned rats relative to controls in recognition for a change within object-spatial feature configuration, complex object-place feature configuration and spatial recognition tasks”. [2, 12].
- Temporal Association: can associate spatial events occurred within 3 min or 30s , may be mediated by neurogenesis [2].

2.1 Scales

- Spatial Scales: the place field diameter for granule cells is about 50 cm, and varies for mossy cells [5].
- Temporal Scales: the location firing of granule cells are heavily theta modulated. If define the time of peak CA1 population activity as phase zero, most of the peak firing happened near phase 270 degree (about 90 degree in advance of CA1 pyramidal cells), and most of granule cells do not fire near phase zero. It is hypothesized that “the phase shift in CA1 could be inherited from the granule cells, and this requires that the synaptic integration time scale must be much shorter than the theta cycle” [6], (generally, the integration time in hippocampus is at least 10 times shorter than the theta cycle) [3]. Besides, “Extended interneuron system provides temporarily distributed inhibition to DG during sharp-wave ripples” [3].

2.2 Evidence of Building Blocks / Modularization

Granule cells remaps more to the change of distal cues, whereas mossy cells remaps more to the change of local cues. [1]

2.3 Evidence of Lateralization

Jordan (2017) reported that object exploration led to the expression of a specific protein in the left DG, suggesting that memories about novel objects may be mainly registered by the left side [13]

2.4 Empirical Lesion studies

- DG lesions impaired the animal to detect changes in the environment [12].
- Rats with damaged DG could not navigate through a familiar maze regardless of more trainings, which shows that DG damage impaired their working memories.

For more lesion studies, see the table in [14].

3 Computational Model

- Rolls et al., (2006) modeled the granule cells using a competitive learning network to “reduce redundancy and produce sparse, orthogonal outputs” [15, 16].
- Faghihi (2015) explored the pattern separation efficiency using simulations of different parameter values in the neural network of the entorhinal cortex and dentate gyrus [17].

References

- [1] Kazu Nakazawa. Dentate Mossy Cell and Pattern Separation. *Neuron*, 93(3):465–467, February 2017.
- [2] Raymond P. Kesner. An analysis of dentate gyrus function (an update). *Behavioural Brain Research*, July 2017.
- [3] Gergely G. Szabo, Xi Du, Mikko Oijala, Csaba Varga, Jack M. Parent, and Ivan Soltesz. Extended Interneuron Network of the Dentate Gyrus. *Cell Reports*, 20(6):1262–1268, August 2017.
- [4] Carolyn R. Houser. Interneurons of the dentate gyrus: an overview of cell types, terminal fields and neurochemical identity. *Progress in Brain Research*, 163:217–232, 2007.

- [5] Douglas GoodSmith, Xiaojing Chen, Cheng Wang, Sang Hoon Kim, Hongjun Song, Andrea Burgalossi, Kimberly M. Christian, and James J. Knierim. Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. *Neuron*, 93(3):677–690.e5, February 2017.
- [6] William E. Skaggs, Bruce L. McNaughton, Matthew A. Wilson, and Carol A. Barnes. Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus*, 6(2):149–172, January 1996.
- [7] Yuta Senzai and Gyrgy Buzski. Physiological Properties and Behavioral Correlates of Hippocampal Granule Cells and Mossy Cells. *Neuron*, 93(3):691–704.e5, February 2017.
- [8] David G. Amaral, Helen E. Scharfman, and Pierre Lavenex. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Progress in brain research*, 163:3–22, 2007.
- [9] Phillip Larimer and Ben W. Strowbridge. Representing information in cell assemblies: persistent activity mediated by semilunar granule cells. *Nature Neuroscience*, 13(2):213, February 2010.
- [10] Shawn F. Sorrells, Mercedes F. Paredes, Arantxa Cebrian-Silla, Kadellyn Sandoval, Dashi Qi, Kevin W. Kelley, David James, Simone Mayer, Julia Chang, Kurtis I. Auguste, Edward F. Chang, Antonio J. Gutierrez, Arnold R. Kriegstein, Gary W. Mathern, Michael C. Oldham, Eric J. Huang, Jose Manuel Garcia-Verdugo, Zhengang Yang, and Arturo Alvarez-Buylla. Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. 555(7696):377–381.
- [11] Toshiaki Nakashiba, Jesse D. Cushman, Kenneth A. Pelkey, Sophie Renaudineau, Derek L. Buhl, Thomas J. McHugh, Vanessa Rodriguez Barrera, Ramesh Chittajallu, Keisuke S. Iwamoto, Chris J. McBain, Michael S. Fanselow, and Susumu Tonegawa. Young Dentate Granule Cells Mediate Pattern Separation whereas Old Granule Cells Contribute to Pattern Completion. *Cell*, 149(1):188–201, March 2012.
- [12] Michael R. Hunsaker, Jenna S. Rosenberg, and Raymond P. Kesner. The role of the dentate gyrus, CA3a,b, and CA3c for detecting spatial and environmental novelty. *Hippocampus*, 18(10):1064–1073, October 2008.
- [13] Jake T. Jordan and Carolyn L. Pytte. Behavior state-dependent lateralization of dorsal dentate gyrus c-Fos expression in mice. *bioRxiv*, page 214239, November 2017.
- [14] Gilberto Fernando Xavier and Valria Catelli Infantozzi Costa. Dentate gyrus and spatial behaviour. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 33(5):762–773, August 2009.
- [15] Edmund T. Rolls, Simon M. Stringer, and Thomas Elliot. Entorhinal cortex grid cells can map to hippocampal place cells by competitive learning. *Network (Bristol, England)*, 17(4):447–465, December 2006.
- [16] Raymond P. Kesner and Edmund T. Rolls. A computational theory of hippocampal function, and tests of the theory: new developments. *Neuroscience and Biobehavioral Reviews*, 48:92–147, January 2015.
- [17] Faramarz Faghihi and Ahmed A. Moustafa. A computational model of pattern separation efficiency in the dentate gyrus with implications in schizophrenia. *Frontiers in Systems Neuroscience*, 9, March 2015.