



TECHNISCHE  
UNIVERSITÄT  
WIEN

B A C H E L O R A R B E I T

**Picard's Theorem**  
**AND**  
**Growth, Zeros and Composition**  
**of Entire Functions**

ausgeführt am

Institut für  
Analysis und Scientific Computing  
TU Wien

unter der Anleitung von

**Ao.Univ.Prof. Dipl.-Ing. Dr.techn. Harald Woracek**

durch

**Ian Hornik**

Matrikelnummer: 12115564

Haizingergasse 18/4

1180 Wien

Wien, am Datum

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Picard's Great Theorem</b>	<b>2</b>
2.1	Bloch's Theorem . . . . .	2
2.2	Schottky's Theorem . . . . .	4
2.3	Picard's Great Theorem . . . . .	6
<b>3</b>	<b>Growth and zeros distribution</b>	<b>10</b>
3.1	Order and Type . . . . .	10
3.2	Hadamard's Theorem . . . . .	11
3.3	Zeros Distribution . . . . .	12
<b>4</b>	<b>Composition of Entire Functions</b>	<b>15</b>
4.1	Pólya's Theorem . . . . .	15
	<b>Bibliography</b>	<b>19</b>

# 1 Introduction

Some words why the subject is of interest.

A reference to the primary literature used.

Overview of used notation.

$$M_f(r) := \max_{|z|=r} |f(z)|$$

## 2 Picard's Great Theorem

### 2.1 Bloch's Theorem

Bloch's Theorem (2.1.4) will show that for any  $f \in H(\mathbb{D})$  with  $f'(0) = 1$  the image of  $\mathbb{D}$  under  $f$  contains a disk of a fixed radius. The point at which said disk is centered at is given in the proof.

If  $G \subseteq \mathbb{C}$  is a domain and  $f \in H(G)$  is non-constant, then  $f(G)$  is a domain as well. Since such an  $f$  is an open mapping, we first observe a simple criterion for the size of disks in their image domain:

**Lemma 2.1.1.** *Let  $G \subset \mathbb{C}$  be a bounded domain,  $f : \overline{G} \rightarrow \mathbb{C}$  continuous and  $f|_G : G \rightarrow \mathbb{C}$  open. If there exists  $a \in G$  such that  $s := \min_{z \in \partial G} |f(z) - f(a)| > 0$ , then  $B_s(f(a)) \subseteq f(G)$ .*

*Proof.* The function  $z \mapsto |z - f(a)|$  is continuous on the compact set  $\partial f(G)$ , hence it assumes its minimum  $m$  at some  $w_* \in \partial f(G)$ . Let  $(z_n)_{n \in \mathbb{N}}$  be a sequence in  $G$ , such that  $\lim_{n \rightarrow \infty} f(z_n) = w_*$ . Since  $\overline{G}$  is compact, we can find a subsequence that converges to some  $z_* \in \overline{G}$  and by continuity of  $f$  we have  $f(z_*) = w_*$ .

Assuming  $z_* \in G$ , since  $f|_G$  is open, the image of any open set in  $G$  containing  $z_*$  under  $f$  must be an open set in  $f(G)$  containing  $w_*$ , which is impossible since  $w_* \in \partial G$ .

Therefore  $z_* \in \partial G$  and we have  $m = |w_* - f(a)| = |f(z_*) - f(a)| \geq s$ , from which it follows that  $B_s(f(a)) \subseteq f(G)$ . ■

**Lemma 2.1.2.** *Let  $a \in \mathbb{C}, r > 0$  and  $B := B_r(a)$ . Suppose further that  $f \in H(\overline{B})$  such that  $\|f'\|_B \leq 2|f'(a)|$ . Then  $B_R(f(a)) \subseteq f(B)$ , where  $R := (3 - 2\sqrt{2})r|f'(a)|$ .*

*Proof.* Without loss of generality we can assume  $a = f(a) = 0$ . Consider the function

$$\alpha_f : \begin{cases} B \rightarrow \mathbb{C}, \\ z \mapsto f(z) - f'(0)z, \end{cases}$$

which satisfies, for the line  $\gamma$  connecting 0 to  $z \in B$ ,

$$|\alpha_f(z)| = \left| \int_{\gamma} f'(\zeta) - f'(0) d\zeta \right| \leq \int_0^1 |f'(tz) - f'(0)| |z| dt. \quad (*)$$

Let  $w \in B$ , then Cauchy's integral formula gives

$$\begin{aligned} |f'(w) - f'(0)| &= \frac{1}{2\pi} \left| \int_{\partial B} \frac{f'(\zeta)}{\zeta - w} - \frac{f'(\zeta)}{\zeta} d\zeta \right| = \frac{1}{2\pi} \left| \int_{\partial B} \frac{wf'(\zeta)}{\zeta(\zeta - w)} d\zeta \right| \leq \\ &\leq \frac{1}{2\pi} \int_{\partial B} \frac{|w|\|f'\|_B}{r(r - |w|)} d\zeta = \frac{|w|}{r - |w|} \|f'\|_B. \end{aligned}$$

Combining the above with (\*) and our estimate on  $\|f'\|_B$  yields

$$|\alpha_f(z)| \leq \int_0^1 \frac{|zt|\|f'\|_B}{r - |zt|} |z| dt \leq \frac{|z|^2}{r - |z|} \|f'\|_B \int_0^1 t dt \leq \frac{|z|^2}{r - |z|} |f'(0)|.$$

Let  $0 < \rho < r$ , then for  $|z| = \rho$  we have

$$\begin{aligned} |f'(0)|\rho - |f(z)| &\leq |\alpha_f(z)| \leq \frac{\rho^2}{r - \rho} |f'(0)| \\ \iff |f(z)| &\geq \left( \rho - \frac{\rho^2}{r - \rho} \right) |f'(0)|. \end{aligned}$$

Considering  $\rho - \rho^2/(r - \rho)$  as a function of  $\rho$ , it assumes its maximum value,  $(3 - 2\sqrt{2})r$ , at  $\rho_* := (1 - \sqrt{2}/2)r \in (0, r)$ . Therefore,

$$|f(z)| \geq (3 - 2\sqrt{2})r |f'(0)| = R, \quad \text{for all } |z| = \rho_*.$$

Invoking Lemma 2.1.1 with  $G := B_{\rho_*}(0)$  thus yields  $B_R(0) \subseteq f(G) \subseteq f(B)$ . ■

**Theorem 2.1.3.** *Let  $f \in H(\overline{\mathbb{D}})$  be non-constant. Then there is a point  $p \in \mathbb{D}$  and a constant  $C_f > 0$  such that  $B_R(f(p)) \subseteq f(\mathbb{D})$ , where  $R := (\frac{3}{2} - \sqrt{2})C_f \geq (\frac{3}{2} - \sqrt{2})|f'(0)|$ .*

*Proof.* The function

$$\alpha_f : \begin{cases} \overline{\mathbb{D}} \rightarrow \mathbb{R} \\ z \mapsto |f'(z)|(1 - |z|) \end{cases}$$

is continuous and assumes its maximum  $C_f > 0$  at some point  $p \in \mathbb{D}$ .

Set  $t := \frac{1}{2}(1 - |p|) > 0$ , then we have  $B_t(p) \subseteq \mathbb{D}$  and  $1 - |z| \geq t$  for  $z \in B_t(p)$ . Since  $|f'(z)|(1 - |z|) \leq C_f = 2t|f'(p)|$ , this implies  $|f'(z)| \leq 2|f'(p)|$ . By Lemma 2.1.2, we obtain  $B_R(f(p)) \subseteq f(\mathbb{D})$ , where

$$R := (3 - 2\sqrt{2})t|f'(p)| = (\frac{3}{2} - \sqrt{2})|f'(p)|(1 - |p|) > \frac{1}{12}|f'(0)|,$$

establishes the assertion. ■

We immediately have:

**Theorem 2.1.4** (Bloch). *Let  $f \in H(\overline{\mathbb{D}})$  and assume that  $f'(0) = 1$ . Then  $f(\mathbb{D})$  contains a disk of radius  $\frac{3}{2} - \sqrt{2}$ .*

In the following we will denote by  $\beta > 0$  any constant satisfying Bloch's Theorem, for example  $\beta = \frac{1}{12} < \frac{3}{2} - \sqrt{2}$ .

**Corollary 2.1.5.** *Let  $G \subset \mathbb{C}$  be a domain and  $f \in H(G)$  with  $f'(c) \neq 0$  for some  $c \in G$ . Then  $f(G)$  contains a disk of every radius  $\beta s |f'(c)|$ , where  $0 < s < d(c, \partial G)$ .*

*Proof.* Without loss of generality we may assume  $c = 0$ . Since  $f$  is analytic on  $\overline{B_s(0)} \subseteq G$ , we have  $g(z) := f(sz)/sf'(0) \in H(\mathbb{D})$ . Since  $g'(0) = 1$ , Bloch's Theorem (2.1.4) yields a disk  $B$  of radius  $\beta$  such that  $B \subseteq g(\mathbb{D})$ . It follows that

$$s|f'(0)|B \subseteq s|f'(0)|g(\mathbb{D}) = f(B_s(0)) \subseteq f(G).$$

■

**Corollary 2.1.6.** *If  $f \in H(\mathbb{C})$  is non-constant, then  $f(\mathbb{C})$  contains a disk of every radius.*

## 2.2 Schottky's Theorem

Holomorphic functions which omit the values 0 and 1 have a universal estimate on the growth of their modulus, given by Schottky's Theorem.

For sets  $G, E \subseteq \mathbb{C}$  we define by  $H(G; E)$  the set of all  $f \in H(G)$  such that  $f(G) \subseteq E$ .

**Lemma 2.2.1.** *It holds that:*

- i. *If  $a, b \in \mathbb{R}$  with  $\cos \pi a = \cos \pi b$ , then  $b = \pm a + 2n$  for some  $n \in \mathbb{Z}$ .*
- ii. *For every  $w \in \mathbb{C}$  there exists a  $v \in \mathbb{C}$  such that  $\cos \pi v = w$  and  $|v| \leq 1 + |w|$ .*

*Proof.* For the first part, it suffices to notice that

$$0 = \cos \pi a - \cos \pi b = -2 \sin \frac{\pi}{2}(a+b) \sin \frac{\pi}{2}(a-b).$$

Since the complex cosine function is surjective and  $\mathbb{R}$ -periodic, we can choose  $v = a + ib$  with  $w = \cos \pi v$  and  $|a| \leq 1$ . Now we have

$$\begin{aligned} |w|^2 &= |\cos(\pi a + i\pi b)|^2 = |\cos \pi a \cos i\pi b + \sin \pi a \sin i\pi b|^2 = \\ &= |\cos \pi a \cosh \pi b - i \sin \pi a \sinh \pi b|^2 = \\ &= \cos^2 \pi a \cosh^2 \pi b + \sin^2 \pi a \sinh^2 \pi b = \\ &= \cos^2 \pi a + \cos^2 \pi a \sinh^2 \pi b + \sin^2 \pi a \sinh^2 \pi b = \\ &= \cos^2 \pi a + \sinh^2 \pi b \geq \sinh^2 \pi b \geq \pi^2 b^2, \end{aligned}$$

where the last inequality holds since  $\sinh x \geq x$  for  $x \geq 0$ . We conclude

$$|v| = \sqrt{a^2 + b^2} \leq \sqrt{1 + |w|^2/\pi^2} \leq 1 + |w|$$

■

**Lemma 2.2.2.** *Let  $G \subseteq \mathbb{C}$  be a simply connected domain and  $f \in H(G; \mathbb{C} \setminus \{-1, 1\})$ . Then there exists an  $F \in H(G)$  such that  $f = \cos F$ .*

*Proof.* Since  $1 - f^2$  vanishes nowhere in  $G$  it has a square root  $g \in H(G)$ . It follows that

$$1 = f^2 + g^2 = (f + ig)(f - ig).$$

Thus  $f + ig$  vanishes nowhere and there exists an  $F \in H(G)$  with  $f + ig = e^{iF}$ . Additionally we have  $f - ig = e^{-iF}$  and therefore

$$f = \frac{1}{2}(e^{iF} + e^{-iF}) = \cos F.$$

■

**Lemma 2.2.3.** *Let  $f \in H(\overline{\mathbb{D}}; \mathbb{C} \setminus \{0, 1\})$ . Then there exists a  $g \in H(\overline{\mathbb{D}})$  such that:*

- i.  $f = \frac{1}{2}(1 + \cos \pi(\cos \pi g))$ , and  $|g(0)| \leq 3 + 2|f(0)|$ .
- ii.  $g(\overline{\mathbb{D}})$  contains no disk of radius 1.
- iii.  $|g(z)| \leq |g(0)| + \theta/b(1 - \theta)$ , for all  $|z| \leq \theta$  where  $0 < \theta < 1$ .

*Proof.* By Lemma 2.2.2 there exists a  $\tilde{F} \in H(\overline{\mathbb{D}})$  such that  $2f - 1 = \cos \pi \tilde{F}$ . By Lemma 2.2.1 there is a  $b \in \mathbb{C}$  such that  $\cos \pi b = 2f(0) - 1$  and  $|b| \leq 1 + |2f(0) - 1| \leq 2 + 2|f(0)|$ . Additionally we have that  $b = \pm \tilde{F}(0) + 2k$  for some  $k \in \mathbb{Z}$ . Then  $F := \pm \tilde{F} + 2k \in H(\overline{\mathbb{D}})$  satisfies  $F(0) = b$ . Since  $F$  must omit all integers, there exists a  $\tilde{g} \in H(\overline{\mathbb{D}})$  such that  $F = \cos \pi \tilde{g}$ . Again, there is an  $a \in \mathbb{C}$  such that  $\cos \pi a = b$  and  $|a| \leq 1 + |b| \leq 3 + 2|f(0)|$ . Like above, we have  $a = \pm \tilde{g}(0) + 2\ell$  for some  $\ell \in \mathbb{Z}$ , thus  $g := \pm \tilde{g} + 2\ell \in H(\overline{\mathbb{D}})$  satisfies  $g(0) = a$ . Ultimately, we obtain

$$f = \frac{1}{2}(1 + \cos \pi(\cos \pi g)), \quad \text{and} \quad |g(0)| = |a| \leq 3 + 2|f(0)|.$$

To show (ii) we consider the set

$$A := \{m \pm i\pi^{-1} \log(n + \sqrt{n^2 - 1}) : m \in \mathbb{Z}, n \in \mathbb{N} \setminus \{0\}\},$$

the points of which can be considered the vertices of a rectangular grid in  $\mathbb{C}$ . The width of said rectangles is 1, and since

$$\begin{aligned} \log((n+1) + \sqrt{(n+1)^2 - 1}) - \log(n + \sqrt{n^2 - 1}) &= \\ &= \log \frac{1 + \frac{1}{n} + \sqrt{1 + \frac{2}{n}}}{1 + \sqrt{1 - \frac{1}{n^2}}} \leq \log(2 + \sqrt{3}) < \pi \end{aligned}$$

their height is strictly bounded above by 1. Let  $a = p + i\pi^{-1} \log(q + \sqrt{q^2 - 1}) \in A$ , then

$$\begin{aligned} \cos \pi a &= \frac{1}{2}(e^{i\pi a} + e^{-i\pi a}) = \frac{1}{2}(-1)^p((q + \sqrt{q^2 - 1})^{-1} + (q + \sqrt{q^2 - 1})) = \\ &= (-1)^p \frac{1}{2} \frac{1 + q^2 + 2q\sqrt{q^2 - 1} + q^2 - 1}{q + \sqrt{q^2 - 1}} = (-1)^p q. \end{aligned}$$

Thus  $\cos \pi(\cos \pi a) = \pm 1$ . But  $0, 1 \notin f(\overline{\mathbb{D}})$ , therefore  $a \notin g(\overline{\mathbb{D}})$ , thus  $g(\overline{\mathbb{D}}) \cap A = \emptyset$  and  $g(\overline{\mathbb{D}})$  cannot contain a disk of radius 1.

For (iii), let  $0 < \theta < 1$ , then for  $|z| \leq \theta$  we have  $1 - \theta \leq d(z, \partial \mathbb{D})$ . By (ii) and Corollary 2.1.5 it follows that  $\beta(1 - \theta)|g'(z)| \leq 1$ . Let  $\gamma$  denote the line connecting 0 and  $z$ , then

$$|g(z)| \leq |g(0)| + |g(z) - g(0)| \leq |g(0)| + \int_{\gamma} |g'(\zeta)| d\zeta \leq |g(0)| + \frac{\theta}{\beta(1 - \theta)}$$

yields our desired estimate. ■

**Remark 2.2.4.** One may notice that in the proof of Lemma 2.2.3 (i) and (ii) we never used the fact that we act on the unit disk. Indeed, applying the proof for  $\mathbb{C}$ , we can quickly obtain Picard's Little Theorem, which states that any  $f \in H(\mathbb{C}; \mathbb{C} \setminus \{a, b\})$ , for distinct points  $a, b \in \mathbb{C}$ , is constant:

Let  $g(z) := \frac{f(z) - a}{b - a} \in H(\mathbb{C}; \mathbb{C} \setminus \{0, 1\})$  then, by the argument above,  $g$  does not contain a disk of radius 1. Corollary 2.1.6 thus implies that  $g$  is constant, and therefore so is  $f$ .

**Theorem 2.2.5** (Schottky). *There exists a function  $\psi(\theta, \omega) : (0, 1) \times (0, \infty) \rightarrow (0, \infty)$  such that for any  $f \in H(\overline{\mathbb{D}}; \mathbb{C} \setminus \{0, 1\})$  with  $|f(0)| \leq \omega$  it holds that*

$$|f(z)| \leq \psi(\theta, \omega), \quad |z| \leq \theta. \quad (2.1)$$

*Proof.* Note that for all  $w \in \mathbb{C}$  we have  $|\cos w| \leq e^{|w|}$  and  $\frac{1}{2}|1 + \cos w| \leq e^{|w|}$ . Hence, from Lemma 2.2.3, we get

$$\begin{aligned} |f(z)| &= \left| \frac{1}{2}(1 + \cos \pi(\cos \pi g(z))) \right| \leq \exp(\pi \exp \pi |g(z)|) \leq \\ &\leq \exp(\pi \exp \pi (|g(0)| + \theta/\beta(1 - \theta))) \leq \\ &\leq \exp(\pi \exp \pi (3 + 2\omega + \theta/\beta(1 - \theta))), \end{aligned}$$

and defining  $\psi(\theta, \omega)$  as the rightmost term establishes the assertion. ■

## 2.3 Picard's Great Theorem

**Definition 2.3.1.** Let  $G \subseteq \mathbb{C}$  be a domain and  $\mathcal{F} \subseteq H(G)$ . Then:

- $\mathcal{F}$  is called *locally bounded* if for every  $w \in G$  there is a neighborhood  $U$  of  $w$  and a constant  $C > 0$  such that  $|f(z)| \leq C$  for all  $f \in \mathcal{F}$  and  $z \in U$ .



- $\mathcal{F}$  is called *normal* in  $G$  if every sequence in  $\mathcal{F}$  has a subsequence which converges compactly in  $G$  to some  $f \in H(G)$  or to  $\infty$ .

The following well-known theorem establishes the equivalence of the two concepts:

**Theorem 2.3.2** (Montel). *Let  $G \subseteq \mathbb{C}$  be a domain. Then a family  $\mathcal{F} \subseteq H(G)$  is normal if and only if it is locally bounded.*

**Theorem 2.3.3** (Hurwitz). *Let  $G \subseteq \mathbb{C}$  be a domain and  $(f_n)_{n \in \mathbb{N}}$  a sequence in  $H(G)$  that converges compactly to  $f$ . If for every  $n \in \mathbb{N}$  the number of  $a$ -points of  $f_n$  is bounded by some  $m \in \mathbb{N}_0$ , then either the number of  $a$ -points of  $f$  are also bounded by  $m$ , or  $f \equiv a$ .*

The following lemma can be interpreted as a sharpened version of Montel's Theorem (2.3.2) and is sometimes referred to as the *fundamental normality test*.

**Lemma 2.3.4.** *For any domain  $G \subseteq \mathbb{C}$  the family  $\mathcal{F} := H(G, \mathbb{C} \setminus \{0, 1\})$  is normal in  $G$ .*

*Proof.* We shall give the proof in three steps:

1. Let  $w \in G$ ,  $c > 0$  and  $\mathcal{F}_*$  be a subfamily of  $\mathcal{F}$  such that  $|f(w)| \leq c$  for all  $f \in \mathcal{F}_*$ . We aim to show that there is a neighborhood of  $w$  in which  $\mathcal{F}_*$  is bounded. Select  $t > 0$  such that  $\overline{B_t(w)} \subset G$ . Without loss of generality we may assume  $w = 0$  and  $t = 1$ . By the maximum modulus principle and Schottky's Theorem we obtain

$$\sup\{\|f\|_{B_{1/2}(0)} : f \in \mathcal{F}_*\} \leq \psi(1/2, c) < \infty.$$

2. Fix some  $w_* \in G$  and set  $\mathcal{F}_1 := \{f \in \mathcal{F} : |f(w_*)| \leq 1\}$ . We aim to show that  $\mathcal{F}_1$  is locally bounded (and by Montel's Theorem therefore normal) in  $G$ . Consider the set

$$U := \{w \in G : \mathcal{F}_1 \text{ is bounded in a neighborhood of } w\},$$

which is open in  $G$ . By (1) we have that  $w_* \in U$ . For sake of contradiction, assume that  $U \neq G$ . Then there exists some  $w \in \partial U \cap G$  and a sequence  $f_n \in \mathcal{F}_1$  with  $\lim_{n \rightarrow \infty} f_n(w) = \infty$ . Set  $g_n := 1/f_n \in \mathcal{F}$ , then  $\lim_{n \rightarrow \infty} g_n(w) = 0$  and by (1) the family  $(g_n)_{n \in \mathbb{N}}$  must be bounded in a neighborhood of  $w$ . By Montel's Theorem it is therefore normal and there exists a subsequence  $(g_{n_k})_{k \in \mathbb{N}}$  which converges compactly to a  $g \in H(B)$ , for some disk  $B$  around  $w$ . The functions  $g_n$  have no zeros, but  $g(w) = 0$ ; by Hurwitz's Theorem we therefore have  $g \equiv 0$ . Then for any  $z \in B \cap U$  we have

$$\lim_{k \rightarrow \infty} f_{n_k}(z) = \lim_{k \rightarrow \infty} 1/g_{n_k}(z) = \infty,$$

contradicting the assumption that  $\mathcal{F}_1$  is bounded in a neighborhood of  $z$ .

3. We can now conclude the proof. Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{F}$ . If it has a subsequence in  $\mathcal{F}_1$ , then the claim follows from (2).

On the other hand, if there are only finitely many  $f_n$  in  $\mathcal{F}_1$ , then  $(1/f_n)_{n \in \mathbb{N}}$  has some subsequence, say  $g_n$ , in  $\mathcal{F}_1$ , which converges compactly in  $G$  to some  $g \in H(G)$ . If  $g$  has no zeros, then the subsequence  $1/g_n$  of the sequence  $f_n$  converges compactly in  $G$  to  $1/g$ . If  $g$  has zeros, then by Hurwitz's Theorem (2.3.3) we have  $g \equiv 0$ , thus  $1/g_n$  converges compactly in  $G$  to  $\infty$ . ■

**Lemma 2.3.5.** *Let  $f \in H(\mathbb{D}^\times; \mathbb{C} \setminus \{0, 1\})$ . Then  $f$  or  $1/f$  is bounded in a punctured neighborhood of zero.*

*Proof.* Set  $f_n(z) := f(z/n) \in H(\mathbb{D}^\times; \mathbb{C} \setminus \{0, 1\})$ . By Lemma 2.3.4 there exists a subsequence  $(f_{n_k})_{k \in \mathbb{N}}$  such that  $(f_{n_k})_{k \in \mathbb{N}}$  or  $(1/f_{n_k})_{k \in \mathbb{N}}$  is bounded on  $\partial B_{1/2}(0)$ .

In the first case, we have  $|f(z/n_k)| \leq M$  for some  $M > 0$  and all  $|z| = \frac{1}{2}$ ,  $k \in \mathbb{N}$ . Thus  $f$  is bounded on every circle of radius  $1/(2n_k)$ . By the maximum modulus principle,  $f$  must therefore be bounded on every annulus  $1/(2n_{k+1}) \leq |z| \leq 1/(2n_k)$ . This yields that  $f$  is bounded in a punctured neighborhood of zero.

In the second case, the same argument shows that  $1/f$  is bounded in a punctured neighborhood of zero. ■

**Theorem 2.3.6** (Picard's Great Theorem). *Let  $G \subseteq \mathbb{C}$  be open,  $w \in G$  and suppose  $f \in H(G \setminus \{w\})$  such that  $f$  has an essential singularity at  $w$ . Then  $f$  assumes all values in  $\mathbb{C}$ , with at most one exception, infinitely often in any punctured neighborhood of  $w$ .*

*Proof.* Aiming for contradiction, assume that  $f$  only takes on  $z_0, z_1 \in \mathbb{C}$  finitely often in some punctured neighborhood  $W$  of  $w$ . Then  $W$  contains a punctured disk of radius  $\varepsilon > 0$  around  $w$ , on which  $f$  does not assume  $z_0, z_1$ . The function

$$g(z) := \frac{f(\varepsilon z + w) - z_0}{z_1 - z_0} \in H(\mathbb{D}^\times; \mathbb{C} \setminus \{0, 1\})$$

has an essential singularity at zero. By Lemma 2.3.5, we have that either  $g$  or  $1/g$  must be bounded in a neighborhood of zero. In the former case the singularity must therefore be removable, whereas in the latter case it must be a pole, yielding a contradiction. ■

**Corollary 2.3.7.** *Every entire transcendental function assumes every value in  $\mathbb{C}$  infinitely often, with at most one exception.*

*Proof.* Let  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  be transcendental and entire, and consider  $g(z) := f(1/z)$ . Then the Laurent series expansion of  $g$  has an infinite principal part, therefore  $g$  has an essential singularity at zero. By Picard's Great Theorem  $g$  assumes all values in  $\mathbb{C}$  on  $\mathbb{C}^\times$ , except at most one. Thus  $f$  does the same on  $\mathbb{C}^\times$ . ■

**Corollary 2.3.8** (Picard's Little Theorem). *Every non-constant entire function omits at most one value.*

*Proof.* A non-constant entire function  $f$  is either a non-constant polynomial or transcendental. In the latter case, the claim follows from Corollary 2.3.7.

Otherwise let  $w \in \mathbb{C}$ , then  $f(z) - w$  has a complex root by the Fundamental Theorem of Algebra and hence  $f$  assumes all values. ■

## 3 Growth and zeros distribution

### 3.1 Order and Type

**Definition 3.1.1.** Let  $f$  be an entire function. The *order* of  $f$  is defined by

$$\rho_f := \limsup_{r \rightarrow \infty} \frac{\log \log M_f(r)}{\log r}. \quad (3.1)$$

Constant functions, by convention, have order 0.

Note that, for any entire function  $f$ , we have  $0 \leq \rho_f \leq \infty$ . Equivalently,  $\rho_f$  can also be defined as the infimum over all  $\rho > 0$  such that  $M_f(r) = O(\exp(r^\rho))$  as  $r \rightarrow \infty$ .

An elementary fact regarding the order of the sum or product of two entire functions of finite order can immediately be obtained by naively estimating their respective maximum modulus:

**Proposition 3.1.2.** *Let  $f, g$  be entire functions of finite order. Then it holds that:*

- i.  $\rho_{f+g} \leq \max\{\rho_f, \rho_g\}$
- ii.  $\rho_{fg} \leq \max\{\rho_f, \rho_g\}$

**Proposition 3.1.3.** *Let  $f$  be an entire function of finite order, then  $\rho_{f'} = \rho_f$ .*

*Proof.* **TODO.** ■

For functions of finite and positive order, we can obtain a natural refinement of the concept of order:

**Definition 3.1.4.** Let  $f$  be an entire function of finite and positive order. The *type* of  $f$  is defined by

$$\tau_f := \limsup_{r \rightarrow \infty} \frac{\log M_f(r)}{r^{\rho_f}} \quad (3.2)$$

For any entire function  $f$  with  $0 < \rho_f < \infty$ , we have  $0 \leq \tau_f \leq \infty$ . Equivalently,  $\tau_f$  can also be defined as the infimum over all  $\tau > 0$  such that  $M_f(r) = O(\exp(\tau r^{\rho_f}))$  as  $r \rightarrow \infty$ .

Imitating the proof of Proposition 3.1.3, we immediately get:

**Proposition 3.1.5.** *Let  $f$  be an entire function of finite type, then  $\tau_{f'} = \tau_f$ .*

**Definition 3.1.6.** Let  $f$  be an entire function. Then  $f$  is said to be of *growth*  $(a, b)$  if

- $\rho_f < a$ , or
- $\rho_f = a$  and  $\tau_f \leq b$ .

**Example 3.1.7.** Maybe give some functions together with their order and type?

## 3.2 Hadamard's Theorem

**Theorem 3.2.1** (Weierstrass [1, 4]). Let  $(z_j)_{j \in \mathbb{N}}$  be a sequence in  $\mathbb{C}$  without accumulation points. Then there exists an entire function  $E$  (called the Weierstrass canonical product formed from said sequence) that has zeros precisely at  $(z_j)_{j \in \mathbb{N}}$ , with multiplicities equal to how often  $z_j$  occurs in the sequence.

In particular, we have

$$E(z) = z^k \prod_{n=1}^{\infty} \left(1 - \frac{z}{z_n}\right) e^{R_n(z/z_n)}, \quad (3.3)$$

where  $k$  is the order of the zero at  $z = 0$  and  $R_n$  is a polynomial, namely a truncation of the power series for  $-\log(1 - \frac{z}{z_n})$  chosen of smallest degree to ensure convergence of the product [3].

Weierstrass' Theorem is also known as the *Weierstrass Factorization Theorem*, due to the following corollary:

**Corollary 3.2.2.** Let  $f$  be an entire function with zeros  $(z_j)_{j \in \mathbb{N}}$ . Then there exists an entire function  $g$ , such that

$$f(z) = e^{g(z)} E(z),$$

where  $E$  is a Weierstrass canonical product formed from the zeros of  $f$ .

*Proof.* Since  $f/E$  has removable singularities at all  $(z_j)_{j \in \mathbb{N}}$ , we have that  $f/E$  is entire and nowhere zero. Thus there exists an entire function  $g$  with  $f/E = e^g$ , which yields  $f = e^g E$ . ■

Hadamard's Theorem will show that, for functions of finite order  $\rho$ , the function  $g$  in Corollary 3.2.2 can be taken to be a polynomial of degree less than  $\rho$  and the degree of the polynomials  $R_n$  in (3.3) can be taken to be independent of  $n$ . To prove this we require the following lemma, which can be interpreted as a version of the maximum modulus principle applied to the real part of an analytic function.

**Lemma 3.2.3** (Borel-Carathéodory). Let  $f$  be analytic in  $\overline{\mathbb{D}}$  and let

$$A_f(r) = \max_{|z|=r} \Re f(z).$$

Then, for  $0 < r < R$ ,

$$M_f(r) \leq \frac{2r}{R-r} A_f(R) + \frac{R+r}{R-r} |f(0)|$$

and, if additionally  $A_f(R) \geq 0$ , then for  $n \in \mathbb{N}$

$$\max_{|z|=r} |f^{(n)}(z)| \leq \frac{2^{n+2} n! R}{(R-r)^{n+1}} (A_f(R) + |f(0)|).$$

*Proof.* **TODO.** ■

**Theorem 3.2.4** (Hadamard). *Let  $f$  be an entire function of finite order with zeros  $(z_j)_{j \in \mathbb{N}}$ . Then there exists a polynomial  $Q$  with  $\deg Q \leq \rho_f$ , such that*

$$f(z) = e^{Q(z)} E(z),$$

where  $E$  is a Weierstrass canonical product formed from the zeros of  $f$ .

*Proof.* **TODO.** ■

### 3.3 Zeros Distribution

We recall a rather explicit connection between the moduli of the zeros of an analytic function and the modulus of the function itself:

**Theorem 3.3.1** (Jensen [4]). *Let  $f$  be analytic on  $B_R(0)$  with  $f(0) \neq 0$  and let  $r_1, r_2, \dots$  denote the moduli of the zeros of  $f$  in  $B_R(0)$  arranged in a non-decreasing sequence. Then, for  $r_n < r < r_{n+1}$ , we have*

$$\frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta = \log |f(0)| + \log \frac{r^n}{r_1 \dots r_n}.$$

**Definition 3.3.2.** Let  $f$  be analytic on  $B_R(0)$ . Then, for  $0 < r < R$ , we denote by  $n_f(r)$  the number of zeros of  $f$  in  $\overline{B_r(0)}$ .

**Corollary 3.3.3.** *Let  $f$  be analytic on  $B_R(0)$  with  $f(0) \neq 0$ . Then, for  $0 < r < R$ , we have*

$$\frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta = \log |f(0)| + \int_0^r \frac{n_f(s)}{s} ds$$

*Proof.* Let  $r_1, r_2, \dots$  denote the moduli of the zeros of  $f$  in  $B_R(0)$  arranged in a non-

decreasing sequence. Then, for any  $r_n < r < r_{n+1}$ , we obtain

$$\begin{aligned} \log \frac{r^n}{r_1 \dots r_n} &= \sum_{k=1}^n \log \frac{r}{r_k} = \sum_{k=1}^n \int_{r_k}^r \frac{1}{s} ds = \\ &= \sum_{k=1}^n \int_0^r \mathbb{1}_{(r_k, \infty)}(s) \frac{1}{s} ds = \int_0^r \left( \sum_{k=1}^n \mathbb{1}_{(r_k, \infty)}(s) \right) \frac{1}{s} ds = \\ &= \int_0^r \frac{n_f(s)}{s} ds \end{aligned}$$

and Theorem 3.3.1 concludes the claim. ■

In particular, we observe that the more zeros a function  $f(z)$  has, the faster its modulus must grow as  $|z| \rightarrow \infty$ . The converse is naturally false, as seen by iterated exponentials [3].

**Definition 3.3.4.** Let  $f$  be an entire function and let  $(r_j)_{j \in \mathbb{N}}$  denote the non-zero moduli of its zeros (if any) arranged in non-decreasing order. Then

$$\lambda_f := \inf \left\{ \lambda > 0 : \sum_{n=1}^{\infty} \frac{1}{r_n^\lambda} < \infty \right\}$$

is called the *exponent of convergence of the zeros of  $f$* . If  $f$  has finitely many zeros, then we set  $\lambda_f = 0$  by convention.

Furthermore, the *exponent of convergence of the  $a$ -points of  $f$*  is defined as exponent of convergence of zeros of  $f(z) - a$ .

**Theorem 3.3.5.** Let  $f$  be an entire function of finite order. Then  $\lambda_f \leq \rho_f$ .

*Proof.* TODO. ■

**Example 3.3.6.** Consider the entire function  $f(z) := \sin(z)$ , we want to calculate  $\lambda_f$  and  $\rho_f$ . First, let  $\lambda > 0$  and consider that  $f$  has zeros at  $(n\pi)_{n \in \mathbb{Z}}$ . Since

$$\sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{|n\pi|^\lambda} = \frac{2}{\pi^\lambda} \sum_{n=1}^{\infty} \frac{1}{|n|^\lambda}$$

is finite if and only if  $\lambda > 1$ , we obtain  $\lambda_f = 1$ . Furthermore, we have  $|\sin(z)| \leq e^{|\Im z|}$  and therefore

$$M_f(r) \leq \max_{|z|=r} |e^z|.$$

Since  $e^z$  is of order 1, this implies  $\rho_f \leq 1$ . Finally, Theorem 3.3.5 concludes  $\rho_f = 1$ .

**Theorem 3.3.7.** Let  $E$  be a Weierstrass canonical product of finite order. Then  $\lambda_E = \rho_E$ .

*Proof.* **TODO.** ■

**Theorem 3.3.8.** *Let  $f$  be an entire function of finite, non-integer order. Then  $\rho_f = \lambda_f$ .*

*Proof.* By Theorem 3.3.5 we have  $\lambda_f \leq \rho_f$ . Invoking Hadamard's Theorem we can write  $f = e^Q E$  for a polynomial  $Q$  with  $\deg Q \leq \rho_f$ . Since  $\rho_f$  is not an integer, this implies  $\deg Q \leq \lfloor \rho_f \rfloor < \rho_f$ . Now, again by Hadamard's Theorem,  $e^Q$  has order  $\deg Q$  and by Theorem 3.3.7  $E$  has order  $\lambda_f$ . Using Proposition 3.1.2 we obtain

$$\rho_f \leq \max\{\deg Q, \lambda_f\} = \lambda_f \leq \rho_f,$$

implying  $\rho_f = \lambda_f$ . ■

**Theorem 3.3.9.** *Let  $f$  be an entire function of finite, non-integer order. Then  $f$  has infinitely many zeros.*

*Proof.* By Theorem 3.3.8 we have  $\rho_f = \lambda_f$ . Since  $\rho_f$  is not an integer,  $\lambda_f > 0$ , which implies that  $f$  has infinitely many zeros. ■

Maybe introduce Borel exceptional values as a definition? But then again, I will never need them again. Maybe also add a remark on the relation to lacunary values (Picard).

**Theorem 3.3.10** (Borel). *Existence of Borel exceptional values.*

*Proof.* **TODO.** ■



## 4 Composition of Entire Functions

As seen by Proposition 3.1.2, the sum or product of two entire functions of finite order is again of finite order. This is not the case when composition is involved; indeed, consider the function  $f(z) := \exp \exp z \in H(\mathbb{C})$ , which is of infinite order, yet is the composition of two functions of order 1.

### 4.1 Pólya's Theorem

Necessary conditions for the order of a composition to be finite will be given by Pólya's Theorem, the proof of which relies on a result that was first proven by Harald Bohr. While Bloch's Theorem dealt with disks contained in the image of holomorphic functions, we now focus on circles contained in such images.

**Remark 4.1.1.** To justify our next definition, let  $G \subset \mathbb{C}$  be a bounded domain with  $0 \in G$ . We claim that  $\overline{G}$  contains a circle (the boundary of a disk) centered at 0 of maximal, non-zero radius; in other words, we claim that the set

$$S := \{r > 0 : \partial B_r(0) \subseteq \overline{G}\}$$

has a maximum. Firstly, since  $G$  is open and contains 0, there is some  $t > 0$  such that  $B_t(0) \subseteq G$ . Since  $\partial B_t(0) \subset \overline{B_t(0)} \subseteq \overline{G}$  we thus have  $t \in S$  and  $S$  is non-empty. It now suffices to show that the set  $S_0 := S \cup \{0\}$  is compact. Clearly  $S_0$  is bounded from below by 0 and since  $\overline{G}$  is bounded it must also be bounded from above. Thus it remains to show that  $S_0$  is closed.

Let  $(r_n)_{n \in \mathbb{N}}$  be a convergent sequence in  $S_0$  with limit  $r \in [0, \infty)$ . If  $r = 0$ , then  $r \in S_0$ . Otherwise we claim that  $\partial B_r(0) \subseteq \overline{G}$ . Let  $z \in \partial B_r(0)$ , and set  $z_n := r_n z / r$  for  $n \in \mathbb{N}$ . Now  $|z_n| = r_n$ , thus  $z_n \in \partial B_{r_n}(0) \subseteq \overline{G}$  and therefore  $(z_n)_{n \in \mathbb{N}}$  is a sequence in  $\overline{G}$ . Clearly  $z_n \rightarrow z$  and since  $\overline{G}$  is closed we therefore have  $z \in \overline{G}$ . Since  $z$  was arbitrary, this concludes the claim and we have  $r \in S_0$ .

Thus  $S_0$  is compact and therefore has a maximum  $m \geq t > 0$ , which is also a maximum of  $S$ .

If  $G \subset \mathbb{C}$  is a bounded domain and  $f \in H(G) \cap C(\overline{G}; \mathbb{C})$  is non-constant with  $f(0) = 0$ , then  $f(G)$  is a bounded domain with  $0 \in f(G)$ . This, together with the above remark, justifies the following definition:

**Definition 4.1.2.** Let  $f \in H(\mathbb{D}) \cap C(\overline{\mathbb{D}}; \mathbb{C})$  be non-constant with  $f(0) = 0$ , then we define

$$r_f := \max\{r > 0 : \partial B_r(0) \subseteq f(\overline{\mathbb{D}})\} > 0. \quad (4.1)$$

**Theorem 4.1.3** (Bohr). *Fix  $0 < \theta < 1$  and let  $f \in H(\mathbb{D}) \cap C(\overline{\mathbb{D}}; \mathbb{C})$  such that  $f(0) = 0$  and  $M_f(\theta) = 1$ . Then  $r_f \geq C_\theta$ , where  $C_\theta > 0$  is a constant depending only on  $\theta$ .*

*Proof.* Suppose  $f$  satisfies the hypothesis of the theorem, let  $\varepsilon > 0$  and set  $R_\varepsilon := r_f + \varepsilon$ . By definition of  $r_f$ , for all  $r \geq R_\varepsilon$  there exists some point  $w_r \in \partial B_r(0)$  with  $w_r \notin f(\overline{\mathbb{D}})$ . Choose such points  $w_{R_\varepsilon}, w_{2R_\varepsilon}$  and define

$$h(z) := \frac{f(z) - w_{R_\varepsilon}}{w_{2R_\varepsilon} - w_{R_\varepsilon}} \in H(\mathbb{D}, \mathbb{C} \setminus \{0, 1\}).$$

Since

$$|h(0)| = \left| \frac{f(0) - w_{R_\varepsilon}}{w_{2R_\varepsilon} - w_{R_\varepsilon}} \right| \leq \frac{|w_{R_\varepsilon}|}{||w_{2R_\varepsilon}| - |w_{R_\varepsilon}||} = \frac{R_\varepsilon}{2R_\varepsilon - R_\varepsilon} = 1,$$

by Schottky's Theorem we have  $|h(z)| \leq \psi(\theta, 1)$  for all  $|z| \leq \theta$ . Therefore

$$|f(z)| - R_\varepsilon \leq |g(z) - w_{R_\varepsilon}| \leq |w_{2R_\varepsilon} - w_{R_\varepsilon}| \psi(\theta, 1) \leq 3R_\varepsilon \psi(\theta, 1)$$

and thus  $|f(z)| \leq R_\varepsilon + 3R_\varepsilon \psi(\theta, 1)$  for all  $|z| \leq \theta$ . Using the hypothesis that  $M_f(\theta) = 1$  and the maximum modulus principle we obtain  $1 \leq R_\varepsilon + 3R_\varepsilon \psi(\theta, 1)$ , and letting  $\varepsilon \rightarrow 0$  we have  $1 \leq r_f + 3r_f \psi(\theta, 1)$ , thus

$$r_f \geq \frac{1}{1 + 3\psi(\theta, 1)} =: C_\theta.$$

■

**Theorem 4.1.4** (Pólya). *Let  $g, h \in H(\mathbb{C})$ . For the order of  $g \circ h$  to be finite, it must hold that either*

- i.  $h$  is a polynomial and  $\rho_g < \infty$ , or
- ii.  $h$  is not a polynomial,  $\rho_h < \infty$  and  $\rho_g = 0$ .

*Proof.* Without loss of generality we can assume  $h(0) = 0$ ; otherwise we just consider  $h_0(z) := h(z) - h(0)$  and  $g_0(w) := g(w + h(0))$ . Set  $f := g \circ h$  and define

$$k_r(z) := \frac{h(rz)}{M_h(r/2)} \in H(\overline{\mathbb{D}}), \quad \text{for } r > 0.$$

Note that by definition we have  $M_{k_r}(1/2) = 1$  and  $k_r(0) = 0$ , thus by Bohr's Theorem there is some constant  $C > 0$  and an  $R/M_h(r/2) > C$  such that  $\partial B_{R/M_h(r/2)}(0) \subseteq k_r(\overline{\mathbb{D}})$  and thus  $\partial B_R(0) \subseteq h(\overline{B_r(0)})$ . By the maximum modulus principle,  $|g|$  assumes its maximum over  $B_R(0)$  at some  $w_0 \in \partial B_R(0)$ . By the above there is a  $z_0 \in B_r(0)$  with  $h(z_0) = w_0$ .

Thus we get

$$M_g(CM_h(r/2)) < M_g(R) = |g(w_0)| = |g(h(z_0))| = |f(z_0)| \leq M_f(r).$$

Assuming  $\rho_f < \infty$ , we have  $M_f(r) < K \exp(r^\alpha)$  for some  $\alpha > \rho_f$ . Consider the power series expansion  $h(z) = \sum_{n=0}^{\infty} a_n z^n$  and let  $a_m$  denote any non-zero coefficient; note that since  $h(0) = 0$  we have  $m \geq 1$ . By Cauchy's integral formula we have, for all  $s > 0$ ,

$$|a_m| = \left| \frac{h^{(m)}(0)}{m!} \right| = \frac{1}{2\pi} \left| \int_{\partial B_s(0)} \frac{h(\zeta)}{\zeta^{m+1}} d\zeta \right| \leq \frac{M_h(s)}{s^m} \quad (*)$$

and thus

$$M_g(C|a_m|(r/2)^m) \leq M_g(CM_h(r/2)) < M_f(r) < K \exp(r^\alpha), \quad \text{for all } r > 0.$$

Replacing  $(r/2)^m$  with  $r$  we obtain  $\rho_g \leq \alpha/m$ . If  $h$  is not a polynomial we may let  $m \rightarrow \infty$ , thus  $\rho_g = 0$ .

Now consider  $g(z) = \sum_{n=0}^{\infty} b_n z^n$ . Replacing  $h$  with  $g$  in  $(*)$  we obtain  $|b_n|s^n \leq M_g(s)$  for all  $s > 0$  and  $n \geq 1$  and thus

$$|b_n|(CM_h(r/2))^n \leq M_g(CM_h(r/2)) < M_f(r) < K \exp(r^\alpha),$$

which implies  $\rho_h \leq \alpha < \infty$ . ■

**Theorem 4.1.5** (Thron). *Let  $g \in H(\mathbb{C})$  be transcendental, of finite order and suppose that  $g$  assumes some value  $w \in \mathbb{C}$  only finitely often. Then there exists no  $f \in H(\mathbb{C})$  with  $f \circ f = g$ .*

*Proof.* Seeking contradiction, suppose there were such a  $f \in H(\mathbb{C})$ . Since  $g$  is not a polynomial,  $f$  is not a polynomial either. Thus Pólya's Theorem implies  $\rho_f = 0$ .

Consider the sets

$$Z := f^{-1}(\{w\}), \quad Z' := \bigcup_{z \in Z} f^{-1}(\{z\}).$$

Note that since pre-images of singletons are disjoint, the union above is actually disjoint. By definition, for each  $z' \in Z'$  there is some  $z \in Z$  with  $z' \in f^{-1}(\{z\})$ , thus

$$g(z') = f(f(z')) = f(z) = w.$$

Our hypothesis on  $g$  implies that  $Z'$  must be finite. Therefore, all points in  $Z$  are only assumed finitely often by  $f$ . But by Corollary 2.3.7,  $f$  assumes all values in the complex plane infinitely often, with at most one exception; therefore  $|Z| \leq 1$ .

If  $Z = \emptyset$ , then  $h(z) := f(z) - w$  is entire, of order 0 and nowhere 0. By Hadamard's Theorem we have  $h = e^Q$  for some constant polynomial  $Q$ . Therefore  $h$  is constant and consequently so is  $f$ , a contradiction.

If  $Z = \{z_0\}$ , then  $h(z) := f(z) - w$  has a single zero of finite order  $n \in \mathbb{N}$  at  $z_0$ . Therefore we can write  $h(z) = (z - z_0)^n p(z)$ , where  $p$  is entire, of order 0 and nowhere 0. Again, this implies that  $p$  is constant, and therefore  $f$  a polynomial, a contradiction. ■

**Example 4.1.6.** A natural application of Thron's Theorem is taking  $g$  to be  $e^z$ , which never assumes zero as a value. Indeed, this implies that there is no entire function  $f$  satisfying

$$f(f(z)) = e^z.$$

On the other hand, there does exist a real-analytic function satisfying the above, as demonstrated by H. Kneser. [I probably still need a citation here.](#)

# Bibliography

- [1] M. Blümlinger. Komplexe Analysis, 2021.
- [2] R. Remmert. *Classical topics in complex function theory*, volume 172 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998. Translated from the German by Leslie Kay.
- [3] S. L. Segal. *Nine introductions in complex analysis*, volume 208 of *North-Holland Mathematics Studies*. Elsevier Science B.V., Amsterdam, revised edition, 2008.
- [4] E. M. Stein and R. Shakarchi. *Complex analysis*, volume 2 of *Princeton Lectures in Analysis*. Princeton University Press, Princeton, NJ, 2003.