

Assignment 7: GLMs (Linear Regressios, ANOVA, & t -tests)

Yimei Du

Fall 2025

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on generalized linear models.

Directions

1. Rename this file <FirstLast>_A07_GLMs.Rmd (replacing <FirstLast> with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

!! AI generated code should NOT be used in this assignment

Set up your session

1. Set up your session. Check your working directory. Load the tidyverse, agricolae and other needed packages. Import the *raw* NTL-LTER raw data file for chemistry/physics (NTL-LTER_Lake_ChemistryPhysics_Raw.csv). Set date columns to date objects.
2. Build a ggplot theme and set it as your default theme.

#1

```
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
```

```
## v dplyr      1.1.4      v readr      2.1.5
```

```
## v forcats    1.0.0      v stringr    1.5.1
```

```
## v ggplot2    4.0.0      v tibble     3.2.1
```

```
## v lubridate  1.9.3      v tidyr      1.3.1
```

```
## v purrr      1.0.2
```

```
## -- Conflicts ----- tidyverse_conflicts() --
```

```
## x dplyr::filter() masks stats::filter()
```

```
## x dplyr::lag()     masks stats::lag()
```

```
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors
```

```
library(lubridate)
```

```
library(agricolae)
```

```
library(here)
```

```
## here() starts at /home/guest/EDE_Fall2025
```

```
getwd()
```

```
## [1] "/home/guest/EDE_Fall2025"
```

```
here() #check working directory
```

```
## [1] "/home/guest/EDE_Fall2025"
```

```
chemphy <- read.csv(  
  file = here('Data', 'Raw', 'NTL-LTER_Lake_ChemistryPhysics_Raw.csv'),  
  stringsAsFactors = T)
```

```
chemphy$sampleddate <- mdy(chemphy$sampleddate)
```

```
str(chemphy$sampleddate)
```

```
## Date[1:38614], format: "1984-05-27" "1984-05-27" "1984-05-27" "1984-05-27" "1984-05-27" ...
```

```
#2  
mytheme <- theme_classic(base_size = 14) +  
  theme(axis.text = element_text(color = "black"),  
        legend.position = "top")
```

Simple regression

Our first research question is: Does mean lake temperature recorded during July change with depth across all lakes?

3. State the null and alternative hypotheses for this question: > Answer: H0: There is no difference in mean July lake temperature among different depths across all lakes. Ha: There is difference in mean July lake temperature among different depths across all lakes.
4. Wrangle your NTL-LTER dataset with a pipe function so that the records meet the following criteria:
 - Only dates in July.
 - Only the columns: lakename, year4, daynum, depth, temperature_C
 - Only complete cases (i.e., remove NAs)
5. Visualize the relationship among the two continuous variables with a scatter plot of temperature by depth. Add a smoothed line showing the linear model, and limit temperature values from 0 to 35 C. Make this plot look pretty and easy to read.

```
#4  
library(dplyr)  
  
tidy_chemphy <- chemphy %>%  
  filter(month(sampleddate) == 7) %>%
```


```

select(lakename, year4, daynum, depth, temperature_C) %>%
na.omit()

#5
sp <-
  ggplot(tidy_chemphy, aes(x = temperature_C, y = depth)) +
  xlim(0, 35) +
  geom_point(size = 0.5) +
  geom_smooth(method = "lm", col = "orange") +
  scale_y_reverse() +
  labs(
    x = "Temperature (C)",
    y = "Depth (m)",
    title = "Temperature vs Depth (July)"
  ) +
  geom_text(
    x = 18,
    y = 1,
    label = "temperature= -1.95*depth+21.96",
    size = 3
  )
print(sp)

```

'geom_smooth()' using formula = 'y ~ x'



```
lr <-
  lm(data = tidy_chemphy, temperature_C ~ depth)
summary(lr)

##
## Call:
## lm(formula = temperature_C ~ depth, data = tidy_chemphy)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.5173 -3.0192  0.0633  2.9365 13.5834
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 21.95597    0.06792   323.3  <2e-16 ***
## depth       -1.94621    0.01174  -165.8  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.835 on 9726 degrees of freedom
## Multiple R-squared:  0.7387, Adjusted R-squared:  0.7387
## F-statistic: 2.75e+04 on 1 and 9726 DF, p-value: < 2.2e-16
```

6. Interpret the figure. What does it suggest with regards to the response of temperature to depth? Do the distribution of points suggest about anything about the linearity of this trend?

Answer: This figure shows a negative relationship between temperature and depth. As depth increases, lake temperature decreases. The slope of the orange line (-1.95 C per meter) suggests that temperature decreases by about 2 C for every additional meter of depth, on average. The distribution of the points show some variability on the surface and as deeper as it goes, the more clustered the points are.

7. Perform a linear regression to test the relationship and display the results.

```
#7
lr <-
  lm(data = tidy_chemphy, temperature_C ~ depth)
summary(lr)

##
## Call:
## lm(formula = temperature_C ~ depth, data = tidy_chemphy)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.5173 -3.0192  0.0633  2.9365 13.5834
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 21.95597    0.06792   323.3  <2e-16 ***
## depth       -1.94621    0.01174  -165.8  <2e-16 ***
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.835 on 9726 degrees of freedom
## Multiple R-squared:  0.7387, Adjusted R-squared:  0.7387
## F-statistic: 2.75e+04 on 1 and 9726 DF,  p-value: < 2.2e-16
```

8. Interpret your model results in words. Include how much of the variability in temperature is explained by changes in depth, the degrees of freedom on which this finding is based, and the statistical significance of the result. Also mention how much temperature is predicted to change for every 1m change in depth.

Answer: The model shows a strong, statistically significant negative relationship between lake temperature and depth. For every 1 meter increase in depth, temperature decreases by about 1.95 C. The model explains roughly 74 % of the variation in temperature based on 9726 degrees of freedom, and the relationship is highly significant ($p < 2.2 \times 10^{-16}$).

Multiple regression

Let's tackle a similar question from a different approach. Here, we want to explore what might the best set of predictors for lake temperature in July across the monitoring period at the North Temperate Lakes LTER.

9. Run an AIC to determine what set of explanatory variables (year4, daynum, depth) is best suited to predict temperature.
10. Run a multiple regression on the recommended set of variables.

```
#9
LTERAIC <- lm(temperature_C ~ year4 + daynum + depth, data = tidy_chemphy)
step(LTERAIC)

## Start:  AIC=26065.53
## temperature_C ~ year4 + daynum + depth
##
##           Df Sum of Sq    RSS   AIC
## <none>                 141687 26066
## - year4      1         101 141788 26070
## - daynum     1         1237 142924 26148
## - depth      1      404475 546161 39189

##
## Call:
## lm(formula = temperature_C ~ year4 + daynum + depth, data = tidy_chemphy)
##
## Coefficients:
## (Intercept)      year4      daynum      depth
##    -8.57556     0.01134     0.03978    -1.94644
```

```
#10
LTER_multi <- lm(temperature_C ~ year4 + daynum + depth, data = tidy_chemphy)

summary(LTER_multi)

##
## Call:
## lm(formula = temperature_C ~ year4 + daynum + depth, data = tidy_chemphy)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6536 -3.0000  0.0902  2.9658 13.6123
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) -8.575564   8.630715  -0.994  0.32044
## year4         0.011345   0.004299   2.639  0.00833 **
## daynum        0.039780   0.004317   9.215 < 2e-16 ***
## depth        -1.946437   0.011683 -166.611 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.817 on 9724 degrees of freedom
## Multiple R-squared:  0.7412, Adjusted R-squared:  0.7411
## F-statistic: 9283 on 3 and 9724 DF,  p-value: < 2.2e-16
```

11. What is the final set of explanatory variables that the AIC method suggests we use to predict temperature in our multiple regression? How much of the observed variance does this model explain? Is this an improvement over the model using only depth as the explanatory variable?

Answer: The best model includes year4 + daynum + depth, explains about 74% of temperature variability. There is a slight increase compare to only depth but depth still remain as the dominant factor.

Analysis of Variance

12. Now we want to see whether the different lakes have, on average, different temperatures in the month of July. Run an ANOVA test to complete this analysis. (No need to test assumptions of normality or similar variances.) Create two sets of models: one expressed as an ANOVA models and another expressed as a linear model (as done in our lessons).

```
#12
anova <- aov(data = tidy_chemphy, temperature_C ~ lakename)
summary(anova)

##              Df Sum Sq Mean Sq F value Pr(>F)
## lakename       8  21642   2705.2     50 <2e-16 ***
## Residuals    9719 525813     54.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
linear <- lm(data = tidy_chemphy, temperature_C ~ lakename)
summary(linear)
```

```
##
## Call:
## lm(formula = temperature_C ~ lakename, data = tidy_chemphy)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -10.769  -6.614  -2.679   7.684  23.832
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    17.6664     0.6501  27.174 < 2e-16 ***
## lakenameCrampton Lake    -2.3145     0.7699   -3.006 0.002653 **
## lakenameEast Long Lake   -7.3987     0.6918 -10.695 < 2e-16 ***
## lakenameHummingbird Lake -6.8931     0.9429  -7.311 2.87e-13 ***
## lakenamePaul Lake       -3.8522     0.6656  -5.788 7.36e-09 ***
## lakenamePeter Lake      -4.3501     0.6645  -6.547 6.17e-11 ***
## lakenameTuesday Lake    -6.5972     0.6769  -9.746 < 2e-16 ***
## lakenameWard Lake       -3.2078     0.9429  -3.402 0.000672 ***
## lakenameWest Long Lake  -6.0878     0.6895  -8.829 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7.355 on 9719 degrees of freedom
## Multiple R-squared:  0.03953,    Adjusted R-squared:  0.03874
## F-statistic:    50 on 8 and 9719 DF,  p-value: < 2.2e-16
```

13. Is there a significant difference in mean temperature among the lakes? Report your findings.


Answer: There is significant difference in mean July temperature across lakes ($F(8, 9719) = 50$, $p < 2 \times 10^{-16}$). The p value is way below the threshold 0.05.

14. Create a graph that depicts temperature by depth, with a separate color for each lake. Add a `geom_smooth` (method = "lm", se = FALSE) for each lake. Make your points 50 % transparent. Adjust your y axis limits to go from 0 to 35 degrees. Clean up your graph to make it pretty.

```
#14.
library(ggplot2)
plot <- ggplot(tidy_chemphy, aes(x = depth, y = temperature_C, color = lakename)) +
  geom_point(alpha = 0.5, size = 1) +
  geom_smooth(method = "lm", se = FALSE, linewidth = 1) +
  coord_cartesian(ylim = c(0, 35)) +
  labs(
    x = "Depth (m)",
    y = "Temperature (C)",
    color = "Lake",
    title = "Temperature by Depth (July)"
  ) +
  mytheme
print(plot)
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```

Temperature by Depth (July)



15. Use the Tukey's HSD test to determine which lakes have different means.

```
#15
```

```
TukeyHSD(anova)
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = temperature_C ~ lakename, data = tidy_chemphy)
##
## $lakename
##
```

	diff	lwr	upr	p adj
## Crampton Lake-Central Long Lake	-2.3145195	-4.7031913	0.0741524	0.0661566
## East Long Lake-Central Long Lake	-7.3987410	-9.5449411	-5.2525408	0.0000000
## Hummingbird Lake-Central Long Lake	-6.8931304	-9.8184178	-3.9678430	0.0000000
## Paul Lake-Central Long Lake	-3.8521506	-5.9170942	-1.7872070	0.0000003
## Peter Lake-Central Long Lake	-4.3501458	-6.4115874	-2.2887042	0.0000000
## Tuesday Lake-Central Long Lake	-6.5971805	-8.6971605	-4.4972005	0.0000000
## Ward Lake-Central Long Lake	-3.2077856	-6.1330730	-0.2824982	0.0193405
## West Long Lake-Central Long Lake	-6.0877513	-8.2268550	-3.9486475	0.0000000
## East Long Lake-Crampton Lake	-5.0842215	-6.5591700	-3.6092730	0.0000000
## Hummingbird Lake-Crampton Lake	-4.5786109	-7.0538088	-2.1034131	0.0000004
## Paul Lake-Crampton Lake	-1.5376312	-2.8916215	-0.1836408	0.0127491

## Peter Lake-Crampton Lake	-2.0356263	-3.3842699	-0.6869828	0.0000999
## Tuesday Lake-Crampton Lake	-4.2826611	-5.6895065	-2.8758157	0.0000000
## Ward Lake-Crampton Lake	-0.8932661	-3.3684639	1.5819317	0.9714459
## West Long Lake-Crampton Lake	-3.7732318	-5.2378351	-2.3086285	0.0000000
## Hummingbird Lake-East Long Lake	0.5056106	-1.7364925	2.7477137	0.9988050
## Paul Lake-East Long Lake	3.5465903	2.6900206	4.4031601	0.0000000
## Peter Lake-East Long Lake	3.0485952	2.2005025	3.8966879	0.0000000
## Tuesday Lake-East Long Lake	0.8015604	-0.1363286	1.7394495	0.1657485
## Ward Lake-East Long Lake	4.1909554	1.9488523	6.4330585	0.0000002
## West Long Lake-East Long Lake	1.3109897	0.2885003	2.3334791	0.0022805
## Paul Lake-Hummingbird Lake	3.0409798	0.8765299	5.2054296	0.0004495
## Peter Lake-Hummingbird Lake	2.5429846	0.3818755	4.7040937	0.0080666
## Tuesday Lake-Hummingbird Lake	0.2959499	-1.9019508	2.4938505	0.9999752
## Ward Lake-Hummingbird Lake	3.6853448	0.6889874	6.6817022	0.0043297
## West Long Lake-Hummingbird Lake	0.8053791	-1.4299320	3.0406903	0.9717297
## Peter Lake-Paul Lake	-0.4979952	-1.1120620	0.1160717	0.2241586
## Tuesday Lake-Paul Lake	-2.7450299	-3.4781416	-2.0119182	0.0000000
## Ward Lake-Paul Lake	0.6443651	-1.5200848	2.8088149	0.9916978
## West Long Lake-Paul Lake	-2.2356007	-3.0742314	-1.3969699	0.0000000
## Tuesday Lake-Peter Lake	-2.2470347	-2.9702236	-1.5238458	0.0000000
## Ward Lake-Peter Lake	1.1423602	-1.0187489	3.3034693	0.7827037
## West Long Lake-Peter Lake	-1.7376055	-2.5675759	-0.9076350	0.0000000
## Ward Lake-Tuesday Lake	3.3893950	1.1914943	5.5872956	0.0000609
## West Long Lake-Tuesday Lake	0.5094292	-0.4121051	1.4309636	0.7374387
## West Long Lake-Ward Lake	-2.8799657	-5.1152769	-0.6446546	0.0021080

```
tukey <- with(tidy_chemphy, interaction(lakename, temperature_C))
```

16. From the findings above, which lakes have the same mean temperature, statistically speaking, as Peter Lake? Does any lake have a mean temperature that is statistically distinct from all the other lakes?

Answer: Peter Lake has statistically similar mean temperatures to Paul Lake and Ward Lake. East Long Lake has a significantly different mean temperature from almost all other lakes.

17. If we were just looking at Peter Lake and Paul Lake. What's another test we might explore to see whether they have distinct mean temperatures?

Answer: A two sample t-test could be used to test the mean difference between Peter Lake and Paul Lake.

18. Wrangle the July data to include only records for Crampton Lake and Ward Lake. Run the two-sample T-test on these data to determine whether their July temperature are same or different. What does the test say? Are the mean temperatures for the lakes equal? Does that match your answer for part 16?

```
july_twolake <- tidy_chemphy %>%
  filter(lakename %in% c("Crampton Lake", "Ward Lake"))

t.test(temperature_C ~ lakename, data = july_twolake)
```

```
##
```

```
## Welch Two Sample t-test
##
## data:  temperature_C by lakename
## t = 1.1181, df = 200.37, p-value = 0.2649
## alternative hypothesis: true difference in means between group Crampton Lake and group Ward Lake is not equal to 0
## 95 percent confidence interval:
##  -0.6821129  2.4686451
## sample estimates:
## mean in group Crampton Lake      mean in group Ward Lake
##                15.35189                14.45862
```

Answer: A two-sample t-test comparing July temperatures at Lake Crampton (mean = 15.35 C) and Ward Lake (mean = 14.46 C) yielded a p-value of 0.26, indicating no significant difference in average temperatures between the two lakes. This result is consistent with Tukey's multiple comparison test in part 16, which similarly showed no significant difference ($p = 0.97$).