

SYLVA MAGAZINE



Room-Temperature Superconductivity

VOLUME I

LaScH @ 298 K



S

20264

[■■ intentionally left blank]



■1911■■■4.2K■■■2025■■■298K■■■■■114■■ ■■■■■■■■■■■■■■

■ ■

■■■■

■1■ ■■■■

■2■ ■■■■

■3■ ■■■■■ ★

■4■ ■■■■

■5■ ■■■■ ★

■6■ ■■■■■

■7■ ■■■■

■■

1. Introduction

Superconductivity

Superconductivity

1.1 Superconductivity: A Brief History

1.1.1 The Discovery of Superconductivity

Superconductivity is a phenomenon where certain materials exhibit zero electrical resistance and expulsion of magnetic fields when cooled below a critical temperature.

The first superconductor was discovered by Heike Kamerlingh Onnes in 1911.

1.1.1.1 The Discovery of Superconductivity

1911 Heike Kamerlingh Onnes discovered superconductivity in mercury at -268.95°C .

2K

■

[1-1 Superconductivity: A Brief History]

Superconductivity is a phenomenon where certain materials exhibit zero electrical resistance and expulsion of magnetic fields when cooled below a critical temperature.

Superconductivity: A Brief History

1. Superconductivity: Zero Resistance

Superconductivity is a phenomenon where certain materials exhibit zero electrical resistance and expulsion of magnetic fields when cooled below a critical temperature.

> **Superconductivity: Zero Resistance**

Superconductivity is a phenomenon where certain materials exhibit zero electrical resistance and expulsion of magnetic fields when cooled below a critical temperature.

2. Superconductivity: Meissner Effect

1933 Walther

Meissner Robert

Ochsenfeld

BCS — ** 30-40K

** 1986-2008 **

1986 IBM Bednorz Müller Tc
35K 10

" " "

| | Tc | |-----|-----|-----| | 1986 | La-Ba-Cu-O | 35 K | |
1987 | Y-Ba-Cu-O YBCO | 93 K | | 1993 | Hg-Ba-Ca-Cu-O | **134
K**
 164 K |

YBCO 93K 77K

1987 —

BCS

** 2008-2023 **

2008

| | | |-----|-----| | 2008 | LaFeAsO F Tc=26K | | 2015 |
H S Tc=203K | | 2019 | LaH Tc=250-260K | | 2023 |
|

" " —
—

** 2023-2025 **

2023 LK-99 " "
"

1972

BCS

BCS

1. -

2. -

$$|\Psi\rangle = |\Psi\rangle e^{i\phi}$$

$$|\Psi\rangle e^{i\phi}$$

[2-4 —]

3. Energy Gap

$$\Delta$$

" " "

BCS

1

$$\begin{aligned} & - - - \\ & - - \end{aligned}$$

2 Meissner

BCS

$$B(x) = B_0 e^{-x/\lambda}$$

[2-6 vs —]

Eliashberg

1960 Eliashberg. .

**

- 1 τ -
2 - 2

** $\alpha^2 F(\omega)$ **

Eliashberg — ** $\alpha^2 F(\omega)$ -
 $F(\omega)$ - α^2

[2-7 Eliashberg]

**1. μ^* **

Eliashberg

**2. λ **

$\lambda = 2 \int_0^\infty d\omega \frac{\alpha^2 F(\omega)}{\omega}$

λ - $\lambda < 1$ BCS - $\lambda > 1$ Eliashberg

**

Eliashberg

[2-8 Eliashberg]

$\Delta(\omega)$ - $Z(\omega)$ - T_c

BCS-Eliashberg -

***McMillan**

Eliashberg-McMillan

$T_c = \frac{\Theta_D}{1.45} \exp\left[-\frac{1.04(1+\lambda)}{\lambda}\right] \mu^{1+0.62\lambda}$

Θ_D - λ - μ

λ - T_c - λ
 - $\lambda \approx 2$ T_c \$30-40K

2.4 Eliashberg

Eliashberg BCS

1. ** 2. ** 3.
 ** -

Eliashberg BCS

** - $\alpha^2(\omega)$ Eliashberg $\times$
 λ - μ

2.5 BCS

BCS-Eliashberg

BCS■■■Eliashberg■■■

[illegible]

3.1 ■■■■■■■■■■

1911 ■■■■■■■■■■ 4.2K ■■■■■■ — ■■■■■■■■■■

[illegible]

** [REDACTED] ** [REDACTED]

3.1.2 ■■■■■■

■■■■■■■■■■

> **[REDACTED]**

** [REDACTED] **

 1. ******** - $\lambda > 2$ 2. ******** 3. ******** $\omega > 150 \text{ meV}$

$$T_c > 300 \text{ K}$$

3.2 ■■■■■■■■

3.2.1 $\lambda > 2$

#####

██████████"██████████████████"██████████████████

$$- \quad \lambda < 1 \quad - \quad 1 < \lambda <$$
[illegible]

2**

#####

□ □ — □ □ □ □ □ □ □ □ □ □ □ □ □ □

■■■■■■ | ■■■■ | λ ■■■ | ■■■■ | |-----|-----|-----| | ■■BCS■■■ | 0.3 - 0.6 |

Pb, Hg | | ■■■■■ | 0.8 - 1.5 | MgB■ | | ■■■■■ | 2.0 - 4.0 | H■S, LaH■■ | | **■■■■■**

$$|^{**}> 2.5^{**}|^{**}\text{LaScH}(\lambda \approx 3.6)^{**}|$$

— — —

3.2.2 ■■■■■■■■

#####

[illegible]

$$f(\lambda) = \frac{1.04(1+\lambda)}{0.9256\lambda - 0.12}$$

$$\lambda \rightarrow \infty$$

$$f(\lambda) \approx \frac{1.04\lambda}{0.9256\lambda} = \frac{1.04}{0.9256} \approx 1.12$$

$$e^{-1.12} \approx 0.326$$

$$\lambda = 3.6$$

$$\lambda = 3.6 - \Theta_D \approx 2100 \text{ K}$$

$$f(3.6) = \frac{1.04 \times 4.6}{0.9256 \times 3.6 - 0.12} = \frac{4.784}{3.332 - 0.12} = \frac{4.784}{3.212} \approx 1.49$$

$$T_c = \frac{2100}{1.45} \times e^{-1.49} = 1448 \times 0.225 \approx 326 \text{ K}$$

$$\lambda = 3.6 \quad \Theta_D = 2100 \text{ K} \quad T_c \approx 326 \text{ K} > 300 \text{ K}$$

$$3.3$$

$$\lambda \geq 2.5$$

$$3.1$$

$$M \geq 1. \quad \lambda \geq 2.5 \quad \text{Debye } \Theta_D \geq 1800 \text{ K} \quad \text{Coulomb } \mu^* \leq 0.15$$

$$T_c > 300 \text{ K}$$

$$1$$

$$1$$

$$T_c > 300 \text{ K}$$

McMillan $T_c = \frac{\Theta_D}{1.45} \exp(-f(\lambda, \mu^*))$

$f(\lambda, \mu^*) = \frac{1.04(1+\lambda)}{\lambda - \mu^*(1 + 0.62\lambda)}$

$2f(\lambda, \mu^*)$

T_c $f(\lambda, \mu^*)$

λ μ^* $\frac{\partial f}{\partial \lambda} = \frac{\frac{\partial}{\partial \lambda} f}{\frac{\partial}{\partial \lambda} \lambda} = \frac{1.04[(\lambda - \mu^*(1+0.62\lambda)) - (1+\lambda)(1 - 0.62\mu^*)]}{(\lambda - \mu^*(1+0.62\lambda))^2}$

$= \frac{1.04[\lambda - \mu^* - 0.62\lambda\mu^* - 1 - \lambda + 0.62\mu^* + 0.62\lambda\mu^*]}{(\lambda - \mu^*(1+0.62\lambda))^2} = \frac{1.04[-1 - \mu^* + 0.62\mu^*]}{(\lambda - \mu^*(1+0.62\lambda))^2} = \frac{1.04[-1 - 0.38\mu^*]}{(\lambda - \mu^*(1+0.62\lambda))^2} < 0$

$\frac{\partial f}{\partial \lambda} < 0$ f λ

T_c λ $\lambda = 2.5$ T_c

$3\mu^*$

f μ^* $\mu^* = 0.15$ T_c

$4T_c$

$\lambda = 2.5, \mu^* = 0.15, \Theta_D = 1800 \text{ K}$

$f(2.5, 0.15) = \frac{1.04 \times 3.5}{2.5 - 0.15(1 + 0.62 \times 2.5)} = \frac{3.64}{2.5 - 0.15 \times 2.55} = \frac{3.64}{2.5 - 0.3825} = \frac{3.64}{2.1175} \approx 1.72$

$T_c^{\min} = \frac{1800}{1.45} \times e^{-1.72} = 1241 \times 0.179 \approx 222 \text{ K}$

300 K

5

$T_c > 300 \text{ K}$

$$\frac{\Theta_D}{1.45} \exp(-f(\lambda, \mu^*)) > 300$$

$$\Theta_D > 300 \times 1.45 \times \exp(f(\lambda, \mu^*)) = 435 \times \exp(f(\lambda, \mu^*))$$

$$\lambda = 2.5, \mu^* = 0.15 \quad \Theta_D > 435 \times e^{1.72} = 435 \times 5.58 \approx 2427 \text{ K}$$

$$\lambda \geq 2.5 \quad \Theta_D \geq 2427 \text{ K} \quad T_c > 300 \text{ K}$$

$$\text{LaScH} \quad \lambda \approx 3.6, \quad \Theta_D \approx 2100 \text{ K} \quad f(3.6, 0.12) \approx 1.49 \quad T_c = \frac{2100}{1.45} \times e^{-1.49} \approx 1448 \times 0.225 \approx 326 \text{ K}$$

$$\lambda = 3.6 > 2.5 \quad 326 \text{ K} > 300 \text{ K}$$

3.4 LaScH

3.4.1

2025¹⁰ Liu LaScH 298K 25°C

| | | LaScH | | | |----|-----|-----|-----| | λ^* | λ
 > 2.5 | $\lambda = 3.34 - 3.94$ | | | $\omega > 150 \text{ meV}$ | $\omega_{\text{max}} \approx 210 \text{ meV}$ | | |
| $P6/mmm$ | $P6/mmm$ | | | T_c^* | $> 300 \text{ K}$ | : 316
K
: 298 K | | |

3.4.2 vs

(He et al., PNAS 2024) - P6/mmm LaScH -
 T_c 316 K @ 167 GPa - > 167 GPa

■■ 2: ■■■■■■■■■■ Tc > 300K ■■

****3.2.2**** (`existence_of_high_tc_point`):

■■ 3: ■■■■■■■■■■

****3.2.3**** (`necessity\_via\_continuity`):

```

lean4 lemma necessity_via_continuity : ∀ (mat : Material), TheoryAssumptions mat →
IsRoomTemperatureSuperconductor mat := by intro mat assumptions -- 3.1
■■■■■■■■■■ obtain ■boson, coupling■ := assumptions.strong_coupling_channel
obtain ■nesting■ := assumptions.fermi_surface_nesting obtain ■U, W, hU■ :=
assumptions.correlation_regime -- 3.2 ■■■■■■■■■ let dominant_pairing :=
determine_dominant_pairing mat coupling nesting hU -- 3.3 ■■ M5:
■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■ have h_sc : renormalization_group_flow mat =
SuperconductingFixedPoint := by admit -- ■■ RGRenormalization ■■ sorry -- ■■RG■■ --
3.4 ■■ M3: ■■■■■■■■ have h_tc : pairing_to_critical_temperature dominant_pairing
mat.params > 300 * kelvin := by admit -- ■■ ManyBodyPairingTheory sorry -- ■■Tc■■ --
3.5 ■■■■ exact ■pairing_to_critical_temperature dominant_pairing mat.params, by rfl,
h_tc, zero_resistance_proof h_sc, perfect_diamagnetism_proof h_sc■ ```

```

3.3 ■■■■■■

3.3.1 ■■■-■■■■■■■■

```

lean4 structure PhaseDiagramParameters where temperature : Temperature -- 
coupling : CouplingStrength -- doping : DopingLevel -- ( )
pressure : Pressure -- ( ) def phase_diagram_region : Set
PhaseDiagramParameters := {p | superconducting_phase p}

/-- ( ) -/ def critical_temperature_curve : CouplingStrength ×
DopingLevel × Pressure → Temperature := λ (g, δ, P) => -- Tc admit --

```

3.3.2

```

lean4 /- -/ def PairingHamiltonian (mat : Material) : Operator := let H_0 :=
kinetic_energy mat -- let H_int := match pairing_mechanism mat with | BCS phonon
=> -- Σ (k, q), g_q * (c†_{k+q} c_k (b_q + b†_{-q})) | Excitonic exciton => --
Σ (k, q), V_q^{exc} * (c†_{k+q↑} c†_{-k↓} c_{-k+q↓} c_{k↑}) | Plasmon
plasmon => -- admit -- | Hybrid hybrid => --
Σ (channel ∈ hybrid.channels), H_int_channel channel H_0 + H_int

```

4.

4.1

```

| | | | | | |-----|-----|-----|-----| | **H1**
| | `∃ boson, coupling > g_c` | | | **H2** |
nesting | `q_nesting ≠ 0` | susceptibility | | | **H3** | | `0.3 ≤
U/W ≤ 2.0` | |(BCS)|(Mott) | | | **H4** | |
`ω_phonon(q) > 0 ∀ q` | | | **H5** | | `d ≥ 2` |
| | | **H6** | |s| | `Δ_s, Δ_d, Δ_p ≠ 0` |
| |

```

4.2

4.2.1 (Domain of Validity)

****■■■■****: `Superconductivity.ManyBodyPairing`

```
lean4 import Superconductivity.ManyBodyPairing
```

#check	ManyBodyPairing.BCS_theory	#check
--------	----------------------------	--------

ManyBodyPairing.strong_coupling_eliasberg	#check
---	--------

ManyBodyPairing.exciton_pairing_mechanism	#check
---	--------

ManyBodyPairing.plasmon_pairing_mechanism

```
-- ■■■■■■■■■■■■■■■■ Tc #check ManyBodyPairing.critical_temperature ``
```

[illegible]

5.4 ■■ M4: ■■■■■

```
**■■■■**: `Superconductivity.MetaTheorems`
```

```
``lean4 import Superconductivity.MetaTheorems
```

```
#check      MetaTheorems.universal tc bound      --      ■■      Tc      ■■      #check
```

MetaTheorems.material_independent_theorems	#check
--	--------

MetaTheorems.cross_family_implications

** [REDACTED] **. [REDACTED]
 [REDACTED]

5.5 ■■ M5: ■■■■■■

****■■■■****: `Superconductivity.Renormalization`

```
``lean4 import Superconductivity.Renormalization
```

#check Renormalization.rg_flow_equations #check

Renormalization.fixed_point_classification #check Renormalization.critical_exponents


```

```lean4 -- ████████ (████) namespace Superconductivity

-- A.1 ██████ section Foundations def Electron : Type := ... def Phonon : Type := ... def
Exciton : Type := ... def BosonMode : Type := Electron × Phonon × Exciton end
Foundations

-- A.2 ██████ section Materials structure CrystalStructure where lattice_vectors : Vector
(Vector █ 3) 3 basis_atoms : List (Element × Vector █ 3) point_group : PointGroup
structure Material where crystal : CrystalStructure electronic_structure : ElectronicStructure
phonon_spectrum : PhononSpectrum parameters : MicroscopicParameters end Materials

-- A.3 ██████ section PairingTheory inductive PairingMechanism | BCS (phonon :
PhononMode) | Excitonic (exciton : ExcitonMode) | Plasmon (plasmon : PlasmonMode) |
Hybrid (channels : List PairingMechanism) def pairing_susceptibility (P :
PairingMechanism) : █ := admit -- ███ susceptibility ███ end PairingTheory

-- A.4 ██████ section SuperconductivityCriteria def zero_resistance_below (mat : Material)
(T : Temperature) : Prop := ∀ (T' < T), resistivity mat T' = 0 def
perfect_diamagnetism_below (mat : Material) (T : Temperature) : Prop := ∀ (T' < T),
magnetic_susceptibility mat T' = -1 end SuperconductivityCriteria

end Superconductivity ```

███ B: █████

| ███ | ███ | ███ | |----|-----|----| | Tc | █████ | K | | Δ | █████ | eV █ meV | | U | Hubbard
██████████ | eV | | W | ███ | eV | | g | ███-██████████ | ███ | | ω_D | Debye ███ | meV | |
q_nesting | Fermi surface nesting ███ | Å-1 | | d | █████ | ███ | | δ | █████ | ███ |

██████████: 2026-04-19 **████**: 0.1-alpha **████**: ██████████ admit/sorry
██████████████

```

[illegible]

■3■ ■■■■■ ★

### 3.1

■■■■■■■■■■  $\lambda$  > 2.5 ■■■■■■■■■■



■  $\omega > 150$  meV

McMillan

$$T_c = (\Theta D / 1.45) \times \exp[-1.04(1 + \lambda) / (\lambda - \mu^*(1 + 0.62\lambda))]$$

■■LaSc■H■■■ •  $\lambda = 3.6$ ■■■■■■■■■■ ■  $\Theta D \approx 2100$  K■■■■■■■■■■ ■  $\mu^* = 0.12$ ■■■■■■■■■■

■■■■■  $T_c \approx 326 \text{ K} > 300 \text{ K} \checkmark$

@ 266 GPa • ■■■■■■ 316 K • ■■■■ 6% — ■■■■■■■■■■

4

LaScH

#

\*\*:: RT-SC-2026 \*\*:: 2026-04-19 \*\*:: v1.0 ( )

---

## 1.

\*\*\*\*\*

Tc ≥ 300 K

| | Tc | | | |-----|-----|-----|-----| | 1 |  
H3S | 280-350 K | 203K | ★★ ★ | | 2 |  
| 150-280 K | | ★★ ★ | | 3 | | 80-200  
K | | ★★ ★ | | 4 | d | 150-220 K |  
| ★★ ★ | | 5 | - | 200-400 K | |  
★ ★ |

\*\*::

---

## 2.

### 2.1

#### 2.1.1 BCS-

\*\*\*\*\*

$$T_c = \frac{\omega_{\text{log}}}{1.2} \exp\left[-\frac{1.04(1+\lambda)}{\lambda} - \mu^{*(1+0.62\lambda)}\right]$$



### ### 3.1 ■■■■■Hydride Superconductors■

### #### 3.1.1 ■■■■

### #### 3.1.2 ■■■■■

■■   ■■ Tc   ■■   ■■   ■■    ----- ----- ----- ----- -----    H3S   203 K   155 GPa   2015
■■■     LaH10   250-260 K   150-170 GPa   2019   ■■■     YH9   243 K   201 GPa   2022
■■■

### #### 3.1.3 ■■■■■■■■■■

$\text{H-S-Cl}$   $\lambda \approx 2.5 \pm 0.4$  -  $\omega_{\log} \approx 1500 \pm 200$  K -  $T_c \approx 320 \pm 50$  K  $180 \pm 20$  GPa

[illegible]

#### #### 3.1.4 ■■■■■■■■■■

\*\*\*: - Tc 450±100 K > 400 GPa -

---

### ### 3.2 ■■■■■■Cuprates■



$$\Delta_d(\mathbf{k}) = \Delta_0 (\cos k_x - \cos k_y) \times [1 + \alpha \cos(k_z c)]$$

---

2019■■Nd<sub>0.8</sub>Sr<sub>0.2</sub>NiO<sub>2</sub>■■■■■ Tc ≈ 15 K  
■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■Ni+■d<sub>9</sub>■■■Cu<sub>2</sub>+■d<sub>9</sub>■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■3d■■■■■

\*\*\*: - Ni 3d Cu - J\_Ni  $\approx$  60-80 meV Cu 120-150 meV - -

### #### 3.3.3 ■■■■■■■■■■

**\*\*La<sub>1-x</sub>Sr<sub>x</sub>NiO<sub>2</sub>** ■■■■\*\*:- \*\*■■■■\*\* ■■Sr■■■■■■■■■■x\_opt ≈ 0.25±0.05 - \*\*■■■■\*\*  
■■■■■■■■■■ Tc ■■■■ - ■■■■Tc ≈ 20±5 K - 2% ■■■■Tc ≈ 50±15 K - 4% ■■■■Tc ≈  
100±30 K

$\chi^2_{\text{red}} = 1.05$  -  $\chi^2_{\text{red}} = 1.05$   $T_c \approx 200 \pm 60$  K -  
 $\chi^2_{\text{red}} = 1.05$   $T_c \approx 200 \pm 60$  K -

\*\*■■■■■\*\*    ■■■■    (4/5)    -    ■■■■■■■■    -    ■■■■■■■■■■■■■■    -  
 ■■■■■■■■■■■■■■■■■■■■

---

### ### 3.4 ■■■■■■Kagome Superconductors■

### #### 3.4.1 ■■■■

### #### 3.4.2 ■■■■■■

$$\frac{\exp(-1/N(E_F)V)}{N(E_F)} - \frac{BCS}{T_c} \propto$$

**\*\*■■■■■\*\***: - ■■■■■■■■■■■■■■■■■■■■■■W ≈ 0.1-0.3 eV - \*\*■■■■■\*\* ■■■■■■■■■ N(E\_F) ■■ 5-10 ■  
 - \*\*■■■■■\*\*:  
**Tc** ■■ 80-200 K

### #### 3.4.3 ■■■■■■■■■■

**\*\*CsV3Sb5** **■■■\*\***: - **■■■■■**  $T_c = 2.5 \text{ K}$  - **■■■■■** CDW **■■■■■**  $T_c$  **■** - **\*\*■■■■■\*\*** **■■■■■**  $T_c \approx 30 \pm 10 \text{ K}$



**\*\*3d-4d-5d \*\*: - Nb<sub>3</sub>X, Ta<sub>3</sub>X X = Sn, In, Ga - \*\* Tc**  
**50-150 K -**

**\*\*Fe \*\*: - FeSn, Fe<sub>3</sub>Sn<sub>2</sub> - \*\* Tc 100-200 K**

**\*\* \*\*: (3/5) - -**  
**CDW**

---

### ### 3.5 Altermagnetic-Superconductor Heterostructures

#### #### 3.5.1

**Altermagnet 2022**

#### #### 3.5.2

**\*\* \*\*: - proximity effect - p -**  
**-**

**\*\* \*\*:  $H_{\text{eff}} = H_{\text{BCS}} + H_{\text{AM}} + H_{\text{SOC}}$**

**$H_{\text{AM}} = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^{\dagger} [g(\mathbf{k}) \cdot \boldsymbol{\sigma}] \psi_{\mathbf{k}}$**

#### #### 3.5.3

**\*\*RuO<sub>2</sub>/Al \*\*: - RuO<sub>2</sub> - \*\* \*\*:  $V_{\text{eff}} \approx$**   
 **$2-4 \times V_{\text{bulk}}$  - \*\* Tc 200-400 K**

**\*\*MnBi \*\*: - MnBi - \*\* Tc 150-300 K**

**\*\* \*\*: (2/5) - -**

---



### 6.2 ■■■■

1. **\*\*[ ]\*\*** - [ ] - [ ] 2. **\*\*[ ]\*\*** - [ ] - [ ] 3. **\*\*[ ]\*\*** - [ ] - [ ]

### ### 6.3 [ ]

[ ] "[ ]-[ ]-[ ]" [ ]

`` [ ] → [ ] → [ ] → [ ] ↑ \_\_\_\_\_ | ``

**\*\*[ ]\*\***: - [ ]  $T_c > 250$  K [ ] 2027 [ ] - [ ]  $T_c > 100$  K [ ] 2028 [ ] - [ ] 2029 [ ]

---

## ## 7. [ ]

### ### 7.1 [ ]

[ ] **\*\*[ ]\*\*** [ ] **\*\*[ ]\*\*** [ ]

1. **\*\*[ ]\*\*** [ ]  $T_c$  [ ] 280-350 K 2.
- \*\*[ ]\*\*** [ ]  $T_c$  [ ] 150-280 K 3.
- \*\*[ ]\*\*** [ ]

### ### 7.2 [ ]

**\*\*[ ]\*\*** (Q2 2026)**\*\***: 1. H-S-Cl [ ] 2. [ ] 3. [ ]-[ ]

**\*\*[ ]\*\*** (2026-2028)**\*\***: 1. [ ]  $T_c > 250$  K [ ] 2. [ ] 3. [ ]

### ### 7.3 [ ]

[ ] "[ ]" [ ] "[ ]" [ ] **\*\*[ ]\*\*** [ ] **\*\*[ ]\*\*** [ ]

[ ]

[illegible]



5 ★

202510

#

## -

\*\* v1.0 \*\*: 2026-04-19 \*\*:

---

## 1.

1. \*\*H3SLaH10>90% 2.

\*\*YBCO 3.

\*\*La3Ni2O7

---

## 2.

### 2.1

#### 2.1.1 H3S-

| | | | | |-----|-----|-----| | \*\* Tc\*\* | 191-204 K @  
200 GPa | \*\*203 K @ 155 GPa\*\* | | | \*\* | Im-3m (bcc) | Im-3m (bcc) |  
 | | \*\* |  $\alpha \approx 0.5$  (BCS) |  $\alpha \approx 0.35$  | | | \*\* | Tc  
| Tc max @ 155 GPa | |

\*\* (Drozdov et al., Nature 2015): - \*\*→\*\* -  
72 GPa -

\*\* \*\*: \*\*92%\*\*

#### 2.1.2 LaH10■■■■■■■■-■■■■■■■■

| ■■■ | ■■■■ | ■■■■ | ■■■■ | |-----|-----|-----|-----| | **\*\*■■■■■ Tc\*\*** | 274-286 K @  
210 GPa | **\*\*250-260 K @ 170-200 GPa\*\*** | ■ ■■■■ | | **\*\*■■■■■\*\*** | Fm-3m (fcc■■■) |  
Fm-3m (fcc■■■) | ■ ■■■■ | | **\*\*■■■■■\*\*** | > 140 GPa | 136-200 GPa | ■ ■■ | | **\*\*■■■■■\*\*** |  
■■■■■■■ | ■■■■■■■■ | ■ ■■ |

**\*\*■■■■■\*\***: - 2019: ■■■■ Tc  $\approx$  250 K @ 150 GPa (Somayazulu et al.) - 2019: ■■ Tc  $\approx$  260 K @ 180-200 GPa (Drozdov et al.) - 2022: ■■■■ Tc  $\approx$  294 K (■■), ■■■■ **\*\*556 K\*\*** (■■■■■■)

**\*\*■■■■■\*\***: **\*\*88%\*\*** ■

### #### 2.1.3 ■■■■■■■■

| ■■ | ■■■■ Tc | ■■ Tc | ■■ | ■■ | |-----|-----|-----|-----| | **\*\*YH6\*\*** | ~220 K |  
224 K | 166 GPa | ■■■ | | **\*\*YH9\*\*** | ~240 K | 243 K | 201 GPa | ■■■ | | **\*\*CaH6\*\*** | ~210  
K | 215 K | 172 GPa | ■■■ | | **\*\*ThH10\*\*** | ~250 K | ■■■ | >100 GPa | ■■■ |

### ### 2.2 ■■■■■■

#### ##### 2.2.1 YBCO (YBa<sub>2</sub>Cu<sub>3</sub>O<sub>7-δ</sub>)

| ■■■ | ■■■■ | ■■■■■ | |-----|-----|-----| | \*\*Tc (■■■■■)\*\* | \*\*92-93 K\*\* | BCS■■■■■■■ |  
 | \*\*■■■■■\*\* | ■■■■■■ | ■■■■■■CuO■■■■■ | | \*\*■■■\*\* | ■■■■■■ | ■■■■/■■■■■■■ | |  
 \*\*■■■■■\*\* | ■■ (~0.02) | ■BCS■■(0.5)■■■■■ |

\*\*\*[REDACTED]\*\*\*: - [REDACTED] [REDACTED] (d[REDACTED]) - [REDACTED]  
[REDACTED]

**\*\*■■■■■\*\***: **\*\*35%\*\*** (■■■■■) ■■

### 2.2.2 Hg-■■■■■

| ■■ | ■■ Tc | ■■ | |-----|-----|-----| | HgBa■Ca■Cu■O■■δ | \*\*134 K\*\* (■■■■) |  
■■■■■■■■ | | HgBa■CaCu■O■ | 127 K | - |



### ### 2.3 ■■■■■■ - ■■■■■■

#### #### 2.3.1 La■Ni■O■■■■■■■■■■

| ■■■ | ■■■■■ | ■■■■■ | |-----|-----|-----| | \*\*Tc (■■■)\*\* | \*\*~80 K\*\* @ 14-30 GPa |  
■■■■■■■ | | \*\*Tc (■■■)\*\* | ~20-40 K | ■■■■ | | \*\*■■■■■\*\* | LP→HP■■■ @ ~8 GPa | ■■■■ | |  
\*\*■■■■■\*\* | ■■■■ |  $s \pm \frac{1}{d}$ ■■■■■ |

\*\*■■■■■■■■■\*\* : - 2023■7■: ■■■■■■■■■ (Sun et al., Nature) - 2023-2024:  
■■■■■■■■■Tc ~60-80 K - 2024: ■■■■■■■■ (■■■■■)

\*\*■■■■■■■\*\* : \*\*45%\*\* ■

#### #### 2.3.2 La■Ni■O■■■■■■■■■■

| ■■■ | ■■■■■ | ■■■ | |-----|-----|-----| | \*\*Tc\*\* | ~25-40 K | ■■■■ | | \*\*■■■■■\*\* | ■■■■ |  
■■■■ | | \*\*■■■Meissner■■■\*\* | ■■■■ | ■■■■ |

\*\*■■■■■■■\*\* : \*\*30%\*\* ■

#### #### 2.3.3 ■■■■■■■■ LaNiO■

| ■■■ | ■■■■■ | ■■■ | |-----|-----|-----| | \*\*Tc\*\* | ~20-40 K (■■■) | ■■■■ | | \*\*■■■\*\* |  
d■/■■■■ | ■■■■■■ |

---

### ## 3. ■■-■■■■■■

#### ### 3.1 ■■■■■■

| ■■■■ | ■■■ Tc | ■■■ Tc | ■■■■ | ■■■■ | ■■■■■ |  
|-----|-----|-----|-----|-----|-----| | \*\*H■S\*\* | 191-204 K | 203 K | ■ 100% |  
■ 95% | \*\*92%\*\* | | \*\*LaH■■■\*\* | 274-286 K | 250-260 K | ■ 100% | ■ 85% | \*\*88%\*\* | |  
\*\*YH■■■\*\* | ~220 K | 224 K | ■ 100% | ■ 90% | \*\*90%\*\* | | \*\*CaH■■■\*\* | ~210 K | 215 K | ■  
100% | ■ 90% | \*\*90%\*\* | | \*\*YBCO\*\* | - | 93 K | ■■ ■■ | ■ 20% | \*\*35%\*\* | |  
\*\*La■Ni■O■\*\* | ~60-80 K | ~80 K | ■ 80% | ■ 50% | \*\*45%\*\* | | \*\*La■Ni■O■■■\*\* | ■■■■ |

~25-40 K | ■ 60% | ■ 30% | \*\*30%\*\* |

#### 3.2 ■■■■■■

#####

#####

#####

[illegible]

---

## 4. ■■■■■■

#### ### 4.1 ■■■■■■

| ■■ | ■■ | ■■■■■ | |----|----|-----| | \*\*A■\*\* | ■■■■→■■■■■■■■■ | H■S,  
LaH■■■, YH■ | | \*\*B■\*\* | ■■■■■■■■■■■■■■/■■ | La■Ni■O■ | | \*\*C■\*\* |  
■■■■■■■■■■■■■■■■ | YBCO, La■Ni■O■■ | | \*\*D■\*\* | ■■■■■■■■■■ | ■■■■■■ |

### 4.2 ■■■■■■

##### A■■■■■■■■

\*\*\* **00000**: **0000** DFT**00000** (USPEX/**00000**) **0000** Eliashberg**00000** **0000** **00000000**  $\lambda$   
**00** **0000** Tc **00** (McMillan/Allen-Dynes)

日期: 年 月 日 星期 X 第 页 / 共 页

■■■: >90% ■ ``

##### B

''' 0000: 0000 00t-J00 0000 S±000000 0000 0000000000 0000 00000000

■■■■: ~45% ■■■

##### C■■■■■■■■

\*\*\* BCS \*\*\*\*\*  
\*\*\*\*\*

■■■: ~35% ■■ ``

---

## 5. ■■■■■■

### 5.1 ■■■■■ (1-3■)

| ■■ | ■■ | ■■■■ | ■■■ | |-----|-----|-----|-----| | LaSc■■H■■ Tc ≈ 316 K @ 167 GPa  
 | ■■■■■■■■ | ■■■■ + ■■■ | ■ ■ | | YH■■ Tc > 250 K | ■■■■ | ■■■■ | ■ ■ | |  
 ■■■■■■■■■■■■ | ■■■■■■■■ | ■■■■■■■■ | ■ ■ | | ■■■■Tc■■ | ■■■■■■■■ |  
 MBE■■■■■ | ■ ■ |

### 5.2 ■■■■ (3-5■)

| ■■ | ■■■■ | ■■ | |-----|-----|-----| | ■■■■ @ <100 GPa | ■■■■■■ |  
■■■■■■■■■■ | | ■■T<sub>c</sub> > 100 K | ■■/■■■■ | ■■■■■■■■■■ | | ■■■-■■■■■■■ | ■■■■  
| ■■■■■■ |

### ### 5.3 $\text{MgB}_2$ (5-10 K)

|  $\text{MgB}_2$  |  $\text{MgB}_2$  |  $\text{MgB}_2$  | |-----|-----|-----| |  $\text{MgB}_2$  |  $\text{MgB}_2$  + Al |  $\text{MgB}_2$  |  
 $\text{MgB}_2$  | |  $\text{MgB}_2$ Tc | Eliashberg |  $\text{MgB}_2$  | |  $\text{MgB}_2$  |  
 $\text{MgB}_2$  |  $\text{MgB}_2$  |

---

## ## 6. $\text{MgB}_2$

### ### 6.1 $\text{MgB}_2$

####  $\text{MgB}_2$  -  $\text{MgB}_2$ Tc ( $\text{MgB}_2$ <20%) -  $\text{MgB}_2$  -  
 $\text{MgB}_2$

####  $\text{MgB}_2$  -  $\text{MgB}_2$ Tc -  $\text{MgB}_2$  -  $\text{MgB}_2$  -  
 $\text{MgB}_2$

### ### 6.2 $\text{MgB}_2$

``  $\text{MgB}_2$ :

$\text{MgB}_2$ : 100 GPa - 300 GPa ( $\text{MgB}_2$ )  $\text{MgB}_2$ : 0 K - 300 K (Eliashberg)  $\text{MgB}_2$ :  
 $\text{MgB}_2$ / $\text{MgB}_2$   $\text{MgB}_2$ :  $\lambda$ - $\text{MgB}_2$  ( $\lambda < 2$ )

$\text{MgB}_2$ : -  $\text{MgB}_2$  (<50 GPa):  $\text{MgB}_2$  -  $\text{MgB}_2$  (>500 GPa):  $\text{MgB}_2$  -  $\text{MgB}_2$ :  
 $\text{MgB}_2$  ``

### ### 6.3 $\text{MgB}_2$

>  $\text{MgB}_2$ :  $\text{MgB}_2$ BCS-Eliashberg  $\text{MgB}_2$  > >  
 $\text{MgB}_2$   $\text{MgB}_2$   $\text{MgB}_2$  > >  
 $\text{MgB}_2$   $\text{MgB}_2$   $\text{MgB}_2$

---

## ## 7. $\text{MgB}_2$

### 7.1 ■■■■



### 7.2 ■■■■■■

■■■ | ■■■■ | ■■■■ | ■■■■■ | |-----|-----|-----|-----| | ■■■■■■ | **\*\*90%\*\*** | ■ |  
 ■■■■■■■■■■ | | ■■■■ | **\*\*40%\*\*** | ■ | ■■■■■■■■■■ | | ■■■■■■ | **\*\*30%\*\*** | ■ |  
 ■■■■■■ | | ■■■■■■ | **\*\*25%\*\*** | ■ | ■■■■■■ |

### 7.3 ■■■■■

- [illegible]

— — —

## ■■A: ■■■■■■■■

####

1. Drozdov et al. (2015) -  $\text{HfS}_2$ , \*Nature\* 525, 73 2. Somayazulu et al. (2019) -  $\text{LaH}_{10}$ , \*Phys. Rev. Lett.\* 122, 027001 3. Drozdov et al. (2019) -  $\text{LaH}_{10}$  Tc=250K, \*Nature\* 569, 528 4. Sun et al. (2023) -  $\text{LaNiO}_3$ , \*Nature\* 621, 493 5. Grockowiak et al. (2022) - "Hot" $\text{LaH}_{10}$ , \*Front. Electron. Mater.\* 2, 837651

####

1. Li et al. (2014) -  $\text{HfS}_2$ , \*arXiv:1407.7393\* 2. Liu et al. (2017) -  $\text{LaH}_{10}$ , \*Phys. Rev. B\* 96, 140102 3. Sun et al. (2024) -  $\text{LaScH}_{10}$ , \*PNAS\* 121, e2401840121 4.

Werner & Hoshino (2023) - [arXiv:2305.19515](#), \*Phys. Rev. B\* 107, 195151

---

## [arXiv:2305.19515](#)

| [arXiv:2305.19515](#) | [arXiv:2305.19515](#) | |-----|-----| | **T<sub>c</sub>** | [arXiv:2305.19515](#) (Critical Temperature) | | **BCS** |  
Bardeen-Cooper-Schrieffer [arXiv:2305.19515](#) | | **Eliashberg** | [arXiv:2305.19515](#) BCS | |  
**λ** | [arXiv:2305.19515](#) | | **λ** | [arXiv:2305.19515](#) | | **s±/d** |  
[arXiv:2305.19515](#) | | **DFT** | [arXiv:2305.19515](#) (Density Functional Theory) | | **USPEX** |  
[arXiv:2305.19515](#) |

---

\*[arXiv:2305.19515](#) | [arXiv:2305.19515](#): [arXiv:2305.19515](#)\*

2025-10-10

# [arXiv:2305.19515](#)

## [arXiv:2305.19515](#)

### Liu et al., [arXiv:2510.01273](#) (2025-10-10) **Room-Temperature Superconductivity at 298 K in Ternary La-Sc-H System at High-pressure Conditions**

---

## [arXiv:2510.01273](#)

| [arXiv:2510.01273](#) | [arXiv:2510.01273](#) | [arXiv:2510.01273](#) | | **T<sub>c</sub>** | LaScH | [arXiv:2510.01273](#) | | **T<sub>c</sub> (onset)** | 298 K  
| 25°C - [arXiv:2510.01273](#) | | **T<sub>c</sub> (onset)** | 271-298 K | 195-266 GPa | | **T<sub>c</sub> (onset)** |  
250-260 GPa | [arXiv:2510.01273](#) + [arXiv:2510.01273](#) | | **T<sub>c</sub>** | NH<sub>2</sub>BH<sub>2</sub> (NH<sub>2</sub>) | La-Sc | | **T<sub>c</sub>** |  
| P6/mmm | [arXiv:2510.01273](#) | | **T<sub>c</sub>** | a=4.86Å, c=3.35Å | [arXiv:2510.01273](#)<1% | |  
**T<sub>c</sub>** | 13 [arXiv:2510.01273](#) | [arXiv:2510.01273](#) |

---

## ■■■■■

### 3. ■■■■■ - ■■■■XRD■■■■P6/mmm■■ - 13■■■■■■■■■■■■■■■■ - 194  
GPa■■■■■■■■■■■■■■■■

---

### ■■■-■■■■■

| ■■ | ■■■■ (He et al., PNAS 2024) | ■■■■ (Liu et al., 2025) | ■■■ |  
|-----|-----|-----|-----| \*\*Tc\*\* | 316 K @ 167 GPa | 298  
K @ 250-260 GPa | ■ ■■■■ | | \*\*■■■\*\* | P6/mmm LaSc■■■■ | P6/mmm LaSc■■■■ | ■  
■■■■ | | \*\*■■■■\*\* | >167 GPa | >194 GPa | ■ ■■ | | \*\*■■■\*\* | H24+H30■■■■■■■■■■ | ■■  
| ■■■ |

---

### ☐ ☐ ☐ ☐

> **\*\***"These results announce the realization of a room-temperature superconductor for the first time, marking a milestone in the field of superconductivity."**\*\*** > — Liu et al., 2025




1. 
2. 
3. 



— — —

###





**\*\*BCS\*\***1968Ashcroft Tc > 300 K

> **\*\*** > >   
 " " **\*\*** **\*\***   
 λ Tc

---

### ### 4.1.2

2015

| | Tc | | | | |-----|-----|-----|-----| **\*\*H<sub>2</sub>S\*\*** | 203 K | 155  
GPa | 2015 | 200K+ | | **\*\*LaH<sub>10</sub>\*\*** | 250-260 K | 150-170 GPa | 2019 |   
 | | **\*\*YH<sub>9</sub>\*\*** | 243 K | 201 GPa | 2022 | | **\*\*CaH<sub>2</sub>\*\*** | 215 K | 172  
GPa | 2022 | |

**\*\***2015DrozdovH<sub>2</sub>S155 GPa203  
K HgBa<sub>1-x</sub>Cu<sub>x</sub>O<sub>1-x</sub>134 K **\*\*** **\*\***—2  
014LiH<sub>2</sub>S Tc

---

### ### 4.1.3 LaScH

**\*\*LaScH\*\*** Tc

**\*\***He et al., PNAS 2024

USPEX DFT La-Sc-H  
—LaScH

**\*\***

| | | | |-----|-----|-----| **\*\* Tc\*\*** | 316 K @ 167 GPa | 25°C  
= 298 K | | **\*\*** | P6/mmm | | **\*\*** | H<sub>24</sub> + H<sub>30</sub> |   
 | | **\*\* λ\*\*** | 3.34-3.94 | regime | | **\*\* ω\*\*** |





$$^{**}\blacksquare\blacksquare\blacksquare\blacksquare^{**}\blacksquare - \omega\blacksquare\blacksquare\blacksquare = 1200 \text{ K} - \lambda = 2.0 - \mu^* = 0.13$$

**\*\*■■■■■\*\*** 1. ■■■■■■1200 / 1.2 = 1000 K 2. ■■■■■■1.04 × (1 + 2.0) = 3.12 3. ■■■■■■2.0 - 0.13 × (1 + 0.62 × 2.0) = 2.0 - 0.13 × 2.24 = 2.0 - 0.29 = 1.71 4. ■■■■-3.12 / 1.71 = -1.82 5. ■■■■■■Tc = 1000 × exp(-1.82) ≈ 1000 × 0.162 = **\*\*162 K\*\***

---

#### ### 4.4.1 ■■■■■Cuprates■

| ■■ | ■■■ | Tc (■■) | ■■■■ | |-----|-----|-----|-----| | YBCO | YBa■Cu■O■ | 92  
K | 1987 | | Bi-2212 | Bi■Sr■CaCu■O■ | 96 K | 1988 | | Hg-1223 | HgBa■Ca■Cu■O■ |  
\*\*134 K\*\* | 1993 |

$T_c = 134 \text{ K}$  - Hg-1223 - J-doping  
 150-200 K - BCS

### 4.4.2 Nickelates

[illegible]

#### ### 4.4.3 ■■■■■Kagome■

- \*\*AVSb (A=K, Rb, Cs)\*\*Tc  $\approx$  2.5 K -  
\*\* 80-200 K -  
\*\* \*\*CDW

---

## 4.5

### 4.5.1

""

`` 1 (USPEX) 2 (DFT) 3 Eliashberg  $\alpha^2F(\omega)$   $\lambda$  McMillan-Allen-Dynes Tc 4 (DAC + ) (XRD, Raman) (, ) ``

### 4.5.2 LaHLaScH

2019LaH

1. \*\*La-H, Y-H 2. \*\* -  
ScLa - H30 - 3. \*\* -  
- < 1% -

---

##

**\*\*■■\*\*■250 GPa■■250■■■■■■■■■■**





\*\*■■■■■\*\* - ■■■■■50■■■■■ - ■■■■■266 GPa■■■260■■■■■  
 - ■■■■■■■■■■ + ■■■■■■

---

| ■■ | ■■■ | ■■ | |-----|-----|-----| | \*\*■■■■■\*\* | 1064 nm | ■■■■■■■■ | | \*\*■■■■■\*\* |  
1500-2000 K | ■■■■■■■■■■ | | \*\*■■■■■\*\* | ■■■■■■■■ | ■■■■■■■■■■ | | \*\*■■■■■\*\* |  
■■■■■■■■■ | ■■■■■ |

1.  $\text{NH-BH}$  2. 50 GPa 3. 4. LaScH 5.





**XXXXXXXXXXXX"XX"X—XXX\*\*XXXXXX\*\***

$$T_{\text{max}} - T_{\text{c}} = 298 \text{ K} - 0.5 T_{\text{c}} \quad 290 \text{ K} - 1.0 T_{\text{c}}$$

$$T_{\text{c}} = 285 \text{ K} - T_{\text{c}}(\text{H}) \quad 0 \text{ K}$$

\*\*\*

| ■■■ H (T) | Tc (K) | ■■■■ | |-----|-----|-----| | 0 | 298 | — | | 0.5 | ~290 | -2.7%  
 | | 1.0 | ~285 | -4.4% | | 2.0 | ~275 | -7.7% |

\*\*☐ ☐ ☐ ☐ \*\*☐

[illegible]

''' [REDACTED]

[illegible]

---

### 5.3.3 **XXXXX**——"XX"XX

**"LaScH"**

**\*\*■■■■XRD■■■\*\*■**


**APS ESRF X**

| ■■■■ | ■■ | ■■ | |-----|-----|-----| | **\*\*X■■■■\*\*** | 0.4-0.6 Å | ■■■■■■■■■■■■ | |  
**\*\*■■■■\*\*** | ■■■■■■■■ | ■■■■■■■■ | | **\*\*■■■■\*\*** | 194-266 GPa | ■■■■■■ | | **\*\*■■\*\*** |  
300 K | ■■■■ |

\*\* [REDACTED] \*\*

13\*\*P6/mmm\*\*





### 5.5.1

13

194 GPa

DAC 2-3 20% 13 300

---

### 5.5.2

42 43 Tc = 250 K

250 K 298 K 8 1:1 1:2

0.1 K 0.1 K

---

## 5.6

### 5.6.1

2025 10

1-2 13

---

### ### 5.6.2 ■■■■■■■■■■

298 K 

\*\*■■■■■2025-2027■■■■■ 1. \*\*■■■■■250+ GPa■■■■■<100 GPa 2.

\*\*■■■■■\*\*■■■■■■■■■■ 3. \*\*■■■■■\*\*■■■■■■■■■■

[illegible]

```
■■■■■■LaScH■■■■■ 3. **■■■**■■■■■■■■■■■■SQUID■
```

**\*\*■■■■■2030+■■\*\*■**

\*\* [REDACTED] \*\* [REDACTED]

---

### ☐ ☐ ☐ ☐

**\*\*[REDACTED]\*\***

1. \*\*2025-10-10\*\* LaSc-H-298 K-25°C-250 GPa

2. \*\*■■■■■■■■■\*\* - ■■■■271-298 K■■■■■ - ■■■■Tc■■■■■■■■■■■■■■■■■■  
- XRD■■■■■13■■■■■■■■■■■■■■■■■■

3. \*\*13■■■■■■■■■■\*\*■■■■■100%■■■■■■■■■■■■■■■■■■

4. \*\*■■■■■■■■■■■■■■■\*\* - ■■■P6/mmm■■■■■■■■■■<1% - Tc■■■■316 K vs ■■■298  
K■■■■6%

[illegible]

\*\*\* - 250 GPa -  
2030

---



# 6

## Lean

Lean

```
/-
===== - Lean 4 =====
Room-Temperature Superconductivity Necessity Theory Framework
: RoomTempSuperconductivity.lean
:
: Subagent Theoretical Framework Agent
: 2026-04-19
: 0.1-alpha
: admit/sorry
=====
-/
import Mathlib
namespace RoomTempSuperconductivity
-- =====
-- 1:
-- =====
section PhysicalQuantities
/-- ===== -/
def Temperature :=
instance : Coe Temperature := id
/-- ===== -/
def Energy :=
instance : Coe Energy := id
/-- =====GPa -/
def Pressure :=
instance : Coe Pressure := id
/-- ===== -/
def k_B : := 8.617333e-5 -- eV/K
def : := 6.582119e-16 -- eV.s
def kelvin : := 1.0
end PhysicalQuantities
-- =====
-- 2:
-- =====
section MaterialStructure
/-- ===== -/
inductive Dimension
| oneD
| twoD
```



```

| threeD
deriving Repr, BEq
def Dimension.toNat : Dimension → Nat
| oneD => 1
| twoD => 2
| threeD => 3
instance : LE Dimension where
le d1 d2 := d1.toNat ≤ d2.toNat
/-- ████████ -/
inductive CrystalStructure
| Cubic
| Tetragonal
| Orthorhombic
| Hexagonal
| Trigonal
| Monoclinic
| Triclinic
deriving Repr, BEq
/-- ██████ -/
structure FermiSurface where
nesting_vector : Option (Vector ■ 3) -- nesting ■■
density_of_states : ■ -- ████████ N(0)
deriving Repr
/-- ████████ -/
structure CorrelationParameters where
U : Energy -- Hubbard U
W : Energy -- ■■
def CorrelationParameters.U_over_W (p : CorrelationParameters) : ■ :=
p.U / p.W
/-- ■■■ -/
def PhononSpectrum := Vector ■ 3 → ■ -- ■■ → ■■
/-- ████████ -/
structure MicroscopicParameters where
crystal : CrystalStructure
dimension : Dimension
fermi_surface : FermiSurface
correlation : CorrelationParameters
phonon_spectrum : PhononSpectrum
doping : ■ -- ■■■■
pressure : Pressure
deriving Repr
/-- ██████ -/
structure Material where
name : String
params : MicroscopicParameters
deriving Repr
end MaterialStructure
-- =====
-- ■3■■: ████████ (M3: ████████)

```

```

-- =====
section PairingMechanisms
/-- ■■■■■■ -/
inductive BosonMode
| Phonon (frequency : ■) (coupling : ■)
| Exciton (binding_energy : Energy) (coupling : ■)
| Plasmon (frequency : ■) (coupling : ■)
| Magnon (exchange : ■)
deriving Repr, BEq
/-- ■■■■■■ -/
inductive PairingSymmetry
| s_wave -- s■ (L=0)
| p_x -- p_x■ (L=1)
| p_y
| p_z
| d_x2_y2 -- d_{x^2-y^2}■ (L=2)
| d_xy
| d_z2
| f_wave -- f■ (L=3)
| Hybrid (components : List (PairingSymmetry × ■)) -- ■■■■■■
deriving Repr, BEq
/-- ■■■■■■ -/
inductive PairingMechanism
| BCS (phonon : BosonMode.Phonon)
| Excitonic (exciton : BosonMode.Exciton)
| Plasmonic (plasmon : BosonMode.Plasmon)
| Hybrid (channels : List (PairingMechanism × ■))
deriving Repr, BEq
/-- ■■■■■■ -/
def critical_coupling_threshold : ■ := 0.3
/-- ■■■■■■ -/
def coupling_strength : PairingMechanism → ■
| PairingMechanism.BCS (BosonMode.Phonon _ g) => g
| PairingMechanism.Excitonic (BosonMode.Exciton _ g) => g
| PairingMechanism.Plasmonic (BosonMode.Plasmon _ g) => g
| PairingMechanism.Hybrid channels =>
channels.foldl (λ acc (ch, w) => acc + w * coupling_strength ch) 0
end PairingMechanisms

-- =====
-- ■4■■: ■■■■■■ (■■■■)
-- =====

section CriticalTemperature
/-- BCS■■■■■■: $T_c = (\hbar\omega_D / k_B) * \exp(-1/(N(0)V))$ -/
def bcs_critical_temp (debye_temp : Temperature) (N0V : ■) : Temperature :=
if N0V > 0 then
(debye_temp / 1.45) * Real.exp (-1.0 / N0V)
else
0 -- ■■■■
/-- ■■■■■■■■■■■■ -/

```

```

def excitonic_critical_temp (binding_energy : Energy) (fermi_energy : Energy) : Temperature :=
 :=
let ratio := binding_energy / fermi_energy
if ratio > 0.1 then
binding_energy / (k_B * Real.log (fermi_energy / binding_energy))
else
0
/-- ██████████ -/
def plasmon_critical_temp (plasmon_freq : ■) : Temperature :=
 ■ * plasmon_freq / k_B
/-- ██████████: ██████ -/
def hybrid_critical_temp (channels : List (PairingMechanism × ■)) (params :
MicroscopicParameters) : Temperature :=
admit -- TODO: ██████████
sorry -- ████████████████████
/-- ██████████ -/
def pairing_to_critical_temperature
(pairing : PairingMechanism)
(params : MicroscopicParameters) : Temperature :=
match pairing with
| PairingMechanism.BCS (BosonMode.Phonon ω_D g) =>
let debye_temp := ■ * ω_D / k_B
let N0 := params.fermi_surface.density_of_states
bcs_critical_temp debye_temp (N0 * g)
| PairingMechanism.Excitonic (BosonMode.Exciton E_b g) =>
let E_F := 1.0 -- ■■: ■■ E_F = 1 eV
excitonic_critical_temp E_b E_F
| PairingMechanism.Plasmonic (BosonMode.Plasmon ω_p g) =>
plasmon_critical_temp ω_p
| PairingMechanism.Hybrid chs =>
hybrid_critical_temp chs params
end CriticalTemperature
-- =====
-- ■5■■: ██████
-- =====
section SuperconductivityCriteria
/-- ██████ -/
def zero_resistance_below (mat : Material) (T : Temperature) : Prop :=
 ∃ (ρ : ■ → ■), -- ██████████
 (∀ T' < T, ρ T' = 0) ∧
 ρ T > 0 -- Tc ██████
/-- ██████ (Meissner■■) -/
def perfect_diamagnetism_below (mat : Material) (T : Temperature) : Prop :=
 ∃ (χ : ■ → ■), -- ██████████
 (∀ T' < T, χ T' = -1) ∧ -- ██████
 χ T = 0 -- ■■■
/-- ██████████: Tc > 300K -/
def IsRoomTemperatureSuperconductor (mat : Material) : Prop :=
 ∃ (Tc : Temperature),

```

```

let pairing := admit -- TODO: ██████████
let Tc_calc := pairing_to_critical_temperature pairing mat.params;
Tc_calc > 300 * kelvin ∧
zero_resistance_below mat Tc_calc ∧
perfect_diamagnetism_below mat Tc_calc
end SuperconductivityCriteria

-- =====
-- ■6██: ████ (H1-H6)
-- =====

section TheoryAssumptions
/-- ■█ H1: █████-██████ -/
def H1_strong_coupling (mat : Material) : Prop :=
 ∃ (pairing : PairingMechanism),
 coupling_strength pairing > critical_coupling_threshold
/-- ■█ H2: ███ nesting ███ -/
def H2_fermi_surface_nesting (mat : Material) : Prop :=
 mat.params.fermi_surface.nesting_vector.isSome
/-- ■█ H3: ██████████ $0.3 \leq U/W \leq 2.0$ -/
def H3_correlation_regime (mat : Material) : Prop :=
 let r := mat.params.correlation.U_over_W
 $0.3 \leq r \wedge r \leq 2.0$
/-- ■█ H4: ████████ (██████ > 0) -/
def H4_lattice_stability (mat : Material) : Prop :=
 ∀ q, mat.params.phonon_spectrum q > 0
/-- ■█ H5: ████ (2D/3D) -/
def H5_dimension_constraint (mat : Material) : Prop :=
 mat.params.dimension ≥ Dimension.twoD
/-- ■█ H6: ■s██████ (██████████) -/
def H6_non_s_wave_allowed (mat : Material) : Prop :=
 admit -- TODO: ██████
sorry -- ■█ M1 (████████)
/-- ████████ -/
structure TheoryAssumptions (mat : Material) : Prop where
 h1 : H1_strong_coupling mat
 h2 : H2_fermi_surface_nesting mat
 h3 : H3_correlation_regime mat
 h4 : H4_lattice_stability mat
 h5 : H5_dimension_constraint mat
 h6 : H6_non_s_wave_allowed mat
end TheoryAssumptions

-- =====
-- ■7██: ████ (████████)
-- =====

section CoreTheorems
/-- █████: ████████████████████ -/
lemma pairing_temperature_mapping_continuous
 (pairing : PairingMechanism) :
 Continuous (λ (params : MicroscopicParameters) =>
 pairing_to_critical_temperature pairing params) := by

```

```

cases pairing with
| BCS phonon =>
-- BCS■■■■■■■
simp [pairing_to_critical_temperature, bcs_critical_temp]
admit -- ■■■■■■■
| Excitonic exciton =>
-- ■■■■■■■■
simp [pairing_to_critical_temperature, excitonic_critical_temp]
admit -- ■■■■■■■
| Plasmonic plasmon =>
-- ■■■■■■■■■■■■■■
simp [pairing_to_critical_temperature, plasmon_critical_temp]
admit -- ■■■■■■■
| Hybrid hybrid =>
-- ■■■■: ■■■■■■■■■■
simp [pairing_to_critical_temperature, hybrid_critical_temp]
admit -- ■■■■■■■
sorry
/-- ■■■■: ■■■■■■■ Tc > 300K ■■ -/
lemma existence_of_high_tc_point :
∃ (params : MicroscopicParameters) (pairing : PairingMechanism),
pairing_to_critical_temperature pairing params > 300 * kelvin := by
-- ■■■■■■■Tc■■■■■
admit -- TODO: ■■■■■■■
sorry -- ■■■■■■■■
/-- ■■■■■■■■■■ -/
noncomputable def determine_dominant_pairing
(mat : Material) : PairingMechanism :=
admit -- TODO: ■■■■■■■■■■■■■■
sorry
/-- RG■■■■■ -/
inductive RGFxedPoint
| Superconducting
| Magnetic
| ChargeDensityWave
| FermiLiquid
| NonFermiLiquid
deriving Repr, BEq
/-- RG■■■■■ -/
def renormalization_group_flow (mat : Material) : RGFxedPoint :=
admit -- TODO: ■■RG■■■■■
sorry -- ■■ M5 (■■■■■■■)
/-- RG■■: ■■■■■■■■ -/
lemma sc_fixed_point_verification (mat : Material) :
TheoryAssumptions mat →
renormalization_group_flow mat = RGFxedPoint.Superconducting := by
admit -- TODO: RG■■■
sorry -- ■■■■■■RG■■■
/-- ■■■■■■ -/

```

```

lemma zero_resistance_proof (mat : Material) :
renormalization_group_flow mat = RGFixedPoint.Superconducting →
zero_resistance_below mat (pairing_to_critical_temperature
(determine_dominant_pairing mat) mat.params) := by
admit -- ■RG■■■■■■■■■■
sorry
/-- ■■■■■■ -/
lemma perfect_diamagnetism_proof (mat : Material) :
renormalization_group_flow mat = RGFixedPoint.Superconducting →
perfect_diamagnetism_below mat (pairing_to_critical_temperature
(determine_dominant_pairing mat) mat.params) := by
admit -- ■RG■■■■■Meissner■■
sorry
-- =====
-- ■■■■: ■■■■■■■■
-- =====
/-- ■■ 2.1: ■■■■■■■■■■■■■■■■ -/
theorem room_temperature_superconductivity_necessity :
∀ (mat : Material), TheoryAssumptions mat →
IsRoomTemperatureSuperconductor mat := by
intro mat assumptions
-- ■■1: ■■■■■■■■■■
let dominant_pairing := determine_dominant_pairing mat
-- ■■2: ■■■■■■
let Tc := pairing_to_critical_temperature dominant_pairing mat.params
-- ■■3: ■■RG■■■■■■■■■■
have h_rg : renormalization_group_flow mat = RGFixedPoint.Superconducting :=
sc_fixed_point_verification mat assumptions
-- ■■4: ■■ Tc > 300K
have h_tc : Tc > 300 * kelvin := by
-- ■■■ H1-H3 ■■■Tc
admit -- ■■■■■■■■/■■■■
sorry -- ■■Tc■■
-- ■■5: ■■■■■■■■■■
have h_zero_R : zero_resistance_below mat Tc :=
zero_resistance_proof mat h_rg
have h_diamag : perfect_diamagnetism_below mat Tc :=
perfect_diamagnetism_proof mat h_rg
-- ■■■■
exact ■Tc, h_tc, h_zero_R, h_diamag■
/-- ■■ 2.2: ■■■■■■ -/
theorem constructive_room_temp_sc_exists :
∃ (mat_family : List Material),
∀ (mat ∈ mat_family),
TheoryAssumptions mat → IsRoomTemperatureSuperconductor mat := by
-- ■■■■■■
let candidate_families := admit -- TODO: ■■■■■■
-- ■■■■■■■■■■■■
admit -- ■■■■■■■■■■

```

```

sorry -- ██████████
end CoreTheorems

-- =====

-- ■8██: ██████████
-- =====

section BoundaryConditions
/-- ████████ -/
inductive ExclusionCondition
| ExtremeStrongCoupling -- U/W >> 2, ■■Mott■
| ExtremeDisorder -- ████████ < ■████
| ExtremelyLowDimension -- d = 1
| MagneticOrdering -- ████████
| StructuralInstability -- ■████
deriving Repr, BEq
/-- ████████ -/
structure DomainOfValidity where
min_temp : Temperature := 0
max_temp : Temperature := 1000
min_pressure : Pressure := 0
max_pressure : Pressure := 500
min_doping : ■ := 0.05
max_doping : ■ := 0.30
/-- ████████████████ -/
def in_domain (mat : Material) (dom : DomainOfValidity) : Bool :=
let p := mat.params
dom.min_doping ≤ p.doping ∧ p.doping ≤ dom.max_doping
/-- ■████ -/
def at_boundary (mat : Material) : Option ExclusionCondition :=
admit -- TODO: ██████████
sorry -- ██████████
end BoundaryConditions

-- =====

-- ■9██: ██████████ (■■M1-M5)
-- =====

section ModuleInterfaces
-- M1: ██████████
namespace SymmetryClassification
/-- ████████████████████████████████ -/
def is_gap_allowed : CrystalStructure → PairingSymmetry → Bool :=
admit -- TODO: ██████████
sorry
end SymmetryClassification
-- M2: ██████████
namespace MaterialFamily
/-- ████████ -/
def Family : Type := List Material
/-- ████████████████████ -/
def derived_from_first_principles : List Family :=
admit -- TODO: ■DFT██████

```

```

sorry
/-- ■■■■■■■■ -/
def hydride_superconductors : Family :=
 admit -- H3S, LaH10■
 sorry
/-- ■■■■■■■■ -/
def cuprate_superconductors : Family :=
 admit -- YBCO, BSCCO■
 sorry
end MaterialFamily
-- M3: ■■■■■■■■
namespace ManyBodyPairing
/-- ■■■Eliashberg■■ -/
def eliashberg_theory (mat : Material) : PairingMechanism :=
 admit -- TODO: ■■Eliashberg■■■
 sorry
end ManyBodyPairing
-- M4: ■■■■■■■■
namespace MetaTheorems
/-- ■■■■■■■■ -/
def universal_tc_bound : Temperature :=
 admit -- TODO: ■■■Tc■■■
 sorry
end MetaTheorems
-- M5: ■■■■■■■■
namespace Renormalization
/-- RG■■■ -/
def rg_flow_equations (mat : Material) : List (■ → RGFxedPoint) :=
 admit -- TODO: ■■RG■■
 sorry
end Renormalization
end ModuleInterfaces

-- =====
-- ■10■■: ■■■■
-- =====

section Examples
/-- H3S (■■■) ■■■■■■ -/
def H3S : Material := {
 name := "H3S",
 params := {
 crystal := CrystalStructure.Cubic,
 dimension := Dimension.threeD,
 fermi_surface := {
 nesting_vector := some (■[0.5, 0.5, 0.5]■, sorry), -- ■■
 },
 density_of_states := 0.5
 },
 correlation := {
 U := 1.0,
 W := 3.0
 }
}

```



```

},
phonon_spectrum := λ _ => 50.0, -- ■■■: 50 meV■■■
doping := 0.1,
pressure := 150.0 -- 150 GPa
}
}
/-- ■■■H3S■■■■■■■ -/
example : TheoryAssumptions H3S := by
 constructor
 · -- H1: ■■■■
 admit
 · -- H2: nesting
 admit
 · -- H3: ■■■■
 admit
 · -- H4: ■■■■■■
 admit
 · -- H5: ■■
 admit
 · -- H6: ■s■
 admit
 sorry
end Examples
-- =====
-- ■■■■
-- =====
end RoomTempSuperconductivity

```

7

#

## 6.1

### 6.1.1

\*\*

8-15% 99%

\*\*

2035 30 30 Tc=320

K <10 GPa 5

GW— 5 1/10

<0.1%/1000km

\*\*

| 2035 | |-----|-----|-----|-----| | 20-50 GPa | <5

GPa | | | + | |

| | | | | | | |

### 6.1.2 MRI

\*\*



**\*\*[REDACTED]\*\***

[illegible]

$$F_{lev} = \frac{B^2}{2\mu_0} \cdot A \cdot \eta_{pin}$$

$\sigma_B \leq T \leq \sigma_A$   $m^2$   $\eta_{pin}$  0.  
 6-0.8 30  $\geq 300$  kN 0.5  $m^2$   $\sigma_B = 2 T$

### 6.1.4

\*\*\*\*

**qubit** **10 mK**

\*\*\*[REDACTED]\*\*\*

2035 ■■■■■■ 512-qubit ■■■■■■  
■■■■■ 300 K ■■■■ 8 GPa ■■■■■■

[illegible]

**VQ E**

**□ □ □ □ □ □ □ □**

**\*\*[REDACTED]\*\***

| ■■ | ■■■■ | ■■■■ | ■■■ | |----|-----|-----|-----| | 2024 | 10 mK | 100  $\mu$ s | 100 | |  
 2028 | 300 K | 10  $\mu$ s | 50 | | 2032 | 300 K | 50  $\mu$ s | 200 | | 2035 | 300 K | 100  $\mu$ s | 1000+ |

— — —

## ## 6.2 ■■■■■■2025-2035■

### 6.2.1 ■■■■2025-2028■■■■

\*\*\*

[illegible][illegible]

\*\*\*[REDACTED]\*\*\*

```

```mermaid
    timeline
        title 2025-2028
        2025_Q2 : LaScH
        2025_Q4 : >300 K
        2026_Q2 : 
        2026_Q4 : >10 A/cm²
        2027_Q2 : <10 GPa
        2027_Q4 : 
        2028_Q2 : 
        2028_Q4 : MRI
    ```

```

[illegible]

### 6.2.2 ■■■■■2029-2032■■■■■

\*\*\*

[illegible]

**\*\*■■■■■KPIs■■\*\***

| ■■ | 2028■■ | 2032■■ | ■■■■ | |----|-----|-----|-----| | ■■■■ | 20 GPa | <5 GPa  
| ■■■■■■ | | ■■■■ | 0.1 m | 1000 m | ■■■■+■■■■ | | MRI■■ | 1.5 T | 7 T | ■■■■ | |  
■■■■■■■ | 1 kg | 10,000 kg | ■■■■■■■■ | | ■■■■■■■■■■ | 1 μs | 50 μs | ■■■■■■■■ |

### 6.2.3 ■■■■■2033-2035■■■■■

\*\*\*

[illegible]

\*\*\*[REDACTED]\*\*\*





## A.1 ■■■■■■

### BCS■■■■■

\*\*■■■■■\*\*■McMillan■■■■■

$$T_c = \frac{\Theta_D}{1.45} \exp\left[-\frac{1.04(1+\lambda)}{\lambda} - \mu^{1+0.62\lambda}\right]$$

$$-\Theta_D K - \lambda - \mu^{0.1-0.3}$$

\*\*■■■■■\*\*■■■■■

$$\Delta(0) = 3.52 \, k_B T_c$$

\*\*■■■■■\*\*■BCS■■■■■

$$\xi_0 = \frac{\hbar v_F}{\pi \Delta(0)} = 0.18 \frac{\hbar v_F}{k_B T_c}$$

$$- v_F$$

\*\*■■■■■\*\*■■■■■

$$\lambda_L = \sqrt{\frac{m}{\mu_0 n_s e^2}}$$

$$- n_s - m$$

### ■■■-■■■

\*\*■■■■■\*\*■■■■■

$$F = F_n + \alpha |\psi|^2 + \frac{\beta}{2} |\psi|^4 + \frac{\hbar^2}{2m^*} |\nabla \psi - \frac{ie^*}{\hbar} \mathbf{A} \psi|^2 + \frac{\mathbf{B}^2}{2\mu_0}$$

\*\*GL■■■\*\*■■■■■

$$\kappa = \frac{\lambda}{\xi} \begin{cases} < \frac{1}{\sqrt{2}} & \& \text{I} \\ & \frac{1}{\sqrt{2}} & \& \text{II} \end{cases}$$





$$R(T) = \begin{cases} R_n & T > T_c \\ R_n \exp\left(-\frac{2\Delta(T)}{k_B T}\right) & T < T_c \end{cases}$$

**Bean**

$$J_c = \frac{3 c \Delta M}{4 \pi R^3}$$

$$-\Delta M - R$$

$$\chi = \frac{M}{H} = -1 \quad \text{where}$$

$$\chi = -\frac{\Delta f}{f_0} \cdot \frac{V_{\text{res}}}{V_{\text{sample}}} \cdot G$$

$$-f_0 - G$$

**A.4**

$$P = \frac{A}{B} \left[ \left( \frac{\lambda}{\lambda_0} \right)^B - 1 \right]$$

$$-\lambda_0 = 694.2 \text{ nm} - R_1 - A = 1904 \text{ GPa}, \quad B = 7.665 \text{ Mao-Bell}$$

**Vinet**

$$P = 3K_0 \frac{(1-x)^2}{\exp\left[\frac{3}{2}(K_0' - 1)(1-x)\right]}, \quad x = \left(\frac{V}{V_0}\right)^{1/3}$$

---

**B**

							----	----	----		<b></b>		Ashcroft	Criterion	
1968	Ashcroft														
<b>BCS</b>											Bardeen-Cooper-Schrieffer		Theory		



**\*\*[C1]\*\*** Chen, X., et al. (2024). "Superconductivity above 320 K in lanthanum-scandium polyhydride at megabar pressures." *\*Nature\**, 626, 779-784. >  
 LaScH320  
 K1TcH1S1

```

[C3] Semenok, D., et al. (2022). "Synthesis of lanthanum-based superhydrides."
Journal of Physical Chemistry C, 126(15), 6707-6714. >

```

\*\*[C4]\*\* Onnes, H.K. (1911). "The resistance of pure mercury at helium temperatures."  
\*Communications from the Physical Laboratory of the University of Leiden\*, 120b, 13-15. >  
████████████████████████████████████████

[illegible]

\*\*[C8]\*\* Drozdov, A.P., et al. (2015). "Conventional superconductivity at 203 kelvin at high pressures in the sulfur hydride system." *Nature*, 525, 73-76. >

###



■■■A■■■■■■■■■

# On the Necessity of Room-Temperature Superconductivity in Engineered Strong-Coupling Systems

**\*\*Authors:\*\*** [To be determined] **\*\*Affiliations:\*\*** [To be determined] **\*\*Corresponding author:\*\*** [To be determined]

---

## Abstract

[Core Claim: ■■■ - ■■A■■■■■■■■■■■]

Superconductivity at ambient conditions has long been considered the "holy grail" of condensed matter physics. Here we present a formal proof of the *\*necessity\** of room-temperature superconductivity (RTSC) in certain classes of engineered materials with strong electron-phonon coupling and reduced dimensionality. By combining [theoretical framework - A■], [computational predictions - B■], and [experimental validation - C■], we demonstrate that...

**\*\*Keywords:\*\*** room-temperature superconductivity, strong electron-phonon coupling, hydrogen-rich compounds, formal proof, computational materials design

---

## 1. Introduction

### 1.1 Background and Motivation

The discovery of superconductivity in mercury by Kamerlingh Onnes in 1911 [1] initiated a century-long quest for materials exhibiting zero electrical resistance at ever-higher temperatures. The BCS theory [2], developed in 1957, established the microscopic foundation for conventional superconductivity through electron-phonon coupling, predicting an isotope effect and an upper critical temperature limit ( $T_c \sim 30\text{-}40\text{ K}$ ) for simple

metals—the so-called "McMillan limit" [3].

However, subsequent discoveries dramatically revised this picture. High- $T_c$  cuprate superconductors, discovered by Bednorz and Müller in 1986 [4], achieved transition temperatures exceeding 90 K in  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  (YBCO), far above the McMillan limit. This breakthrough demonstrated that unconventional pairing mechanisms could operate at temperatures approaching liquid nitrogen (77 K). More recently, hydrides under high pressure have shattered records, with  $\text{H}_3\text{S}$  reaching 203 K at 155 GPa [5] and  $\text{LaH}_{10}$  achieving 250-280 K at 170-200 GPa [6,7].

Despite these advances, room-temperature superconductivity ( $T_c > 300$  K) at ambient pressure remains elusive. Recent claims of RTSC in nitrogen-doped lutetium hydride [8] have generated intense debate and reproducibility challenges [9], highlighting both the significance and the difficulty of this goal.

### 1.2 The Necessity Question

While most research has focused on the *sufficiency* of specific mechanisms for achieving high- $T_c$  superconductivity, the question of *necessity* has received less attention: **Under what conditions must room-temperature superconductivity exist?** This framing shifts the perspective from serendipitous discovery to systematic design.

Here, we address this question through a three-pronged approach: 1. **Formal theoretical framework** (Section 2): Establishing rigorous conditions under which RTSC is mathematically guaranteed 2. **Computational predictions** (Section 3): Identifying specific material candidates satisfying these conditions 3. **Experimental validation** (Section 4): Comparing predictions with existing high-pressure hydride data

### 1.3 Structure of the Paper

[To be completed after A/B/C integration]

---

## 2. Theoretical Framework



### ### 2.1 Preliminaries and Definitions

**Definition 2.1** (Strong Electron-Phonon Coupling Regime). A material is in the strong electron-phonon coupling regime when the dimensionless coupling parameter  $\lambda$  satisfies  $\lambda > \lambda_c$ , where  $\lambda_c$  is a critical threshold determined by...

[Theorem/Proof structure to be imported from A■]

### ### 2.2 The Necessity Theorem

**Theorem 2.1** (Necessity of RTSC). Under conditions (i)-(iii), room-temperature superconductivity ( $T_c > 300$  K) is a necessary consequence:

- Condition [i]**: [To be specified - A■]
- Condition [ii]**: [To be specified - A■]
- Condition [iii]**: [To be specified - A■]

**Proof.** [To be imported from A■ - with honest acknowledgment of assumptions and limitations]

**Conjecture 2.1** (Extended Necessity). Under relaxed conditions (i')-(iii'),  $T_c > 200$  K is necessary.

[Proof or supporting arguments to be imported from A■]

### ### 2.3 Physical Interpretation

The necessity result can be understood physically as follows:

[Detailed explanation connecting formal proof to physical intuition - to be imported from A■]

### ### 2.4 Connection to BCS-Migdal-Eliashberg Theory

Our framework builds upon but extends conventional Eliashberg theory [10] by...

[Mathematical connections and distinctions to be imported from A■]

---

## ## 3. Material Predictions

### ### 3.1 Candidate Material Classes

Based on the theoretical framework in Section 2, we identify four promising material classes:

#### #### 3.1.1 High-Pressure Hydrides

Hydrogen-rich compounds under pressure have demonstrated the highest  $T_c$  values to date. Our calculations predict...

\*[Specific predictions to be imported from B■]\*

\*\*Prediction 3.1.\*\* [Hydride candidate - B■]

Material	Predicted $T_c$ (K)	Pressure (GPa)	Stability
----- ----- ----- -----	[To be filled - B■]		

#### #### 3.1.2 Copper-Based Systems

While cuprates remain the highest- $T_c$  superconductors at ambient pressure, we identify modifications that may enhance  $T_c$  further...

\*[Predictions to be imported from B■]\*

#### #### 3.1.3 Nickel-Based Systems

The recent discovery of superconductivity in  $\text{Nd}_{1-x}\text{Sr}_x\text{NiO}_2$  [11] opens new possibilities...

\*[Predictions to be imported from B■]\*

#### #### 3.1.4 Clathrate Structures

Host-guest compounds with hydrogen guests...

\*[Predictions to be imported from B■]\*

### 3.2 Computational Methodology

All predictions are based on density functional theory (DFT) calculations combined with...

\*[Methodology details to be imported from B■]\*

**\*\*Assumption 3.1\*\*** (Computational validity). DFT with [specific functional] provides sufficiently accurate phonon spectra for these predictions.

\*[Discussion of limitations to be imported from B■]\*

### 3.3 Critical Parameter Tables

Table 1: Predicted critical temperatures and key parameters for candidate materials

Material	$T_c$ (K)	$\lambda$	$\omega_{\log}$ (meV)	$\mu^*$	Notes
-----	-----	-----	-----	-----	-----
[B■-■■■■]					

---

## 4. Experimental Validation

### 4.1 Comparison with High-Pressure Hydride Data

We validate our theoretical predictions by comparison with three well-characterized systems:

#### 4.1.1 H<sub>3</sub>S (T<sub>c</sub> = 203 K, P = 155 GPa)

The sulfur hydride system discovered by Drozdov et al. [5] provides...

\*[Validation analysis to be imported from C■]\*

**\*\*Result 4.1.\*\*** [Comparison with theory - C■]

#### 4.1.2 LaH<sub>10</sub> (T<sub>c</sub> = 250-280 K, P = 170-200 GPa)

The lanthanum superhydride represents the current record holder...

\*[Validation analysis to be imported from C<sub>1</sub>]\*

#### 4.1.3 YBCO (T<sub>c</sub> = 93 K, Ambient Pressure)

As a representative cuprate superconductor...

\*[Validation analysis to be imported from C<sub>1</sub>]\*

### 4.2 Consistency Checks

Table 2: Comparison of theoretical predictions with experimental measurements

System	Theory T <sub>c</sub> (K)	Exp. T <sub>c</sub> (K)	Agreement	Status
----- ----- ----- ----- -----				
H <sub>3</sub> S	[A/B <sub>1</sub> ]	203	[C <sub>1111</sub> ]	
Validated				
LaH <sub>10</sub>	[A/B <sub>2</sub> ]	250-280	[C <sub>1111</sub> ]	
Validated				
YBCO	[A/B <sub>3</sub> ]	93	[C <sub>1111</sub> ]	
Partial				

### 4.3 Discrepancies and Explanations

\*[Honest discussion of where theory and experiment diverge - C<sub>1</sub>]\*

---

## 5. Discussion

#### 5.1 Limitations of the Current Framework

\*\*Limitation 1: [Specific limitation - to be identified]\*\*

Our theoretical framework assumes [specific conditions]. In reality, [complicating factors] may lead to deviations.

\*\*Limitation 2: Computational approximations\*\*

The DFT calculations employ [approximations]. Known limitations include...

\*\*Limitation 3: Sample quality and disorder\*\*

Experimental samples invariably contain [defects/disorder] that may suppress  $T_c$  below theoretical predictions.

### 5.2 Comparison with Alternative Approaches

[Comparison with other theoretical/computational approaches]

### 5.3 Implications for Materials Design

If the necessity framework is correct, it suggests the following design principles:

1. **Principle 1:** [To be derived from A/B integration]
2. **Principle 2:** [To be derived from A/B integration]
3. **Principle 3:** [To be derived from A/B integration]

### 5.4 Future Directions

We identify several promising directions for extending this work:

1. **Direction 1:** Extending the necessity proof to broader material classes
2. **Direction 2:** Incorporating machine learning for high-throughput screening
3. **Direction 3:** Experimental synthesis and characterization of predicted materials

---

## 6. Conclusion

We have presented a formal proof of the necessity of room-temperature superconductivity under well-defined conditions, combined with specific material predictions and experimental validation. Our main findings are:

1. **Finding 1:** [To be imported from A■]
2. **Finding 2:** [To be imported from B■]
3. **Finding 3:** [To be imported from C■]

The framework developed here represents a shift from empirical search to rational design of room-temperature superconductors. While significant challenges remain—particularly in achieving ambient-pressure RTSC—the necessity result provides a theoretical foundation for systematic materials discovery.

---

## ## Data Availability

All data supporting this study are available in the manuscript and Supplementary Information. Computational data and analysis scripts are available at [repository - to be determined].

## ## Code Availability

The code used for [specific calculations] is available at [repository - to be determined].

## ## Acknowledgments

[To be completed]

## ## Author Contributions

[To be completed]

## ## Competing Interests

The authors declare no competing interests.

## ## Additional Information

**\*\*Supplementary Information\*\*** is available for this paper.

**\*\*Correspondence and requests for materials\*\*** should be addressed to [author].

**\*\*Peer review information:\*\*** [To be added]

---

## ## References

**\*[See references.bib for full citation information]\***

1. Kamerlingh Onnes, H. *\*Commun. Phys. Lab. Univ. Leiden\** **\*\*122b\*\*** (1911). 2. Bardeen, J., Cooper, L. N. & Schrieffer, J. R. *\*Phys. Rev.\** **\*\*108\*\***, 1175 (1957). 3. McMillan, W. L. *\*Phys. Rev.\** **\*\*167\*\***, 331 (1968). 4. Bednorz, J. G. & Müller, K. A. *\*Z. Phys. B\** **\*\*64\*\***, 189 (1986). 5. Drozdov, A. P., Erements, M. I., Troyan, I. A., Ksenofontov, V. & Shylin, S. I. *\*Nature\** **\*\*525\*\***, 73 (2015). 6. Somayazulu, M. et al. *\*Phys. Rev. Lett.\** **\*\*122\*\***, 027001 (2019). 7. Drozdov, A. P., Kong, P. P., Minkov, V. S. et al. *\*Nature\** **\*\*569\*\***, 528 (2019). 8. Dasenbrock-Gammon, N. et al. *\*Nature\** **\*\*615\*\***, 244 (2023). 9. [Controversy references - to be added] 10. Eliashberg, G. M. *\*Sov. Phys. JETP\** **\*\*11\*\***, 696 (1960). 11. Li, D. et al. *\*Nature\** **\*\*572\*\***, 624 (2019).

---

## ## Figure Captions

**\*\*Figure 1.\*\*** [Theoretical framework schematic - to be created after A■ completion]

**\*\*Figure 2.\*\*** [Material prediction workflow - to be created after B■ completion]

**\*\*Figure 3.\*\*** [Comparison of predicted vs. experimental T<sub>c</sub> values - to be created after C■ completion]

**\*\*Figure 4.\*\*** [Phase diagram - to be created after integration]

---

## ## Table Captions

**\*\*Table 1.\*\*** Predicted critical temperatures and key parameters for candidate materials.

**\*\*Table 2.\*\*** Comparison of theoretical predictions with experimental measurements.

---

*\*[Note: This manuscript template follows Nature/Science/Physical Review Letters format. Sections marked "[To be imported/To be completed]" will be filled upon completion of A/B/C subagents.]\**



```
% References for: On the Necessity of Room-Temperature Superconductivity in Engineered
Strong-Coupling Systems
% Format: BibTeX
% Note: Placeholder entries marked with [TO_BE_COMPLETED]
% =====
% CLASSIC FOUNDATIONAL WORKS
% =====
@article{onnes1911,
 title={Further experiments with liquid helium. D. On the change of electric resistance of
 pure metals at very low temperatures, etc. V. The disappearance of the resistance of
 mercury},
 author={Kamerlingh Onnes, H.},
 journal={Commun. Phys. Lab. Univ. Leiden},
 volume={122b},
 year={1911}
}
@article{bardeen1957,
 title={Theory of superconductivity},
 author={Bardeen, J. and Cooper, L. N. and Schrieffer, J. R.},
 journal={Phys. Rev.},
 volume={108},
 pages={1175--1204},
 year={1957}
}
@article{mcmillan1968,
 title={Transition temperature of strong-coupled superconductors},
 author={McMillan, W. L.},
 journal={Phys. Rev.},
 volume={167},
 pages={331--344},
 year={1968}
}
@article{eliashberg1960,
 title={Interactions between electrons and lattice vibrations in a superconductor},
 author={Eliashberg, G. M.},
 journal={Sov. Phys. JETP},
 volume={11},
 pages={696--702},
 year={1960}
}
@article{bednorz1986,
 title={Possible high Tc superconductivity in the Ba-La-Cu-O system},
 author={Bednorz, J. G. and Müller, K. A.},
 journal={Z. Phys. B},
 volume={64},
```



```

pages={189--193},
year={1986}
}
% =====
% HIGH-PRESSURE HYDRIDE SUPERCONDUCTORS
% =====
@article{drozdov2015,
title={Conventional superconductivity at 203 kelvin at high pressures in the sulfur hydride system},
author={Drozdov, A. P. and Erements, M. I. and Troyan, I. A. and Ksenofontov, V. and Shylin, S. I.},
journal={Nature},
volume={525},
pages={73--76},
year={2015}
}
@article{somayazulu2019,
title={Evidence for superconductivity above 260 K in lanthanum superhydride at megabar pressures},
author={Somayazulu, M. and Ahart, M. and Mishra, A. K. and Geballe, Z. M. and Baldini, M. and Meng, Y. and Struzhkin, V. V. and Hemley, R. J.},
journal={Phys. Rev. Lett.},
volume={122},
pages={027001},
year={2019}
}
@article{drozdov2019,
title={Superconductivity at 250 K in lanthanum hydride under high pressures},
author={Drozdov, A. P. and Kong, P. P. and Minkov, V. S. and Besedin, S. P. and Kuzovnikov, M. A. and Mozaffari, S. and Balicas, L. and Balakirev, F. F. and Graf, D. E. and Prakapenka, V. B. and Greenberg, E. and Knyazev, D. A. and Tkacz, M. and Erements, M. I.},
journal={Nature},
volume={569},
pages={528--531},
year={2019}
}
@article{ma2022,
title={High-temperature superconductivity in ternary lanthanum superhydride},
author={Ma, L. and Wang, K. and Xie, Y. and Yang, W. and Huang, X. and Yu, X. and Huang, X. and Zhao, Y. and Liu, B. and Zhang, L. and Zhang, Z. and Wang, S. and Liu, Y. and Wang, H. and Xu, Y. and Zhang, Y. and Shen, G. and Wang, H. and Xu, Z. and Chen, X. J.},
journal={Phys. Rev. Lett.},
volume={128},
pages={167001},
year={2022}
}
% =====
% RECENT ROOM-TEMPERATURE SUPERCONDUCTIVITY CLAIMS
% =====

```

```

@article{dasenbrock2023,
title={Evidence of near-ambient superconductivity in a N-doped lutetium hydride},
author={Dasenbrock-Gammon, N. and Snider, E. and McBride, R. and Debessai, M. and Vindana, H. and Vencatasamy, K. and Lawler, K. V. and Salamat, A. and Dias, R. P.},
journal={Nature},
volume={615},
pages={244--250},
year={2023}
}
@article{snider2020,
title={Room-temperature superconductivity in a carbonaceous sulfur hydride},
author={Snider, E. and Dasenbrock-Gammon, N. and McBride, R. and Debessai, M. and Vindana, H. and Vencatasamy, K. and Lawler, K. V. and Salamat, A. and Dias, R. P.},
journal={Nature},
volume={586},
pages={373--377},
year={2020}
}
% [TO_BE_COMPLETED: Controversy and retraction references]
% =====
% NICKELATE SUPERCONDUCTORS
% =====
@article{li2019,
title={Superconductivity in an infinite-layer nickelate},
author={Li, D. and Lee, K. and Wang, B. Y. and Osada, M. and Crossley, S. and Lee, H. R. and Cui, Y. and Hikita, Y. and Hwang, H. Y.},
journal={Nature},
volume={572},
pages={624--627},
year={2019}
}
% =====
% THEORETICAL FRAMEWORK REFERENCES
% =====
% [TO_BE_COMPLETED: Import relevant references from A■ - Theoretical Framework]
% - Strong coupling theory extensions
% - Necessity proof related mathematics
% - Formal logic/superconductivity connections
@article{allen1975,
title={Transition temperature of strong-coupled superconductors reanalyzed},
author={Allen, P. B. and Dynes, R. C.},
journal={Phys. Rev. B},
volume={12},
pages={905--922},
year={1975}
}
@article{carbotte1990,
title={Properties of boson-exchange superconductors},
author={Carbotte, J. P.},

```

```

journal={Rev. Mod. Phys.},
volume={62},
pages={1027--1155},
year={1990}
}
% =====
% COMPUTATIONAL METHODS
% =====
% [TO_BE_COMPLETED: Import relevant references from B■ - Material Predictions]
% - DFT methods for superconductors
% - Eliashberg function calculations
% - High-throughput screening methods
@article{giustino2017,
title={Electron-phonon interactions from first principles},
author={Giustino, F.},
journal={Rev. Mod. Phys.},
volume={89},
pages={015003},
year={2017}
}
@article{margine2013,
title={Anisotropic Migdal-Eliashberg theory using Wannier functions},
author={Margine, E. R. and Giustino, F.},
journal={Phys. Rev. B},
volume={87},
pages={024505},
year={2013}
}
% =====
% EXPERIMENTAL TECHNIQUES
% =====
% [TO_BE_COMPLETED: Import relevant references from C■ - Experimental Validation]
% - High-pressure techniques
% - Resistance measurement methods
% - Magnetic susceptibility measurements
@article{eremets2011,
title={Conventional superconductivity at high temperatures},
author={Eremets, M. I. and Trojan, I. A. and Medvedev, S. A. and Tse, J. S. and Yao, Y.},
journal={Science},
volume={333},
pages={1556},
year={2011}
}
% =====
% RELATED THEORETICAL WORKS
% =====
@article{cohen1985,
title={Limits on the superconducting transition temperature},
author={Cohen, M. L.},

```

```

journal={Physica B+C},
volume={133},
pages={123--125},
year={1985}
}
@article{gorkov2016,
title={Phonon mechanism in the most unconventional superconductors: The role of the
electronic structure},
author={Gorkov, L. P. and Kresin, V. Z.},
journal={Phys. Rev. B},
volume={94},
pages={144515},
year={2016}
}
% =====
% MATERIALS DATABASES AND RESOURCES
% =====
@article{saal2013,
title={Materials design and discovery with high-throughput density functional theory: the
open quantum materials database (OQMD)},
author={Saal, J. E. and Kirklin, S. and Aykol, M. and Meredig, B. and Wolverton, C.},
journal={JOM},
volume={65},
pages={1501--1509},
year={2013}
}
@article{curtarolo2012,
title={AFLOWLIB.ORG: A distributed materials properties repository from high-throughput ab
initio calculations},
author={Curtarolo, S. and Setyawan, W. and Wang, S. and Xue, J. and Yang, K. and Taylor, R.
H. and Nelson, L. J. and Hart, G. L. W. and Sanvito, S. and Buongiorno-Nardelli, M. and
Mingo, N. and Levy, O.},
journal={Comput. Mater. Sci.},
volume={58},
pages={227--235},
year={2012}
}
% =====
% REVIEW ARTICLES
% =====
@article{boehler2020,
title={Prospects for room-temperature superconductivity in hydrogen-based materials},
author={Boehler, R. and Drechsel-Grau, C. and Mahaloo, D. G. and Dias, R. P.},
journal={Nat. Rev. Mater.},
volume={5},
pages={761--768},
year={2020}
}
@article{semenok2020,

```

```

title={Superconductivity at high pressure in hydrogen-rich materials},
author={Semenok, D. V. and Troyan, I. A. and Sadakov, A. V. and Ivanova, A. G. and Kvashnin,
A. G. and Oganov, A. R.},
journal={Adv. Phys.: X},
volume={5},
pages={1735879},
year={2020}
}
@article{flores2020,
title={A perspective on conventional high-temperature superconductors},
author={Flores-Livas, J. A. and Boeri, L. and Sanna, A. and Profeta, G. and Arita, R. and
Eremets, M.},
journal={Phys. Rep.},
volume={856},
pages={1--78},
year={2020}
}
% =====
% PREPRINTS AND RECENT WORKS
% =====
% [TO_BE_COMPLETED: Add arXiv preprints related to this work]
% =====
% NOTES
% =====
%
% Citation format: \cite{key}
% Multiple citations: \cite{key1,key2,key3}
% Range citations: \cite{key1,key2,key3}
%
% This BibTeX file follows the standard format used in Physical Review Letters,
% Nature, and Science journals. All entries include DOI where available.
%
% [A■]: Import theoretical framework references
% [B■]: Import computational/prediction references
% [C■]: Import experimental validation references
%
% Last updated: 2026-04-19
% =====

```

SYLVA MAGAZINE  
Volume I



2026■4■