

Analytic Geometry 1

Liang Zheng
Australian National University
liang.zheng@anu.edu.au

3.1 Norms

- A norm on a vector space V is a function

$$\|\cdot\| : V \rightarrow \mathbb{R},$$

$$x \mapsto \|x\|, \text{ vector to real number}$$

which assigns each vector x its length $\|x\| \in \mathbb{R}$.

Examples

- The Manhattan norm on \mathbb{R}^n is defined for $\mathbf{x} \in \mathbb{R}^n$ as

L1 范式

$$\|\mathbf{x}\|_1 := \sum_{i=1}^n |x_i|, \quad \text{每个元素的绝对值的和 必定} \geq 0$$

$$\begin{bmatrix} x_1 \\ x_1 \\ \vdots \\ x_n \end{bmatrix}$$



where $|\cdot|$ is the absolute value. It is also called ℓ_1 norm.

- The Euclidean norm of $\mathbf{x} \in \mathbb{R}^n$ is defined as

$$\|\mathbf{x}\|_2 := \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{\mathbf{x}^T \mathbf{x}} \quad \text{每个元素的平方的和再开方}$$

It is the Euclidean distance of \mathbf{x} from the origin; also called


ℓ_2 norm




3.1 Norms

For all $\lambda \in \mathbb{R}$, and $\mathbf{x}, \mathbf{y} \in V$ the following holds:

- Absolutely homogeneous: $\|\lambda \mathbf{x}\| = |\lambda| \|\mathbf{x}\|$
- Triangle inequality: $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$
- Positive definite: $\|\mathbf{x}\| \geq 0$ and $\|\mathbf{x}\| = 0 \Leftrightarrow \mathbf{x} = \mathbf{0}$



Sparse representation Wright et al., TPAMI, 2009



Input

$$\underline{y} = \alpha_{i,1} \mathbf{v}_{i,1} + \alpha_{i,2} \mathbf{v}_{i,2} + \cdots + \alpha_{i,n_i} \mathbf{v}_{i,n_i}$$

$$A = [A_1, A_2, \dots, A_k] = [\mathbf{v}_{1,1}, \mathbf{v}_{1,2}, \dots, \mathbf{v}_{k,n_k}]$$

$$\underline{y} = A \underline{x_0} \text{ coefficients/weights}$$

$$\hat{x}_0 = \arg \min \|x\|_0 \text{ subject to } Ax = y$$

L0 norm, the number of non-zero elements

3.2.1 Dot Product

- Scalar product/dot product in \mathbb{R}^n is given by

$$\underbrace{x^T}_{1 \times n} \underbrace{y}_{n \times 1} = \sum_{i=1}^n x_i y_i \quad \mathbb{R}$$

点积是内积的一个特例，内积是双线性映射的一个特例。

Bilinear mapping

- A bilinear mapping Ω is a mapping with two arguments, and it is linear in each argument. Consider a vector space V , for all $x, y, z \in V, \lambda, \varphi \in \mathbb{R}$,

$$\Omega(\lambda x + \varphi y, z) = \lambda \Omega(x, z) + \varphi \Omega(y, z)$$

Ω is linear in the first argument

$$\Omega(x, \lambda y + \varphi z) = \lambda \Omega(x, y) + \varphi \Omega(x, z).$$

Ω is linear in the second argument

Inner product

两个参数的定义域

- Let V be a vector space and $\Omega: \boxed{V \times V} \rightarrow \mathbb{R}$ be a bilinear mapping.
- Ω is called symmetric if $\Omega(x, y) = \Omega(y, x)$
- Ω is called positive definite if

$$\underline{\forall x \in V \setminus \{0\} : \Omega(x, x) > 0, \quad \Omega(0, 0) = 0} \quad \text{即恒满足 } \geq 0$$

- A positive definite, symmetric bilinear mapping $\Omega: V \times V \rightarrow \mathbb{R}$ is called an **inner product** on V . We write $\langle x, y \rangle$ instead of $\Omega(x, y)$.
- The **pair** $(V, \langle \cdot, \cdot \rangle)$ is called is called an inner product vector space. If we use the dot product, we call $(V, \langle \cdot, \cdot \rangle)$ a Euclidean vector space.

Example

- Consider $V = \mathbb{R}^2$. If we define

$$\langle \mathbf{x}, \mathbf{y} \rangle := x_1 y_1 - \underbrace{(x_1 y_2 + x_2 y_1)} + 2x_2 y_2$$

- then $\langle \cdot, \cdot \rangle$ is an inner product but different from the dot product.

This mapping is symmetric: it is easy to derive $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$

Is it positive definite?

$$\forall \mathbf{x} \in V \setminus \{\mathbf{0}\}, \langle \mathbf{x}, \mathbf{x} \rangle = x_1^2 - (x_1 x_2 + x_2 x_1) + 2x_2^2 = (x_1 - x_2)^2 + x_2^2 > 0$$

3.2.3 Symmetric, Positive Definite Matrices

- Consider an n -dimensional vector space V with an inner product $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$, and a basis $B = (\mathbf{b}_1, \dots, \mathbf{b}_n)$ of V .

$$\langle \mathbf{x}, \mathbf{y} \rangle = \left\langle \sum_{i=1}^n \varphi_i \mathbf{b}_i, \sum_{j=1}^n \lambda_j \mathbf{b}_j \right\rangle = \sum_{i=1}^n \sum_{j=1}^n \varphi_i \langle \mathbf{b}_i, \mathbf{b}_j \rangle \lambda_j = \hat{\mathbf{x}}^T \mathbf{A} \hat{\mathbf{y}}$$

Basis vectors

where $A_{ij} := \langle \mathbf{b}_i, \mathbf{b}_j \rangle$ and $\hat{\mathbf{x}}, \hat{\mathbf{y}}$ are the coordinates of \mathbf{x}, \mathbf{y} with respect to the basis B .

- The inner product $\langle \cdot, \cdot \rangle$ is uniquely determined through \mathbf{A} . The symmetry of the inner product also means that \mathbf{A} is symmetric.
- The positive definiteness of the inner product implies that

$$\forall \mathbf{x} \in V \setminus \{\mathbf{0}\} : \langle \mathbf{x}, \mathbf{x} \rangle = \mathbf{x}^T \mathbf{A} \mathbf{x} > 0$$

3.2.3 Symmetric, Positive Definite Matrices

- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ that satisfies $\forall x \in V \setminus \{0\} : x^T A x > 0$ is called **symmetric, positive definite**, or just **positive definite**. If only \geq holds, then A is called **symmetric, positive semidefinite**.
- Example

$$A_1 = \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 3 \\ 3 & 3 \end{bmatrix}$$

- A_1 is positive definite because it is symmetric and

$$\begin{aligned} x^T A_1 x &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\ &= 3x_1^2 + 2x_1x_2 + 4x_2^2 = (x_1 + x_2)^2 + 2x_1^2 + 3x_2^2 > 0 \end{aligned}$$

for all $x \in V \setminus \{0\}$.

- A_2 is symmetric but not positive definite

$$x^T A_2 x = x_1^2 + 6x_1x_2 + 3x_2^2 = (x_1 + 3x_2)^2 - 6x_2^2 \text{ can be less than 0}$$

3.2.3 Symmetric, Positive Definite Matrices

- For a real-valued, finite-dimensional vector space V and a basis B of V , it holds that $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ is an inner product if and only if there exists a symmetric, positive definite matrix $A \in \mathbb{R}^{n \times n}$ with

$$\langle x, y \rangle = \hat{x}^T A \hat{y} \quad \text{一个内积有且只有一个正定矩阵.}$$

- If $A \in \mathbb{R}^{n \times n}$ is symmetric and positive definite,
the diagonal elements a_{ii} of A are positive because $a_{ii} = e_i^T A e_i = \langle e_i, e_i \rangle > 0$, where e_i is the i th vector of the standard basis in \mathbb{R}^n .

3.3 Lengths and Distances

- Any inner product induces a norm

$$\|x\| := \sqrt{\langle x, x \rangle}$$

- Cauchy-Schwarz Inequality
- For an inner product vector space $(V, \langle \cdot, \cdot \rangle)$ the induced norm $\|\cdot\|$ satisfies the Cauchy-Schwarz inequality

$$|\langle x, y \rangle| \leq \|x\| \|y\|$$

Example - Lengths of Vectors Using Inner Products

- We can now use an inner product to compute vector lengths, using $\|\mathbf{x}\| := \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$. Consider $\mathbf{x} = [1, 1]^T \in \mathbb{R}^2$. If we use the dot product as the inner product, we obtain

$$\|\mathbf{x}\| = \sqrt{\mathbf{x}^T \mathbf{x}} = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$\mathbf{x}^T \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{y} \text{ is dot product}$$

as the length of \mathbf{x} . Let us now choose a different inner product:

$$\langle \mathbf{x}, \mathbf{y} \rangle := \mathbf{x}^T \begin{bmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix} \mathbf{y} = x_1 y_1 - \frac{1}{2}(x_1 y_2 + x_2 y_1) + x_2 y_2$$

With this inner product, we obtain

$$\langle \mathbf{x}, \mathbf{x} \rangle = x_1^2 - x_1 x_2 + x_2^2 = 1 - 1 + 1 = 1 \Rightarrow \|\mathbf{x}\| = \sqrt{1} = 1$$

\mathbf{x} is “shorter” with this inner product than with the dot product.

3.3 Lengths and Distances

- Consider an inner product space $(V, \langle \cdot, \cdot \rangle)$, then

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}$$

is called the distance between x and y for $x, y \in V$.

- If we use the dot product as the inner product, then the distance is called Euclidean distance.

3.3 Lengths and Distances

- The mapping

$$\begin{aligned} d : V \times V &\rightarrow \mathbb{R} \\ (x, y) &\mapsto d(x, y) \end{aligned}$$

is called a metric.

- A metric d satisfies the following:
- d is positive definite, i.e., $d(x, y) \geq 0$ for all $x, y \in V$ and $d(x, y) = 0 \Leftrightarrow x = y$
- d is symmetric, i.e., $d(x, y) = d(y, x)$ for all $x, y \in V$
- Triangle inequality: $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in V$
- Very similar x and y will result in a large value for the inner product and a small value for the metric.


3.4 Angles and Orthogonality

$$|\langle x, y \rangle| \leq \|x\| \|y\|$$

- According to Cauchy-Schwarz inequality, assume $x \neq 0$, $y \neq 0$. Then,

$$-1 \leq \frac{\langle x, y \rangle}{\|x\| \|y\|} \leq 1$$

Therefore, there exists a unique $\omega \in [0, \pi]$, with



$$\cos \omega = \frac{\langle x, y \rangle}{\|x\| \|y\|}$$

The number ω is the angle between the vectors x and y .


度量方向相似性

- The angle between two vectors tells us how similar their orientations are.
- Using the dot product, the angle between x and $y = 4x$ is 0, so their orientation is the same.

$$\cos \omega = \frac{\langle x, 4x \rangle}{\|x\| \|4x\|} = \frac{4\langle x, x \rangle}{\sqrt{x^T x} \sqrt{(4x)^T (4x)}} = \frac{4\langle x, x \rangle}{4\|x\| \|x\|} = \frac{\langle x, x \rangle}{\|x\| \|x\|} = 1$$

Example (Angle between Vectors)

- Let us compute the angle between $\mathbf{x} = [-1, 1]^T \in \mathbb{R}^2$ and $\mathbf{y} = [2, 1]^T \in \mathbb{R}^2$. We use the dot product as the inner product. We get



$$\cos \omega = \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\sqrt{\langle \mathbf{x}, \mathbf{x} \rangle \langle \mathbf{y}, \mathbf{y} \rangle}} = \frac{\mathbf{x}^T \mathbf{y}}{\sqrt{\mathbf{x}^T \mathbf{x} \mathbf{y}^T \mathbf{y}}} = \frac{-1}{\sqrt{10}}$$

- and the angle between the two vectors is $\arccos\left(\frac{-1}{\sqrt{10}}\right) \approx 1.89\text{rad}$, which corresponds to about 108.4° .
- We then use inner product to characterize orthogonality.

3.4 Angles and Orthogonality


- Two vectors \mathbf{x} and \mathbf{y} are orthogonal if and only if $\langle \mathbf{x}, \mathbf{y} \rangle = 0$, and we write $\mathbf{x} \perp \mathbf{y}$. If additionally $\|\mathbf{x}\| = \|\mathbf{y}\| = 1$, i.e., the vectors are unit vectors, then \mathbf{x} and \mathbf{y} are orthonormal.
标准正交的
- $\mathbf{0}$ -vector is orthogonal to every vector in the vector space

- Example (Orthogonal Vectors)
- Consider $\mathbf{x} = [1, 2]^T$ and $\mathbf{y} = [-4, 2]^T$
- Using dot product as inner product, we have
 - $\langle \mathbf{x}, \mathbf{y} \rangle = 0$, so $\mathbf{x} \perp \mathbf{y}$.
- if we choose the inner product

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} \mathbf{y}$$

- the angle ω between \mathbf{x} and \mathbf{y} is given by 使用的内积会影响角度的度量

$$\cos \omega = \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\|\mathbf{x}\| \|\mathbf{y}\|} = -\frac{2}{\sqrt{17 \times 12}} \Rightarrow \omega \approx 1.43 \text{ rad} \approx 81.95^\circ$$



3.4 Angles and Orthogonality

- A square matrix $A \in \mathbb{R}^{n \times n}$ is an orthogonal matrix if and only if its columns are orthonormal, such that

$$\underline{A A^T = I = A^T A}$$

which implies that

$$\underline{A^{-1} = A^T}$$

i.e., the inverse is obtained by simply transposing the matrix

Properties - length

- The length of a vector \mathbf{x} is not changed when transforming it using an orthogonal matrix \mathbf{A} . For dot product, we obtain

$$\|\mathbf{Ax}\|^2 = (\mathbf{Ax})^T (\mathbf{Ax}) = \mathbf{x}^T \mathbf{A}^T \mathbf{Ax} = \mathbf{x}^T \mathbf{I} \mathbf{x} = \mathbf{x}^T \mathbf{x} = \|\mathbf{x}\|^2$$

Properties - angle

- The **angle** between any two vectors \mathbf{x} and \mathbf{y} as measured by their inner product, is also unchanged when transforming both of them using an orthogonal matrix \mathbf{A} . We use the dot product as inner product

$$\cos \omega = \frac{(\mathbf{Ax})^T (\mathbf{Ay})}{\|\mathbf{Ax}\| \|\mathbf{Ay}\|} = \frac{\mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{y}}{\sqrt{\mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{x} \mathbf{y}^T \mathbf{A}^T \mathbf{A} \mathbf{y}}} = \frac{\mathbf{x}^T \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

使用同一个标准正交矩阵同时对 \mathbf{x}, \mathbf{y} 进行变换

- Orthogonal matrices A with $A^{-1} = A^T$ preserve both angles and distances.
- Orthogonal matrices define transformations that are rotations

3.5 Orthonormal Basis

- Consider an n -dimensional vector space V and a basis $\{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ of V . For all $i, j = 1, \dots, n$, if

$$\langle \mathbf{b}_i, \mathbf{b}_j \rangle = 0 \quad \text{for } i \neq j \quad (1)$$

$$\langle \mathbf{b}_i, \mathbf{b}_i \rangle = 1$$

then the basis is called an orthonormal basis (ONB).

If only (1) is satisfied, the basis is called an orthogonal basis.

Example (Orthonormal Basis)


- The canonical/standard basis for a Euclidean vector space \mathbb{R}^n is an orthonormal basis, where the inner product is the dot product of vectors

For \mathbb{R}^3 : $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$

- In \mathbb{R}^2 , the vectors $b_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, b_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ form an orthonormal basis since $b_1^\top b_2 = 0$ and $\|b_1\| = 1 = \|b_2\|$

3.6 Orthogonal Complement

- A hyperplane U in a three-dimensional vector space can be described by its normal vector, which spans its orthogonal complement U^\perp 法向量



- Generally, orthogonal complements can be used to describe hyperplanes in n -dimensional vector and affine spaces

$(n - 1)$ dimensional

3.6 Orthogonal Complement

- We now look at vector spaces that are orthogonal to each other
- Consider a D -dimensional vector space V and an M -dimensional subspace $U \subseteq V$. The **orthogonal complement** U^\perp is a $(D - M)$ -dimensional subspace of V and contains all vectors in V that are orthogonal to every vector in U .
- $U \cap U^\perp = \{\mathbf{0}\}$ so that any vector $\mathbf{x} \in V$ can be uniquely decomposed into

$$\mathbf{x} = \sum_{m=1}^M \lambda_m \mathbf{b}_m + \sum_{j=1}^{D-M} \psi_j \mathbf{b}_j^\perp, \lambda_m, \psi_j \in \mathbb{R}$$

- Where $(\mathbf{b}_1, \dots, \mathbf{b}_M)$ is a basis of U and $(\mathbf{b}_1^\perp, \dots, \mathbf{b}_{D-M}^\perp)$ is a basis of U^\perp .

i.e. $U + (U^\perp) = \text{original vector space } V$

可以使用 \mathbf{b} 与 (\mathbf{b}^\perp) 的线性组合表示 V 中的任意向量

Check your understanding

- (A) Norm characterises the length of a vector. ✓
- (B) The norm of a vector can be a Real complex number. ✗
- (C) The inner product takes 2 ~~one~~ vector as input and outputs a real number. ✗
- (D) A metric characterises the similarity between two vectors. ✓
- (E) Any bilinear mapping introduces an inner product ✗
- (F) Any inner product introduces a norm ✓
- (G) Any vector in U^\perp is orthogonal to any vector in U . ✓
- (H) In \mathbb{R}^2 there can be infinitely many bases, but only a finite number of orthogonal / orthonormal bases ✗