

# YK\_Final\_P3

Yinan Kang


5/13/2019



```
require(dplyr)
require(leaps)
require(neuralnet)
require(readxl)
require(caret)
require(lmtest)
```


```
rm(list=ls())
df.3 <- read.csv("/cloud/project/Question 3.csv")
```

## Fit Model

```
model.3 <- lm(Y~., data=df.3)
plot(model.3)
```







**Analysis:** We see unequal variance (Megaphone shape) in the Residuals Plot

## Using Weighted Least Squares

```
ei <- model.3$residuals
abs.ei <- abs(ei)
model.3.1 <- lm(abs.ei ~ ., data=df.3)

s <- model.3.1$fitted.values
wi <- 1/(s^2)

model.3.new <- lm(Y~., weights=wi, data=df.3)
summary(model.3.new)
```

```
##
## Call:
## lm(formula = Y ~ ., data = df.3, weights = wi)
##
## Weighted Residuals:
##      Min       1Q   Median       3Q      Max
## -2.3041 -0.4933  0.2369  0.9425  2.3723
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   16.2356     1.5265  10.636 1.60e-15 ***
## X1             12.4000     0.7930  15.636 < 2e-16 ***
## X2              1.2718     0.1861   6.833 4.49e-09 ***
## X3              1.5929     0.3179   5.011 4.93e-06 ***
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.129 on 61 degrees of freedom
## Multiple R-squared:  0.8878, Adjusted R-squared:  0.8823
## F-statistic: 160.9 on 3 and 61 DF,  p-value: < 2.2e-16
```

```
plot(model.3.new)
```

