

學名	商品名	外滲特性	處置	備註
Arsenic trioxide	Asadin	None	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Asparaginase	Leunase	None	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Azacitidine	Vidaza	None	Cold compression with ice pack for 20 min QID for 1-2 days (4)	Given SC, IF
Bendamustine	Innomustine	Irritant with vesicant potential	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Bleomycin	Bleocin	None	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Bortezomib	Velcade	Possible irritant	Cold compression with ice pack for 20 min QID for 1-2 days (4)	Given SC, IV
Brentuximab vedotin	Adcetris	Irritant (vesicant reactions reported)	Cold compression (8)	Treat as vesicant
Busulfan	Busulfex	Vesicant	Cold compression with ice pack for 20 min QID for 1-2 days (1,4)	
Cabazitaxel	Jevtana	None	Cold compression with ice pack for 20 min QID for 1-2 days (1,4) OR warm compression (3,5) *	
Carboplatin	Paraplatin	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (1,4) OR warm compression (3,5) *	
Carfilzomib	Kyprolis	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (5)	
Carmustine	Carmuther	Vesicant	Cold compression with ice pack for 20 min QID for 1-2 days (1, 4)	
Cisplatin	Kemoplat	Irritant (vesicant if undiluted)	1. Cold compression with ice pack for 20 min QID for 1-2 days (1,4) OR warm compression (3,5) * 2. 大體積未稀釋 cisplatin 外滲: 可用 Sodium thiosulfate (1/6 M or 4%) 2mL into IV line. 4% Sodium thiosulfate: 配製見註記**	Exfoliant reactions
Cladribine	Leustatin	None	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Clofarabine	Evoltra	None	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Cyclophosphamide	Endoxan	None	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Cytarabine	Cytosar	None	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Dacarbazine	Dacarbazine	Irritant (vesicant reactions reported)	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Dactinomycin	Dactilon, Cosmegen	Vesicant	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Decitabine	Dacogen	None	Cold compression with ice pack for 20 min QID for 1-2 days (5)	
Docetaxel	Taxotere	Irritant (vesicant reactions reported)		

學名	商品名	外滲特性	處置	備註
Doxorubicin	Adriamycin	Vesicant	1. 99% DMSO, topical, to twice the affected area, allow to air dry. Repeat QID x 7 days (1,3,4) 2. Cold compression with ice pack for 20 min QID for 1-2 days (1,2)	
Doxorubicin, liposomal	Lipodox	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (2, 4)	
Enfortumab vedotin	Padcev	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (6)	
Epirubicin	Pharmarubicin	Vesicant	1. 99% DMSO, topical, to twice the affected area, allow to air dry. Repeat QID x 7 days (1,3,4) 2. Cold compression with ice pack for 20 min QID for 1-2 days (1,2)	
Eribulin	Halaven	None	Cold compression with ice pack for 20 min QID for 1-2 days (4, 5)	
Etoposide	Fytosid, Vepesid, Eposin	Irritant	Warm compression (44-50°C) for 20-30 min, QID for 1-2 days. (5, 6)	
Fludarabine	Fludara	None	Cold compression with ice pack for 20 min QID for 1-2 days (4, 5)	
Fluorouracil	5-FU injection	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (4, 5)	
Gemcitabine	Gemmis, Gemzar	None	Cold compression with ice pack for 20 min QID for 1-2 days (4, 5)	
Gemtuzumab ozogamicin	Mylotarg	None	Cold compression with ice pack for 20 min QID for 1-2 days (4, 5)	
Idarubicin	Zavedos	Vesicant	1. 99% DMSO, topical, to twice the affected area, allow to air dry. Repeat QID x 7 days (1,3,4) 2. Cold compression with ice pack for 20 min QID for 1-2 days (1,2)	
Ifosfamide	Holoxan	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Inotuzumab ozogamicin	Besponsa	None	Not indicated (7)	
Irinotecan	Campto	None	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Irinotecan, liposomal	Onivyde	Not documented	Cold compression with ice pack for 20 min QID for 1-2 days (7)	
Ixabepilone	Ixemptra	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (5)	
Melphalan	Alkeran	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Methotrexate	Methotrexate	None	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Mitomycin	Vesimycin, Mitonco,	Vesicant	1. 99% DMSO, topical, to twice the affected area, allow to air dry. Repeat QID x 7 days (1,3,4) 2. Cold compression with ice pack for 20 min QID for 1-2 days (1,2)	
Mitoxantrone	Mitoxantrone, Novantrone	Irritant	1. 99% DMSO, topical, to twice the affected area, allow to air dry. Repeat QID x 7 days (1,3,4) 2. Cold compression with ice pack for 20 min QID for 1-2 days (1,2)	

學名	商品名	外滲特性	處置	備註
Nelarabine	Arranon, Atriance	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (2,4)	
Oxaliplatin	Oxalip, Eloxatin	Irritant,	Warm compression (44-50°C) for 20-30 min, QID for 1-2 days. (2, 4)	Treat as vesicant
Paclitaxel	Phyxol, Taxol	Irritant,	Cold compression with ice pack for 20 min QID for 1-2 days (1,4) OR warm compression (3,5) *	Treat as vesicant
Paclitaxel, albumin-bound	Abraxane	Irritant,	Cold compression with ice pack for 20 min QID for 1-2 days (1,4) OR warm compression (3,5) *	Treat as vesicant
Pemetrexed	Alimta	None	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Polatuzumab vedotin	Polivy	Irritant	Cold compression with ice pack for 20 min QID for 1-2 days (7)	
Pralatrexate	Folotylin	None	Not indicated (7)	
Sacituzumab govitecan	Trodelvy	Irritant (7)	Cold compression with ice pack for 20 min QID for 1-2 days (7)	
Thiotepa	Tepadina	None	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Topotecan	Hycamtin	None	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Trabectedin	Yondelis	Vesicant	Cold compression with ice pack for 20 min QID for 1-2 days (4)	
Trastuzumab deruxtecan	Enhertu	No data	Cold compression with ice pack for 20 min QID for 1-2 days (4)	*無資料，依 trastuzumab emtansine 和 irinotecan 處置
Trastuzumab emtansine	Kadcyla	Irritant (vesicant reactions reported)	Cold compression with ice pack for 20 min QID for 1-2 days (4,5)	
Vinblastine	Vinblastine	Vesicant	Warm compression (44-50°C) for 20-30 min, QID for 1-2 days. (2, 4)	
Vincristine	Vincristine	Vesicant	Warm compression (44-50°C) for 20-30 min, QID for 1-2 days. (2, 4)	
Vinorelbine	Navelbine	Vesicant	Warm compression (44-50°C) for 20-30 min, QID for 1-2 days. (2, 4)	

註記：

冷敷處置：0°C用冰敷袋或冰塊包毛巾，每次 20 分鐘，一天 3-4 次，持續 24-48 小時。

熱敷處置：44-50°C，每次 20-30 分鐘，一天 3-4 次持續 24-48 小時。

*部分藥品(platinum, 紫杉醇類)該冷敷或熱敷目前無定論。

**1/6 M (約 4%) 硫代硫酸鈉 sodium thiosulfate 配製：取 12.5g/50 mL (25%) sodium thiosulfate inj 1.6 mL，加入 8.4 mL 注射用水

Ref:

1. BC Cancer Extravasation guidelines (Canada)
2. Drug Information Handbook for Oncology 17th Ed.
3. ESMO-EONS Guidelines
4. Support Care Cancer (2015) 23:1459–1471
5. NHS Trust (Salisbury, Thames Valley)(UK)
6. UptoDate
7. Cancer Institute NSW (Australia)
8. Dermatol Ther. 2022 Oct; 35(10): e15751.