


Advanced Topics in Astrodynamics and Trajectory Design

Circular Restricted Three Body Problem

Dr Joan Pau Sánchez

MSc in Astronautics and Space Engineering



Bibliography

Some good books ordered by increasing depth and breath :

- *Orbital Mechanics*, V.A Chobotov, AIAA Education Series
- *Analytical Mechanics of Space Systems*, H.Shcaub, J.L.Junkins, AIAA Education Series
- *An Introduction to the Mathematics and Methods of Astrodynamics*, R.H.Battin, AIAA Educationan Series
- *Fundamentals of Astrodynamics and Applications*, D.A.Vallado, Space Technology Library

Circular Restricted Three Body Problem (CR3BP)

- “**Three-body**” refers to a problem that considers the motion of 3 bodies. Generally, 2 celestial bodies + spacecraft; Earth+Moon+S/C, Sun+Jupiter+asteroid.
- “**Restricted**” refers to the consideration that two of these bodies have masses that far outweigh the mass of the third.
- “**Circular**” refers to the fact that these two large bodies move in circular orbits around their common centre of mass.

CR3BP: General Description




Figure 9.1: Illustration of Three-Body Problem
 Courtesy (Shcaub & Junkins, 2002)

CR3BP: General Description

- “**Restricted**” refers to the consideration that two of these bodies have masses that far outweigh the mass of the third.
 - ✓ $m_3 \sim 0$
- “**Circular**” refers to the fact that these two large bodies move in circular orbits around their common centre of mass.
 - ✓ **Constant angular velocity vector ω**

$$n = \frac{2\pi}{\tau} = \sqrt{\frac{GM}{a^3}} \quad \longrightarrow \quad \omega = \sqrt{\frac{G(m_1 + m_2)}{r_{12}^3}}$$

CR3BP: Synodic or rotating reference frame

- A reference frame F centred at the barycentre and rotating with an angular velocity $\omega = \omega \hat{\mathbf{k}}$






Figure 9.8: Illustration of Circular Restricted Three-Body Problem


Courtesy (Shcaub & Junkins, 2002)

CR3BP: Synodic or rotating reference frame

Inertial orbit in inertial RF




Inertial orbit in non-inertial RF



CR3BP: Synodic or rotating reference frame

Non-inertial orbit in non-inertial RF



Exercise 1

DemoEx1_CR3BP_Session4.m

1. Open file
“DemoEx1_CR3BP_Session4.m”
2. Run each section in order one time.

1. Change the orbit by choosing a different apoapsis and periapsis distance.

1. Attempt the following example:

$$r_p=0.4u; r_a=0.86u$$

2. Attempt the following example:

$$r_p=0.2u; r_a=0.7615u$$

What is happening in these strange orbits?

CR3BP: Equations of motion – Vector form

- Express \mathbf{r} in $F : \{\hat{\mathbf{e}}_r, \hat{\mathbf{e}}_\theta, \hat{\mathbf{e}}_z\}$

$$\omega = \sqrt{\frac{G(m_1 + m_2)}{r_{12}^3}}$$

$$\mathbf{r} = r_x \hat{\mathbf{e}}_r + r_y \hat{\mathbf{e}}_\theta + r_z \hat{\mathbf{e}}_z$$




Figure 9.8: Illustration of Circular Restricted Three-Body Problem

$$\ddot{\mathbf{r}} + \underbrace{2\boldsymbol{\omega} \times \dot{\mathbf{r}}}_{\text{Coriolis}} + \underbrace{\boldsymbol{\omega} \times \boldsymbol{\omega} \times \mathbf{r}}_{\text{Centripetal}} = \frac{\mathbf{F}_1}{m} + \frac{\mathbf{F}_2}{m}$$

Coriolis Centripetal

$$\mathbf{F}_1 = -\frac{GmM_1}{\xi_1^3} \boldsymbol{\xi}_1$$

$$\mathbf{F}_2 = -\frac{GmM_2}{\xi_2^3} \boldsymbol{\xi}_2$$

$$\xi_i = \sqrt{(r_x - r_i)^2 + r_y^2 + r_z^2}$$

CR3BP: Equations of motion – scalar form

$$\left. \begin{aligned} \ddot{r}_x - 2\omega\dot{r}_y - \omega^2 r_x &= -G \left(\frac{m_1}{\xi_1^3} (r_x - r_1) + \frac{m_2}{\xi_2^3} (r_x - r_2) \right) \\ \ddot{r}_y + 2\omega\dot{r}_x - \omega^2 r_y &= -G \left(\frac{m_1}{\xi_1^3} + \frac{m_2}{\xi_2^3} \right) r_y \\ \ddot{r}_z &= -G \left(\frac{m_1}{\xi_1^3} + \frac{m_2}{\xi_2^3} \right) r_z \end{aligned} \right\}$$

$$\mathbf{a} = -\frac{Gm}{r^2} \left(\frac{\mathbf{r}}{r} \right)$$




Figure 9.8: Illustration of Circular Restricted Three-Body Problem

CR3BP: Equations of motion – Potential function

$$U(r_x, r_y, r_z) = \frac{\omega^2}{2} (r_x^2 + r_y^2) + \frac{Gm_1}{\xi_1} + \frac{Gm_2}{\xi_2}$$

$$\ddot{\mathbf{r}} + 2\boldsymbol{\omega} \times \dot{\mathbf{r}} = \frac{\partial U}{\partial \mathbf{r}}$$




Figure 9.8: Illustration of Circular Restricted Three-Body Problem

CR3BP: Jacobi Integral

$$\ddot{\mathbf{r}} + 2\boldsymbol{\omega} \times \dot{\mathbf{r}} = \frac{\partial U}{\partial \mathbf{r}}$$

$$(\ddot{\mathbf{r}} + 2\boldsymbol{\omega} \times \dot{\mathbf{r}}) \cdot \dot{\mathbf{r}} = \frac{\partial U}{\partial \mathbf{r}} \cdot \dot{\mathbf{r}}$$

$$\ddot{\mathbf{r}} \cdot \dot{\mathbf{r}} + 2\boldsymbol{\omega} \times \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} = \frac{\partial U}{\partial \mathbf{r}} \cdot \frac{d\mathbf{r}}{dt}$$

$$\ddot{\mathbf{r}} \cdot \dot{\mathbf{r}} = \frac{dU}{dt}$$

$$\frac{1}{2} \frac{d}{dt} (\dot{\mathbf{r}} \cdot \dot{\mathbf{r}}) = \frac{dU}{dt}$$

$$v^2 = 2U - C$$

$$C = 2U - v^2$$

- Jacobi integral/constant = C

CR3BP: Jacobi Integral

$$C = \omega^2 (r_x^2 + r_y^2) + \frac{2Gm_1}{\xi_1} + \frac{2Gm_2}{\xi_2} - v^2$$

- Basically, C is the classical energy integral expressed in rotating reference frame.
- Jacobi integral has two important uses:
 - ✓ Verifying accuracy of numerical integration.
 - ✓ Indicating regions of feasible motion.

CR3BP: Non-dimensional nomenclature

- The problem can be made non-dimensional by choosing the following units:
 - ✓ $m_1 + m_2$ as the unit of mass
 - ✓ r_{12} as the unit of length
 - ✓ $P/(2\pi)$ as the unit of time

$$\mu = \frac{m_2}{m_1 + m_2}$$

$$x_2 - x_1 = 1$$

$$x_1 = -\mu$$

$$x_2 = 1 - \mu$$




Figure 9.8: Illustration of Circular Restricted Three-Body Problem

CR3BP: Non-dimensional nomenclature

| System | μ | L | V | T |
|------------------|------------------------|---------------------|--------|---------------------|
| Sun-Jupiter | 9.537×10^{-4} | 7.784×10^8 | 13.102 | 3.733×10^8 |
| Sun-(Earth+Moon) | 3.036×10^{-6} | 1.496×10^8 | 29.784 | 3.147×10^7 |
| Earth-Moon | 1.215×10^{-2} | 3.850×10^5 | 1.025 | 2.361×10^6 |
| Mars-Phobos | 1.667×10^{-8} | 9.380×10^3 | 2.144 | 2.749×10^4 |
| Jupiter-Io | 4.704×10^{-5} | 4.218×10^5 | 17.390 | 1.524×10^5 |
| Jupiter-Europa | 2.528×10^{-5} | 6.711×10^5 | 13.780 | 3.060×10^5 |
| Jupiter-Ganymede | 7.804×10^{-5} | 1.070×10^6 | 10.909 | 6.165×10^5 |
| Jupiter-Callisto | 5.667×10^{-5} | 1.883×10^6 | 8.226 | 1.438×10^6 |
| Saturn-Mimas | 6.723×10^{-8} | 1.856×10^5 | 14.367 | 8.117×10^4 |
| Saturn-Titan | 2.366×10^{-4} | 1.222×10^6 | 5.588 | 1.374×10^6 |
| Neptune-Triton | 2.089×10^{-4} | 3.548×10^5 | 4.402 | 5.064×10^5 |
| Pluto-Charon | 1.097×10^{-1} | 1.941×10^4 | 0.222 | 5.503×10^5 |

TABLE 2.2.1. **Table of m_1 - m_2 systems in the solar system.** Source: The first three are the values used in Koon, Lo, Marsden, and Ross [2000, 2001b]. The others are from the Jet Propulsion Laboratory's solar system dynamics website: <http://ssd.jpl.nasa.gov/>.

CR3BP: Non-dimensional dynamics

$$\left. \begin{aligned} \ddot{x} - 2\dot{y} - x &= -\frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{-\mu}{\rho_2^3}(x-x_2) \\ \ddot{y} + 2\dot{x} - y &= -\left(\frac{1-\mu}{\rho_1^3} + \frac{-\mu}{\rho_2^3}\right)y \\ \ddot{z} &= -\left(\frac{1-\mu}{\rho_1^3} + \frac{-\mu}{\rho_2^3}\right)z \end{aligned} \right\} \quad \rho_i = \sqrt{(x-x_i)^2 + y^2 + z^2}$$

$$U(x, y, z) = \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{\rho_1} + \frac{\mu}{\rho_2}$$

$$v^2 = x^2 + y^2 + \frac{2(1-\mu)}{\rho_1} + \frac{2\mu}{\rho_2} - C$$

CR3BP: Equilibrium Positions

- Let say we have a trajectory with a state vector such as:

$$(\mathbf{x}, \dot{\mathbf{x}}) = (x, y, z, 0, 0, 0)$$

$$\ddot{x} = 2\dot{y} + x - \frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{\mu}{\rho_2^3}(x-x_2)$$

$$\ddot{y} = -2\dot{x} + y - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) y$$

$$\ddot{z} = - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) z$$

CR3BP: Equilibrium Positions

- Let say we have a trajectory with a state vector such as:

$$(\mathbf{x}, \dot{\mathbf{x}}) = (x, y, z, 0, 0, 0)$$

$$\ddot{x} = 0$$

$$\ddot{y} = 0$$

$$\ddot{z} = 0$$

- We then have an equilibrium point. (or stationary point).

CR3BP: Equilibrium Positions

- Let's look for: $(\mathbf{x}, \dot{\mathbf{x}}, \ddot{\mathbf{x}}) = (x, y, z, 0, 0, 0, 0, 0, 0)$

$$\ddot{x} = 2\dot{y} + x - \frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{\mu}{\rho_2^3}(x-x_2)$$

$$\ddot{y} = -2\dot{x} + y - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) y$$

$$\ddot{z} = - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) z \quad \longrightarrow \quad \ddot{z} = 0 \rightarrow z = 0$$


- All stationary points must lie in the orbital plane of m_1 & m_2

CR3BP: Equilibrium Positions

- Let's look for: $(\mathbf{x}, \dot{\mathbf{x}}, \ddot{\mathbf{x}}) = (x, y, 0, 0, 0, 0, 0, 0)$

$$\ddot{x} = 2\dot{y} + x - \frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{\mu}{\rho_2^3}(x-x_2)$$

$$\ddot{y} = -2\dot{x} + y - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) y$$



$y = 0$ *Collinear points*


$$y - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) y = 0$$

$$\left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) = 1; \rho_1 = \rho_2 \quad \text{Equilateral points}$$

CR3BP: Equilibrium Positions (Collinear Solutions)

- Let's look for: $(\mathbf{x}, \dot{\mathbf{x}}, \ddot{\mathbf{x}}) = (x, 0, 0, 0, 0, 0, 0, 0, 0)$
- $$\ddot{x} = 2\dot{y} + x - \frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{\mu}{\rho_2^3}(x-x_2)$$
- $$\rho_1 = \sqrt{(x-x_1)^2} \quad x_1 = -\mu$$
- $$\rho_2 = \sqrt{(x-x_2)^2} \quad x_2 = 1-\mu$$

$$x - \frac{1-\mu}{|x-x_1|^3}(x-x_1) - \frac{\mu}{|x-x_2|^3}(x-x_2) = 0$$



L3: $x - x_1 < 0$ & $x - x_2 < 0$

$$x + \frac{1-\mu}{(x-x_1)^2} + \frac{\mu}{(x-x_2)^2} = 0$$

L1: $x - x_1 > 0$ & $x - x_2 < 0$

$$x - \frac{1-\mu}{(x-x_1)^2} + \frac{\mu}{(x-x_2)^2} = 0$$

L2: $x - x_1 > 0$ & $x - x_2 > 0$

$$x - \frac{1-\mu}{(x-x_1)^2} - \frac{\mu}{(x-x_2)^2} = 0$$

Exercise 2A

Part A in:

DemoEx2_CR3BP_Session4.m

- For the Earth-Moon system, find the three collinear equilibrium points.

Hint: Use *fzero* function (i.e. Matlab) to solve the equations below.

$$\text{L3: } x - x_1 < 0 \ \& \ x - x_2 < 0 \quad x + \frac{1 - \mu}{(x - x_1)^2} + \frac{\mu}{(x - x_2)^2} = 0$$

$$\text{L1: } x - x_1 > 0 \ \& \ x - x_2 < 0 \quad x - \frac{1 - \mu}{(x - x_1)^2} + \frac{\mu}{(x - x_2)^2} = 0$$

$$\text{L2: } x - x_1 > 0 \ \& \ x - x_2 > 0 \quad x - \frac{1 - \mu}{(x - x_1)^2} - \frac{\mu}{(x - x_2)^2} = 0$$

Constants:

$$m_{\text{Earth}} = 5.9737 \times 10^{24} \text{ kg}$$

$$m_{\text{Moon}} = 7.3476 \times 10^{22} \text{ kg}$$

Exercise 2B


Part B in:

DemoEx2_CR3BP_Session4.m

For each equilibrium solution, propagate its initial condition and plot the results, both in rotating reference frame and inertial reference frame.

- Can you explain what you see?

CR3BP: Five Fixed Points (Lagrange or Libration Points)



CR3BP: Zero velocity curves

$$\left. \begin{aligned} \ddot{x} - 2\dot{y} - x &= -\frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{-\mu}{\rho_2^3}(x-x_2) \\ \ddot{y} + 2\dot{x} - y &= -\left(\frac{1-\mu}{\rho_1^3} + \frac{-\mu}{\rho_2^3}\right)y \\ \ddot{z} &= -\left(\frac{1-\mu}{\rho_1^3} + \frac{-\mu}{\rho_2^3}\right)z \end{aligned} \right\} \quad \rho_i = \sqrt{(x-x_i)^2 + y^2 + z^2}$$

$$U(x, y, z) = \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{\rho_1} + \frac{\mu}{\rho_2}$$

$$v^2 = x^2 + y^2 + \frac{2(1-\mu)}{\rho_1} + \frac{2\mu}{\rho_2} - C$$

CR3BP: Zero velocity curves

$$v(x, y, z)^2 = x^2 + y^2 + \underbrace{\frac{2(1-\mu)}{\rho(x, y, z)_1} + \frac{2\mu}{\rho(x, y, z)_2}}_{\text{Potential Energy}} - C$$

- If $C > 2U$ then $v^2 < 0 \longrightarrow v \equiv \text{Does not exist}$

Exercise 3

DemoEx3_CR3BP_Session4.m

- Compute the zero velocity curves for a trajectory with Jacobi Constant $C=L_1$.

CR3BP: Zero velocity curves at $C_1=L_1$

1. Compute (x,y,z) for the L1 point in the Earth-Moon System.
2. Create a grid in 3D and analyse the values of the following equation:

$$v^2 = x^2 + y^2 + \frac{2(1-\mu)}{\rho(x,y,z)_1} + \frac{2\mu}{\rho(x,y,z)_2} - C_1$$

3. Plot the isosurface for $v^2=0$

CR3BP: Zero velocity curves




Figure 9.9: Zero Relative Velocity Surface Contours of the Earth-Moon System in the $x - y$ Plane
Courtesy (Shcaub & Junkins, 2002)

CR3BP: Regimes of Motion




Figure 9.10: Critical Zero Relative Velocity Surface Contours of the Earth-Moon System Touching the Lagrange Stationary Points

Courtesy (Shcaub & Junkins, 2002)



www.cranfield.ac.uk

T: +44 (0)1234 750111



@cranfielduni



@cranfielduni




/cranfielduni

Advanced Topics in Astrodynamics and Trajectory Design

Libration Point Orbits (LPOs)


Dr Joan Pau Sánchez

MSc in Astronautics and Space Engineering



www.cranfield.ac.uk

CR3BP: Five Fixed Points (Lagrange or Libration Points)




Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L1 equilibrium point?

Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L1 equilibrium point?




Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L1 equilibrium point, but with 1 cm/s error in its velocity?

Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L1 equilibrium point, but with 1 cm/s error in its velocity?




Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L5 equilibrium point?

Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L5 equilibrium point?




Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L5 equilibrium point, but with 1 cm/s error in its velocity?

Motion of a spacecraft in an equilibrium point

- What happens when we place a spacecraft at the L5 equilibrium point, but with 1 cm/s error in its velocity?




Motion of a spacecraft in an equilibrium point

- Would you say then that L5 is more stable than L1 point?
- Is L1 stable or unstable?
- Is L5 stable?

Linear Stability Analysis near fixed points

- Fixed points or Equilibrium points have an important “organizing role” of a system phase space.



- In particular, if we linearize the system in a neighbourhood of the fixed points:

$$\delta \dot{\mathbf{y}} = \mathbf{M} \delta \mathbf{y}$$

- The eigenvalues of the matrix \mathbf{M} determine the stability of the fixed point

Linear Stability Analysis near fixed points

- Consider our dynamical model: $\ddot{\mathbf{r}} = f(\mathbf{r}, \dot{\mathbf{r}})$
- Now, let us consider the motion near an equilibrium point \mathbf{r}_0 : $\ddot{\mathbf{r}}(\mathbf{r}_0, \mathbf{0}) = \mathbf{0}$

$$\ddot{\mathbf{r}}(\mathbf{r}_0 + \delta\mathbf{r}, \mathbf{0} + \delta\mathbf{v}) = \mathbf{0} + \mathbf{M} \begin{pmatrix} \delta\mathbf{r} \\ \delta\mathbf{v} \end{pmatrix}$$

$$\mathbf{M} = \left(\begin{array}{ccc|ccc} \frac{\partial \dot{x}}{\partial x} & \frac{\partial \dot{x}}{\partial y} & \frac{\partial \dot{x}}{\partial z} & \frac{\partial \dot{x}}{\partial \dot{x}} & \frac{\partial \dot{x}}{\partial \dot{y}} & \frac{\partial \dot{x}}{\partial \dot{z}} \\ \frac{\partial \dot{y}}{\partial x} & \dots & & & & \\ \vdots & & & & & \\ \hline \vdots & & & & \ddots & \vdots \\ \frac{\partial \ddot{z}}{\partial x} & \frac{\partial \ddot{z}}{\partial y} & \frac{\partial \ddot{z}}{\partial z} & \frac{\partial \ddot{z}}{\partial \dot{x}} & \frac{\partial \ddot{z}}{\partial \dot{y}} & \frac{\partial \ddot{z}}{\partial \dot{z}} \end{array} \right)$$

Linear Stability Analysis near fixed points

- Consider our dynamical model: $\ddot{\mathbf{r}} = f(\mathbf{r}, \dot{\mathbf{r}})$
- Now, let us consider the motion near an equilibrium point \mathbf{r}_0 : $\ddot{\mathbf{r}}(\mathbf{r}_0, \mathbf{0}) = \mathbf{0}$

$$\ddot{\mathbf{r}}(\mathbf{r}_0 + \delta\mathbf{r}, \mathbf{0} + \delta\mathbf{v}) = \mathbf{0} + \mathbf{M} \begin{pmatrix} \delta\mathbf{r} \\ \delta\mathbf{v} \end{pmatrix}$$

$$\mathbf{M} = \left(\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ \hline \frac{\partial \ddot{x}}{\partial x} & \frac{\partial \ddot{x}}{\partial y} & \frac{\partial \ddot{x}}{\partial z} & 0 & 2 & 0 \\ \frac{\partial \ddot{y}}{\partial x} & \frac{\partial \ddot{y}}{\partial y} & \frac{\partial \ddot{y}}{\partial z} & -2 & 0 & 0 \\ \frac{\partial \ddot{z}}{\partial x} & \frac{\partial \ddot{z}}{\partial y} & \frac{\partial \ddot{z}}{\partial z} & 0 & 0 & 0 \end{array} \right)$$

Linear Stability Analysis near fixed points

$$\left. \begin{array}{l} \dot{x} = v_x \\ \dot{y} = v_y \\ \dot{z} = v_z \\ \hline \ddot{x} = x + 2\dot{y} - \frac{1-\mu}{\rho_1^3}(x-x_1) - \frac{\mu}{\rho_2^3}(x-x_2) \\ \ddot{y} = y - 2\dot{x} - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) y \\ \ddot{z} = - \left(\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3} \right) z \end{array} \right\} \quad \rho_i = \sqrt{(x-x_i)^2 + y^2 + z^2}$$

Linear Stability Analysis near fixed points

$$x_1 = -\mu$$

$$x_2 = 1 - \mu$$

$$\rho_i = \sqrt{(x - x_i)^2 + y^2 + z^2}$$

$$\mathbf{M} = \left(\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ \hline & & \mathbf{Q} & 0 & 2 & 0 \\ & & & -2 & 0 & 0 \\ & & & 0 & 0 & 0 \end{array} \right) \quad \mathbf{Q} = \begin{pmatrix} \frac{\partial \ddot{x}}{\partial x} & \frac{\partial \ddot{x}}{\partial y} & \frac{\partial \ddot{x}}{\partial z} \\ \frac{\partial \ddot{y}}{\partial x} & \frac{\partial \ddot{y}}{\partial y} & \frac{\partial \ddot{y}}{\partial z} \\ \frac{\partial \ddot{z}}{\partial x} & \frac{\partial \ddot{z}}{\partial y} & \frac{\partial \ddot{z}}{\partial z} \end{pmatrix}$$

$$\frac{\partial \ddot{x}}{\partial x} = 1 - \frac{1-\mu}{\rho_1^3} \left(1 - \frac{3}{\rho_1^2} (x - x_1)^2 \right) - \frac{\mu}{\rho_2^3} \left(1 - \frac{3}{\rho_2^2} (x - x_2)^2 \right)$$

$$\frac{\partial \ddot{y}}{\partial y} = 1 + \frac{1-\mu}{\rho_1^3} \left(\frac{3y^2}{\rho_1^2} - 1 \right) + \frac{\mu}{\rho_2^3} \left(\frac{3y^2}{\rho_2^2} - 1 \right)$$

$$\frac{\partial \ddot{x}}{\partial y} = \frac{\partial \ddot{y}}{\partial x} = 3 \left(\frac{1-\mu}{\rho_1^5} (x - x_1) + \frac{\mu}{\rho_2^5} (x - x_2) \right) y$$

$$\frac{\partial \ddot{y}}{\partial z} = \frac{\partial \ddot{z}}{\partial y} = 3 \frac{1-\mu}{\rho_1^5} yz + 3 \frac{\mu}{\rho_2^5} yz$$

$$\frac{\partial \ddot{x}}{\partial z} = \frac{\partial \ddot{z}}{\partial x} = 3 \frac{1-\mu}{\rho_1^5} (x - x_1) z + 3 \frac{\mu}{\rho_2^5} (x - x_2) z$$

$$\frac{\partial \ddot{z}}{\partial z} = \frac{1-\mu}{\rho_1^3} \left(\frac{3z^2}{\rho_1^2} - 1 \right) + \frac{\mu}{\rho_2^3} \left(\frac{3z^2}{\rho_2^2} - 1 \right)$$

Linear Stability Analysis near fixed points

$$\begin{pmatrix} \delta \dot{\mathbf{r}} \\ \delta \dot{\mathbf{v}} \end{pmatrix} = \mathbf{M} \begin{pmatrix} \delta \mathbf{r} \\ \delta \mathbf{v} \end{pmatrix}$$

- Recall, an eigenvector of a matrix \mathbf{M} is any vector that when multiplied to \mathbf{M} , it only gets scaled.

$$\mathbf{M}\mathbf{d} = \lambda\mathbf{d} \qquad \det|\mathbf{M} - \lambda\mathbf{I}| = 0$$

- A linear ordinary differential equation such as this can then be solved as:


$$\delta \mathbf{x} = \sum_i c_i \mathbf{d}_i e^{\lambda_i t}$$

- Thus, only if λ are pure imaginary numbers the displacement will just rotate about the associated equilibrium point.

Classification of Fixed Points

$$\lambda_1 = +i\omega$$


$$\lambda_2 = -i\omega$$




Center

$$\lambda_1 < 0$$

$$\lambda_2 < 0$$



Stable node (sink)




Stable spiral

$$\lambda_1 = -\alpha + i\beta$$


$$\lambda_2 = -\alpha - i\beta$$

$$\lambda_1 < 0$$


$$\lambda_2 > 0$$



Saddle point



Unstable node (source)



Unstable spiral

$$\lambda_1 = \alpha + i\beta$$

$$\lambda_2 = \alpha - i\beta$$

$$\lambda_1 > 0$$

$$\lambda_2 > 0$$

Periodic Orbits near L1 and L2

- Let us prove the existence of periodic motion near the L1/L2 points.

Periodic Orbits near L1 and L2

$$(\mathbf{x}_{Li}, \dot{\mathbf{x}}_{Li}) = (x_{Li}, 0, 0, 0, 0, 0)$$

$$x_1 = -\mu$$

$$x_2 = 1 - \mu$$

$$\rho_i = \sqrt{(x - x_i)^2}$$

$$\Omega = +\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3}$$

$$\mathbf{M} = \left(\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ \hline 1+2\Omega & 0 & 0 & 0 & 2 & 0 \\ 0 & 1-\Omega & 0 & -2 & 0 & 0 \\ 0 & 0 & -\Omega & 0 & 0 & 0 \end{array} \right)$$

$$\begin{pmatrix} \dot{x}' \\ \dot{y}' \\ \dot{z}' \\ \ddot{x}' \\ \ddot{y}' \\ \ddot{z}' \end{pmatrix} = \mathbf{M} \begin{pmatrix} x' \\ y' \\ z' \\ \dot{x}' \\ \dot{y}' \\ \dot{z}' \end{pmatrix}$$

$$\ddot{x}' - 2\dot{y}' - (1 + 2\Omega)x' = 0$$

$$\ddot{y}' + 2\dot{x}' + (\Omega + 1)y' = 0$$

$$\ddot{z}' + \Omega z' = 0$$

Periodic Orbits near L1 and L2

- Let us prove the existence of periodic motion near the L1/L2 points

$$\ddot{x}' - 2\dot{y}' - (1 + 2\Omega)x' = 0$$

$$\ddot{y}' + 2\dot{x}' + (\Omega + 1)y' = 0$$

$$\ddot{z}' + \Omega z' = 0$$

- In the linearized approximation, the out-of-plane component has no influence on the in-plane motion.

$$\ddot{z}' + \Omega z' = 0 \longrightarrow z' = A_z \sin(\sqrt{\Omega}t + \varphi)$$

- While in the X-Y plane, arbitrarily chosen initial conditions will give rise to unbounded motion. However, periodic motion is also possible if starting conditions are chosen suitably.

Periodic Orbits near L1 and L2

- Let us prove the existence of periodic motion near the L1/L2 points

$$\ddot{x}' - 2\dot{y}' - (1 + 2\Omega)x' = 0$$

$$\ddot{y}' + 2\dot{x}' + (\Omega + 1)y' = 0$$

$$\begin{aligned} x' &= -A_x \cos(\omega_p t + \phi) \\ y' &= kA_x \sin(\omega_p t + \phi) \end{aligned} \quad \begin{aligned} k &= \frac{\omega_p^2 + 1 + 2\Omega}{2\omega_p} \\ \omega_p^2 &= \frac{2 - \Omega + \sqrt{9\Omega^2 - 8\Omega}}{2} \end{aligned}$$

Periodic Orbits near L1 and L2

$$\ddot{x}' - 2\dot{y}' - (1 + 2\Omega)x' = 0$$

$$\ddot{y}' + 2\dot{x}' + (\Omega + 1)y' = 0$$

$$\begin{array}{lll} x' = -A_x \cos(\omega_p t + \phi) & \dot{x}' = A_x \omega_p \sin(\omega_p t + \phi) & \ddot{x}' = A_x \omega_p^2 \cos(\omega_p t + \phi) \\ y' = k A_x \sin(\omega_p t + \phi) & \dot{y}' = k \omega_p A_x \cos(\omega_p t + \phi) & \ddot{y}' = -k \omega_p^2 A_x \sin(\omega_p t + \phi) \end{array}$$

$$A_x \omega_p^2 \cos(\omega_p t + \phi) = (2k \omega_p - 2\Omega - 1) A_x \cos(\omega_p t + \phi)$$

$$\omega_p^2 = 2k \omega_p - 2\Omega - 1$$

$$k = \frac{\omega_p^2 + 1 + 2\Omega}{2\omega_p}$$

Periodic Orbits near L1 and L2

$$\ddot{x}' - 2\dot{y}' - (1 + 2\Omega)x' = 0$$

$$\ddot{y}' + 2\dot{x}' + (\Omega + 1)y' = 0$$

$$\begin{aligned} x' &= -A_x \cos(\omega_p t + \phi) & \dot{x}' &= A_x \omega_p \sin(\omega_p t + \phi) & \ddot{x}' &= A_x \omega_p^2 \cos(\omega_p t + \phi) \\ y' &= k A_x \sin(\omega_p t + \phi) & \dot{y}' &= k \omega_p A_x \cos(\omega_p t + \phi) & \ddot{y}' &= -k \omega_p^2 A_x \sin(\omega_p t + \phi) \end{aligned}$$

$$-k \omega_p^2 A_x \sin(\omega_p t + \phi) = -(2\omega_p + (\Omega + 1)k) A_x \sin(\omega_p t + \phi)$$

$$-k \omega_p^2 = -2\omega_p - (\Omega + 1)k \qquad k = \frac{\omega_p^2 + 1 + 2\Omega}{2\omega_p}$$


$$\omega_p^2 = \frac{2 - \Omega + \sqrt{9\Omega^2 - 8\Omega}}{2}$$

Periodic Orbits near L1 and L2


$$x' = -A_x \cos(\omega_p t + \phi)$$

$$y' = kA_x \sin(\omega_p t + \phi)$$

$$z' = A_z \sin(\sqrt{\Omega} t + \phi)$$




Periodic Orbits near L1 and L2




$$x' = -A_x \cos(\omega_p t + \phi)$$

$$y' = kA_x \sin(\omega_p t + \phi)$$

$$z' = A_z \sin(\sqrt{\Omega} t + \varphi)$$



Lissajous Orbits



Exercise 2

Compute a planar Lyapunov Orbit for the Earth-Moon L1 point with an A_x of 1000km.

Periodic Orbits near L1 and L2

- Let us start computing a Lyapunov orbit by propagating the initial conditions as given by equations:

$$\begin{aligned}x' &= -A_x \cos(\omega_p t + \phi) & \dot{x}' &= A_x \omega_p \sin(\omega_p t + \phi) \\y' &= k A_x \sin(\omega_p t + \phi) & \dot{y}' &= k \omega_p A_x \cos(\omega_p t + \phi)\end{aligned}$$

$$\mathbf{x}_0 = \begin{pmatrix} -A_x & 0 & 0 & 0 & k \omega_p A_x & 0 \end{pmatrix}$$

$$t = 0$$

$$\phi = 0$$


$$\omega_p = \sqrt{\frac{2 - \Omega + \sqrt{9\Omega^2 - 8\Omega}}{2}}$$

$$k = \frac{\omega_p^2 + 1 + 2\Omega}{2\omega_p}$$


$$\Omega = +\frac{1-\mu}{\rho_1^3} + \frac{\mu}{\rho_2^3}$$

Periodic Orbits near L1 and L2

- Let us start computing a Lyapunov orbit by propagating the initial conditions.



Periodic Orbits near L1 and L2



Transition Matrix:

$$\delta \mathbf{x} = \Phi \left(\frac{T}{2}, 0 \right) \delta \mathbf{x}_0$$

$$\dot{\Phi}(t, t_0) = \mathbf{M}(t) \Phi(t, t_0)$$


$$\Phi(t_0, t_0) = \mathbf{I}$$

$$\begin{pmatrix} \Delta x_f \\ \Delta y_f \\ \Delta z_f \\ \Delta v_{xf} \\ \Delta v_{yf} \\ \Delta v_{zf} \end{pmatrix} = \Phi_{6 \times 6} \left(\frac{T}{2}, 0 \right) \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \Delta v_{y0} \\ 0 \end{pmatrix}$$


$$\Delta v_{xf} = \Phi_{(4 \ 5)} \Delta v_{y0} \longrightarrow \Phi_{(4 \ 5)} \Delta v_{y0} = -v_{xf} \longrightarrow \Phi_{(4 \ 5)} \Delta v_{y0} = \frac{-v_{xf}}{\Phi_{(4 \ 5)}}$$

Periodic Orbits near L1 and L2



Lyapunov Orbits (Planar & Vertical)



Halo Orbits




ISEE-3 / ICE





Solar-Rotating Coordinates, Ecliptic Plane Projection

| | |
|-------------------------|--|
| Mission: | Investigate Solar-Terrestrial relationships, Solar Wind, Magnetosphere, and Cosmic Rays |
| Launch: | Sept., 1978, Comet Encounter Sept., 1985 |
| Lissajous Orbit: | L1 Libration Halo Orbit, $A_x \sim 175,000\text{km}$, $A_y = 660,000\text{km}$, $A_z \sim 120,000\text{km}$, Class I |
| Spacecraft: | Mass=480Kg, Spin stabilized, |
| Notable: | First Ever Libration Orbiter, First Ever Comet Encounter |

ISEE-3 / ICE



WIND



Mission: Investigate Solar-Terrestrial relationships, Solar Wind, Magnetosphere



Launch: Nov., 1994, Multiple Lunar Gravity Assist

Lissajous Orbit: Originally an L1 Lissajous Constrained Orbit, $A_x \sim 10,000\text{km}$, $A_y \sim 350,000\text{km}$, $A_z \sim 250,000\text{km}$, Class I

Spacecraft: Mass=1254Kg, Spin stabilized,



Notable: First Ever Multiple Gravity Assist towards L1

ACE



| | |
|-------------------------|--|
| Mission: | Investigate low-energy solar and high-energy galactic particles and Solar Wind |
| Launch: | August, 1997, Direct Transfer |
| Lissajous Orbit: | L1 Lissajous Constrained Orbit, $A_y \sim <264,000$, $A_x \sim \text{tbd km}$, $A_z \sim 157,000\text{km}$, |
| Class I | |
| Spacecraft: | Mass=757Kg, Three Axis Stabilized, 'Solar' Pointing |
| Notable: | First Constrained Transfer Orbit |

James Webb Space Telescope



Mission: JWST Is Part of Origins Program. Designed to Be the Successor to the Hubble Space Telescope. NGST Observations in the Infrared Part of the Spectrum.


Launch: ~2011, Direct Transfer

Lissajous Orbit: L2 large lissajous, $A_y \sim 294,000\text{km}$, $A_x \sim 800,000\text{km}$, $A_z \sim 131,000\text{km}$, Class I or II

Spacecraft: Mass~6000kg, Three Axis Stabilized, 'Star' Pointing

Notable: Observations in the Infrared Part of the Spectrum. Important That the Telescope Be Kept at Low Temperatures, $\sim 3^0\text{K}$. Large Solar Shade/Solar Sail

Periodic Orbits near L1 and L2





www.cranfield.ac.uk

T: +44 (0)1234 750111



@cranfielduni



@cranfielduni



/cranfielduni