

Mathematics Methods for Computer Science

Instructor: Xubo Yang

SJTU-SE DALAB

Lecture

Eigenproblems I

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Given: Collection of data points \vec{x}_i

- Age
- Weight
- Blood pressure
- Heart rate

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Given: Collection of data points \vec{x}_i

- Age
- Weight
- Blood pressure
- Heart rate

Find: Correlations between different dimensions

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

One-dimensional subspace

$$\vec{x}_i \approx c_i \vec{v}, \vec{v} \text{ unknown}$$

One-dimensional subspace

$$\vec{x}_i \approx c_i \vec{v}, \vec{v} \text{ unknown}$$

Equivalently:

$$\vec{x}_i \approx c_i \hat{v}$$

$$\hat{v} \text{ unknown with } \|\hat{v}\|_2 = 1$$

$$\begin{aligned} & \text{minimize}_{\hat{v}} \sum_i \|\vec{x}_i - \text{proj}_{\hat{v}} \vec{x}_i\|_2^2 \\ & \text{such that } \|\hat{v}\|_2 = 1 \end{aligned}$$

$$\begin{aligned} & \text{minimize}_{\hat{v}} \sum_i \|\vec{x}_i - \text{proj}_{\hat{v}} \vec{x}_i\|_2^2 \\ & \text{such that } \|\hat{v}\|_2 = 1 \end{aligned}$$

What does the constraint do?

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\begin{aligned} & \text{minimize}_{\hat{v}} \sum_i \|\vec{x}_i - \text{proj}_{\hat{v}} \vec{x}_i\|_2^2 \\ & \text{such that } \|\hat{v}\|_2 = 1 \end{aligned}$$

What does the constraint do?

- Does not affect optimal \hat{v}
- Removes scaling ambiguity

Statistical Motivation


Properties

Spectral Theorem


Other

ODE Theory


Spectral Embedding



(a) Input data



(b) Principal axis



(c) Projection error

$$\min_{c_i} ||\vec{x}_i - c_i \hat{v}||_2$$

What is c_i ?

$$\min_{c_i} ||\vec{x}_i - c_i \hat{v}||_2$$

What is c_i ?

$$c_i = \vec{x}_i \cdot \hat{v}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\begin{aligned} &\text{maximize } ||X^T \hat{v}||_2^2 \\ &\text{such that } ||\hat{v}||_2^2 = 1 \end{aligned}$$

Eigenvector of XX^T with largest
eigenvalue.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Eigenvector of XX^T with largest
eigenvalue.

”First principal component”
More after SVD!

Eigenvalue and eigenvector

An eigenvector $\vec{x} \neq \vec{0}$ of $A \in \mathbb{R}^{n \times n}$ satisfies $A\vec{x} = \lambda\vec{x}$ for some $\lambda \in \mathbb{R}$; λ is an eigenvalue.

Complex eigenvalues and eigenvectors instead have $\lambda \in \mathbb{C}$ and $\vec{x} \in \mathbb{C}^n$.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Eigenvalue and eigenvector

An eigenvector $\vec{x} \neq \vec{0}$ of $A \in \mathbb{R}^{n \times n}$ satisfies

$A\vec{x} = \lambda\vec{x}$ for some $\lambda \in \mathbb{R}$; λ is an eigenvalue.

Complex eigenvalues and eigenvectors instead have $\lambda \in \mathbb{C}$ and $\vec{x} \in \mathbb{C}^n$.

Scale doesn't matter!

→ can constrain $||\vec{x}||_2 \equiv 1$

- Optimize $||A\vec{x}||_2$ such that $||\vec{x}||_2 = 1$
(important!)

- Optimize $||A\vec{x}||_2$ such that $||\vec{x}||_2 = 1$ (important!)
- ODE/PDE problems: Closed solutions and approximations for $\vec{y}' = B\vec{y}$

- Optimize $||A\vec{x}||_2$ such that $||\vec{x}||_2 = 1$
(important!)
- ODE/PDE problems: Closed solutions and approximations for $\vec{y}' = B\vec{y}$
- Critical points of Rayleigh quotient:
$$\frac{\vec{x}^T A \vec{x}}{||\vec{x}||_2^2}$$

Two Basic Properties

Proved in textbook

Lemma

Every matrix $A \in \mathbb{R}^{n \times n}$ has at least one (complex) eigenvector.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Two Basic Properties

Proved in textbook

Lemma

Every matrix $A \in \mathbb{R}^{n \times n}$ has at least one (complex) eigenvector.

Lemma

Eigenvectors corresponding to distinct eigenvalues must be linearly independent.

Two Basic Properties

Proved in textbook

Lemma

Every matrix $A \in \mathbb{R}^{n \times n}$ has at least one (complex) eigenvector.

Lemma

Eigenvectors corresponding to distinct eigenvalues must be linearly independent.

→ at most n eigenvalues

Nondefective

$A \in \mathbb{R}^{n \times n}$ is nondefective or diagonalizable if its eigenvectors span \mathbb{R}^n .

Nondefective

$A \in \mathbb{R}^{n \times n}$ is nondefective or diagonalizable if its eigenvectors span \mathbb{R}^n .

$$D = X^{-1}AX$$

A is diagonalized by a similarity transformation $A \rightarrow X^{-1}AX$

Spectrum and spectral radius

The spectrum of A is the set of eigenvalues of A .

The spectral radius $\rho(A)$ is the eigenvalue λ maximizing $|\lambda|$.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Complex conjugate

The complex conjugate of a number

$z = a + bi \in \mathbb{C}$ is $\bar{z} \equiv a - bi$.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Complex conjugate

The complex conjugate of a number

$$z = a + bi \in \mathbb{C} \text{ is } \bar{z} \equiv a - bi.$$

Complex transpose

The conjugate transpose of $A \in \mathbb{C}^{m \times n}$ is

$$A^H \equiv \overline{A}^T.$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$A = A^H$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Lemma

All eigenvalues of Hermitian matrices are real.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Lemma

All eigenvalues of Hermitian matrices are real.

Lemma

Eigenvectors corresponding to distinct eigenvalues of Hermitian matrices must be orthogonal.

Spectral Theorem

Suppose $A \in \mathbb{C}^{n \times n}$ is Hermitian (if $A \in \mathbb{R}^{n \times n}$, suppose it is symmetric). Then, A has exactly n orthonormal eigenvectors $\vec{x}_1, \dots, \vec{x}_n$ with (possibly repeated) eigenvalues $\lambda_1, \dots, \lambda_n$.

Spectral Theorem

Suppose $A \in \mathbb{C}^{n \times n}$ is Hermitian (if $A \in \mathbb{R}^{n \times n}$, suppose it is symmetric). Then, A has exactly n orthonormal eigenvectors $\vec{x}_1, \dots, \vec{x}_n$ with (possibly repeated) eigenvalues $\lambda_1, \dots, \lambda_n$.

$$\text{Full set: } D = X^T A X$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\vec{b} = c_1 \vec{x}_1 + \cdots + c_k \vec{x}_k$$
$$A\vec{x} = \vec{b}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\vec{b} = c_1 \vec{x}_1 + \cdots + c_k \vec{x}_k$$
$$A\vec{x} = \vec{b}$$

$$\Rightarrow \vec{x} = \frac{c_1}{\lambda_1} \vec{x}_1 + \cdots + \frac{c_n}{\lambda_n} \vec{x}_n$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\vec{b} = c_1 \vec{x}_1 + \cdots + c_k \vec{x}_k$$
$$A\vec{x} = \vec{b}$$

$$\Rightarrow \vec{x} = \frac{c_1}{\lambda_1} \vec{x}_1 + \cdots + \frac{c_n}{\lambda_n} \vec{x}_n$$

$$A = XDX^{-1} \Rightarrow A^{-1} = XD^{-1}X^{-1}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

- Given symmetric positive semi-definite (PSD) matrix, U
- Can compute matrix square root, $U^{1/2}$

- Given real $n - by - n$ matrix, A
- There exists a unique factorization called the Polar Decomposition

$$A = RU$$

where R is an $n - by - n$ orthogonal matrix, and U is an $n - by - n$ symmetric PSD right "stretch" matrix.

- Also a left stretch matrix, W , such that $A = WR$.
- Geometric interpretation.

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding



- Fast Lattice Shape Matching (Fast LSM)
- SIGGRAPH 2007 [Rivers and James 2007]
- <http://www.alecrivers.com/fastlsm>
- Need to compute orientation, R , of local particle groups
- Millions of polar decompositions (and eigenvalue decomp) per second

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Newton:

$$\vec{F} = m \frac{d^2 \vec{x}}{dt^2}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Newton:

$$\vec{F} = m \frac{d^2 \vec{x}}{dt^2}$$

Hooke:

$$\vec{F}_s = k(\vec{x} - \vec{y})$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$M\vec{X}'' = K\vec{X}$$
$$\longrightarrow \frac{d}{dt} \begin{pmatrix} \vec{X} \\ \vec{V} \end{pmatrix} = \begin{pmatrix} 0 & I_{3n \times 3n} \\ M^{-1}K & 0 \end{pmatrix} \begin{pmatrix} \vec{X} \\ \vec{V} \end{pmatrix}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\vec{Y}' = B\vec{Y}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\begin{aligned}\vec{y}' &= B\vec{y} \\ B\vec{y}_i &= \lambda_i\vec{y}_i \\ \vec{y}(0) &= c_1\vec{y}_1 + \cdots + c_k\vec{y}_k\end{aligned}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\vec{y}' = B\vec{y}$$

$$B\vec{y}_i = \lambda_i\vec{y}_i$$

$$\vec{y}(0) = c_1\vec{y}_1 + \cdots + c_k\vec{y}_k$$

$$\rightarrow \vec{y}(t) = c_1e^{\lambda_1 t}\vec{y}_1 + \cdots + c_ke^{\lambda_k t}\vec{y}_k$$

Application: Modal Sound Synthesis

Statistical Motivation


Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding



Major role in physics-based sound synthesis

<https://www.youtube.com/watch?v=dMUHp8i6E5E>

Organizing a Collection

Statistical Motivation


Properties

Spectral Theorem


Other

ODE Theory

Spectral Embedding



(a) Database of photos



(b) Spectral embedding

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Have: n items in a dataset

$w_{ij} \geq 0$ similarity of items i and j

$$w_{ij} = w_{ji}$$

Want: x_i embedding on \mathbb{R}

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$E(\vec{x}) = \sum_{ij} w_{ij} (x_i - x_j)^2$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\text{minimize } E(\vec{x})$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\begin{aligned} &\text{minimize } E(\vec{x}) \\ &\text{such that } ||\vec{x}||_2^2 = 1 \end{aligned}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$\begin{aligned} &\text{minimize } E(\vec{x}) \\ &\text{such that } \|\vec{x}\|_2^2 = 1 \\ &\quad \vec{1}\vec{x} = 0 \end{aligned}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

$$E(\vec{x}) = 2\vec{x}^T(A - W)\vec{x}$$

Statistical Motivation

Properties

Spectral Theorem

Other

ODE Theory

Spectral Embedding

Eigenvector of $A - W$ with
second smallest eigenvalue.