

Table 2
Rotated Factor Loadings for Grief Reaction Measures (N = 250)

	Physiological	Behavioral	Cognitive	Attitudes toward Deceased	Affective	h^{2a}
Loss of sleep	.02	.27	<u>.69</u>	.20	.03	.61
Oversleep	<u>.72</u>	.02	.02	.02	.02	.52
Loss of appetite	<u>.44</u>	.03	.37	<u>.50</u>	.03	.58
Loss of interest	<u>.52</u>	.14	.25	.03	.13	.38
Feelings of numbness	<u>.52</u>	.02	.39	.12	.02	.44
Lack of concentration	.27	.20	<u>.56</u>	.03	.32	.52
Inability to control emotions in public	.11	<u>.77</u>	.02	.02	.03	.62
Inability to control emotions in private	.03	<u>.71</u>	<u>.41</u>	.03	.03	.68
Anger	.03	.02	.35	.02	<u>.61</u>	.51
Guilt	.03	.03	.02	.12	<u>.79</u>	.65
Frustration	.39	.02	<u>.47</u>	.02	<u>.42</u>	.55
Thinking at times the dead person was still alive	.17	.13	-.16	<u>.78</u>	.13	.70
Feeling the presence of dead person	-.16	.03	.19	<u>.75</u>	.03	.63
Dwelling on events leading up to the death	.35	<u>.47</u>	-.32	.03	.36	.58
Feeling that life had little or no meaning	<u>.66</u>	.12	.21	.19	.13	.55
Crying	.03	<u>.72</u>	.25	.16	.18	.64
Percentage of variance	13.12	12.70	11.96	9.87	9.54	

Note. Reproduced with permission of the T. Bell (1981) Bereavement Study. Underlined coefficients indicate variables used to provide substantive meaning to rotated factors.

^aCommunality estimates for 5-factor model.