

# On the deletion channel with small deletion probability

Yashodhan Kanoria

Stanford University

joint work with Andrea Montanari

# Outline

- 1 Introduction
- 2 Main results
- 3 Proof Sketch

# Outline

- 1 Introduction
- 2 Main results
- 3 Proof Sketch

# The deletion channel

- $n$  input bits  $X^n$ , each bit deleted independently w.p.  $d$
- Output  $Y(X^n)$  of length  $\text{Binomial}(n, 1 - d)$


# The deletion channel

- $n$  input bits  $X^n$ , each bit deleted independently w.p.  $d$
- Output  $Y(X^n)$  of length  $\text{Binomial}(n, 1 - d)$

$X^n$ : 1 1 1 0 1 1 0 0 0 1 0 ...

# The deletion channel

- $n$  input bits  $X^n$ , each bit deleted independently w.p.  $d$
- Output  $Y(X^n)$  of length  $\text{Binomial}(n, 1 - d)$



# Capacity of the deletion channel

$$C_n = \frac{1}{n} \max_{p_{X^n}} I(X^n; Y(X^n))$$

Lemma (Dobrushin '67)

$\lim_{n \rightarrow \infty} C_n$  exists and is equal to  $\inf_{n \geq 1} C_n$ .

Define capacity  $C$  as this limit.

# Capacity of the deletion channel

$$C_n = \frac{1}{n} \max_{p_{X^n}} I(X^n; Y(X^n))$$

Lemma (Dobrushin '67)

$\lim_{n \rightarrow \infty} C_n$  exists and is equal to  $\inf_{n \geq 1} C_n$ .

**Define capacity  $C$  as this limit.**



Problems:

- Don't know how to compute capacity
- Optimal  $p_{X^n}$  not known for any  $d > 0$

What's so hard?

LACK OF SYNCHRONIZATION!

Problems:

- Don't know how to compute capacity
- Optimal  $p_{X^n}$  not known for any  $d > 0$

What's so hard?

LACK OF SYNCHRONIZATION!


Problems:

- Don't know how to compute capacity
- Optimal  $p_{X^n}$  not known for any  $d > 0$

What's so hard?

**LACK OF SYNCHRONIZATION!**

# Bounds on Capacity



- $(1-d)/9 \leq C \leq (1-d)$  [Mitzenmacher et al '06]
- Upper bounds: augmented channels [Diggavi et al '07, Fertoni-Duman '09].
- Best computed lower bds: Markov sources + Jigsaw decoding [Mitzenmacher-Drinea '07]

# New Idea

# New Idea

**I know how to solve  $d = 0$ !**

# New Idea

**I know how to solve  $d = 0$ !**

Can we expand  $C$  for  $d \rightarrow 0^+$ ?

Optimal input distribution for small  $d$ ?

- [Kalai, Mitzenmacher & Sudan, ISIT '10] addresses **same problem!**
- Shows  $C = 1 - d \log(1/d) + o(d \log(1/d))$
- Very different proof technique

We obtain in addition:

- (i) Order  $d$  term (in paper)
- (ii) Order  $d^2$  term and optimal coding scheme (updated result)



- [Kalai, Mitzenmacher & Sudan, ISIT '10] addresses **same problem!**
- Shows  $C = 1 - d \log(1/d) + o(d \log(1/d))$
- Very different proof technique

We obtain in addition:

- (i) Order  $d$  term (in paper)
- (ii) Order  $d^2$  term and optimal coding scheme (updated result)

- [Kalai, Mitzenmacher & Sudan, ISIT '10] addresses **same problem!**
- Shows  $C = 1 - d \log(1/d) + o(d \log(1/d))$
- Very different proof technique

We obtain in addition:

- (i) Order  $d$  term (in paper)
- (ii) Order  $d^2$  term and optimal coding scheme (updated result)

# Outline

- 1 Introduction
- 2 Main results
- 3 Proof Sketch

# Main result: Capacity expansion

Result in our ISIT paper:

## Theorem

For small  $d$ ,

$$\textcircled{1} \quad C(d) = 1 - d \log(1/d) - A_1 d + O(d^{1.4})$$

where  $A_1 = \log(2e) - \sum_{\ell=1}^{\infty} 2^{-\ell-1} \ell \log_2 \ell \approx 1.154$

$\textcircled{2}$  The iid Bernoulli( $1/2$ ) process achieves rate  $C - O(d^{1.4})$ .

# Main result: Capacity expansion

Result in our ISIT paper:

## Theorem

For small  $d$ ,

$$\textcircled{1} \quad C(d) = 1 - d \log(1/d) - A_1 d + O(d^{1.4})$$

where  $A_1 = \log(2e) - \sum_{\ell=1}^{\infty} 2^{-\ell-1} \ell \log_2 \ell \approx 1.154$

$$\textcircled{2} \quad \text{The iid Bernoulli}(1/2) \text{ process achieves rate } C - O(d^{1.4}).$$

**But the expansion can go further!**

# Main result: Capacity expansion

## Updated result:

### Theorem

For small  $d$ ,

$$\textcircled{1} \quad C(d) = 1 - d \log(1/d) - A_1 d + A_2 d^2 + O(d^{2.9})$$

where  $A_1 = \log(2e) - \sum_{\ell=1}^{\infty} 2^{-\ell-1} \ell \log_2 \ell \approx 1.154$

$A_2 = \dots$  (multi-line expression)  $\approx 1.792$

$\textcircled{2}$  The process  $X^*$  (coming up!) achieves  $C - O(d^{2.9})$ .

# Implications of expansion

- Previous best upper bound off by  $(1/4)d \log(1/d)$
- Previous computed lower bound off by  $0.904d^2$ :  
 Bounds based on Markov sources + Jigsaw decoding  
 [Diggavi et al '01, Mitzenmacher-Drinea '07]

## Fact

- *The maximum rate achieved by a first order Markov source is*

$$R_{\text{Mkv}} = C - 0.100d^2 + O(d^{2.9}).$$

- *'Jigsaw decoding' incurs asymptotic rate loss of  $0.804d^2 + O(d^{2.9})$ .*

Thus, previous computed lower bounds off by  $0.904d^2$  asymptotically.

# Implications of expansion

- Previous best upper bound off by  $(1/4)d \log(1/d)$
- Previous computed lower bound off by  $0.904d^2$ :  
Bounds based on Markov sources + Jigsaw decoding  
[Diggavi et al '01, Mitzenmacher-Drinea '07]

## Fact

- *The maximum rate achieved by a first order Markov source is*

$$R_{\text{Mkv}} = C - 0.100d^2 + O(d^{2.9}).$$

- *'Jigsaw decoding' incurs asymptotic rate loss of  $0.804d^2 + O(d^{2.9})$ .*

Thus, previous computed lower bounds off by  $0.904d^2$  asymptotically.



# Implications of expansion

- Previous best upper bound off by  $(1/4)d \log(1/d)$
- Previous computed lower bound off by  $0.904d^2$ :  
Bounds based on Markov sources + Jigsaw decoding  
[Diggavi et al '01, Mitzenmacher-Drinea '07]

## Fact

- *The maximum rate achieved by a first order Markov source is*

$$R_{\text{Mkv}} = C - 0.100d^2 + O(d^{2.9}).$$

- *'Jigsaw decoding' incurs asymptotic rate loss of  $0.804d^2 + O(d^{2.9})$ .*

Thus, previous computed lower bounds off by  $0.904d^2$  asymptotically.

# Implications of expansion

- Previous best upper bound off by  $(1/4)d \log(1/d)$
- Previous computed lower bound off by  $0.904d^2$ :  
Bounds based on Markov sources + Jigsaw decoding  
[Diggavi et al '01, Mitzenmacher-Drinea '07]

## Fact


- *The maximum rate achieved by a first order Markov source is*

$$R_{\text{Mkv}} = C - 0.100d^2 + O(d^{2.9}).$$

- *'Jigsaw decoding' incurs asymptotic rate loss of  $0.804d^2 + O(d^{2.9})$ .*

Thus, **previous computed lower bounds off by  $0.904d^2$  asymptotically.**

# Capacity Expansion



# Capacity Expansion

$d$	LB	$C_{\text{exp}}$ upto $d^2$	$C_{\text{exp}}$ upto $d$	UB
0.05	0.7283	0.7307	0.7262	0.8160
0.10	0.5620	0.5703	0.5524	0.6890
0.15	0.4392	0.4566	0.4163	0.5790

# Optimal coding

- 'Runs' of 0s and 1s

$\dots$  0 1 1 1 0 1 1 0 0 0 1 0  $\dots$   
 $\mathcal{R}_1$   $\mathcal{R}_2$   $\mathcal{R}_3$   $\mathcal{R}_4$   $\mathcal{R}_5$

- $L \equiv$  Length of randomly selected run in stationary  $\mathbb{X}$
- $\mathbb{X}$  is iid Bernoulli(1/2):  $L \sim \text{Geo}(1/2)$ , i.e.  $p_L(\ell) = 2^{-\ell}$ .

# Optimal coding

- 'Runs' of 0s and 1s

$\mathcal{R}_1$     $\mathcal{R}_2$     $\mathcal{R}_3$     $\mathcal{R}_4$     $\mathcal{R}_5$   
 ... 0 1 1 1 0 1 1 0 0 0 1 0 ...

- $L \equiv$  Length of randomly selected run in stationary  $\mathbb{X}$
- $\mathbb{X}$  is iid Bernoulli(1/2):  $L \sim \text{Geo}(1/2)$ , i.e.  $p_L(\ell) = 2^{-\ell}$ .

# Optimal coding

- 'Runs' of 0s and 1s

$\mathcal{R}_1$     $\mathcal{R}_2$     $\mathcal{R}_3$     $\mathcal{R}_4$     $\mathcal{R}_5$   
 $\dots 0 \ 1 \ 1 \ 1 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 1 \ 0 \ \dots$

- $L \equiv$  Length of randomly selected run in stationary  $\mathbb{X}$
- $\mathbb{X}$  is iid Bernoulli(1/2):  $L \sim \text{Geo}(1/2)$  , i.e.  $p_L(\ell) = 2^{-\ell}$ .

# Optimal coding

Input distribution to achieve  $d^2$  term:

## Theorem

The stationary process  $\mathbb{X}^*$  consisting of *iid runs* with distribution

$$p_L^*(\ell) = 2^{-\ell}(1 + d(\ell \ln \ell - c\ell))$$

(where  $c = \sum_{\ell=1}^{\infty} 2^{-\ell-1} \ell \ln \ell \approx 0.893$ .)

achieves rate  $R_* = C - O(d^{2.9})$ .

First optimal coding result for deletion channel.



# Optimal coding

Input distribution to achieve  $d^2$  term:

## Theorem

The stationary process  $\mathbb{X}^*$  consisting of *iid runs* with distribution

$$p_L^*(\ell) = 2^{-\ell}(1 + d(\ell \ln \ell - c\ell))$$

(where  $c = \sum_{\ell=1}^{\infty} 2^{-\ell-1} \ell \ln \ell \approx 0.893$ .)

achieves rate  $R_* = C - O(d^{2.9})$ .

**First optimal coding result** for deletion channel.

# Outline

- 1 Introduction
- 2 Main results
- 3 Proof Sketch**

## Theorem

*For small  $d$  and any  $\epsilon > 0$ ,*

$$C(d) = 1 - d \log(1/d) - 1.154 d + O(d^{1.4})$$

*and the iid Bernoulli(1/2) process achieves rate  $C - O(d^{1.4})$ .*

# Preliminaries

## Lemma

*Stationary ergodic sources suffice to achieve  $C$ .*

# Preliminaries

$$I(X^n; Y(X^n)) = H(Y) - H(Y|X^n)$$

Let  $D^n \equiv$  channel realization.

$$\begin{aligned} H(Y|X^n) &= H(D^n, Y|X^n) - H(D^n|X^n, Y) \\ &= nh(d) - H(D^n|X^n, Y) \end{aligned}$$

since  $Y = f(X^n, D^n)$ , and  $D^n$  is iid Bernoulli( $d$ ) independent of  $X^n$ .

Main problem:  $H(D^n|X^n, Y)$ .

# Preliminaries

$$I(X^n; Y(X^n)) = H(Y) - H(Y|X^n)$$

Let  $D^n \equiv$  channel realization.

$$\begin{aligned} H(Y|X^n) &= H(D^n, Y|X^n) - H(D^n|X^n, Y) \\ &= nh(d) - H(D^n|X^n, Y) \end{aligned}$$

since  $Y = f(X^n, D^n)$ , and  $D^n$  is iid Bernoulli( $d$ ) independent of  $X^n$ .

**Main problem:**  $H(D^n|X^n, Y)$ .

# Preliminaries

$$I(X^n; Y(X^n)) = H(Y) - H(Y|X^n)$$

Let  $D^n \equiv$  channel realization.

$$\begin{aligned} H(Y|X^n) &= H(D^n, Y|X^n) - H(D^n|X^n, Y) \\ &= nh(d) - H(D^n|X^n, Y) \end{aligned}$$

since  $Y = f(X^n, D^n)$ , and  $D^n$  is iid Bernoulli( $d$ ) independent of  $X^n$ .

Main problem:  $H(D^n|X^n, Y)$ .

# Preliminaries

$$I(X^n; Y(X^n)) = H(Y) - H(Y|X^n)$$

Let  $D^n \equiv$  channel realization.

$$\begin{aligned} H(Y|X^n) &= H(D^n, Y|X^n) - H(D^n|X^n, Y) \\ &= nh(d) - H(D^n|X^n, Y) \end{aligned}$$

since  $Y = f(X^n, D^n)$ , and  $D^n$  is iid Bernoulli( $d$ ) independent of  $X^n$ .

**Main problem:**  $H(D^n|X^n, Y)$ .



Many possibilities for  $D^n$  given  $(X^n, Y(X^n))$ , e.g.

⋮

What is conditional entropy rate  $\lim_{n \rightarrow \infty} H(D^n | X^n, Y)/n$ ?

Many possibilities for  $D^n$  given  $(X^n, Y(X^n))$ , e.g.


$X^n$ : 1 1 1 0 1 1 0 0 0 1 0 ...

$Y(X^n)$ : 1 1 1 1 0 1 0 ...

⋮

What is conditional entropy rate  $\lim_{n \rightarrow \infty} H(D^n | X^n, Y)/n$ ?



Many possibilities for  $D^n$  given  $(X^n, Y(X^n))$ , e.g.



...

What is conditional entropy rate  $\lim_{n \rightarrow \infty} H(D^n | X^n, Y)/n$ ?



Many possibilities for  $D^n$  given  $(X^n, Y(X^n))$ , e.g.



...

What is conditional entropy rate  $\lim_{n \rightarrow \infty} H(D^n | X^n, Y) / n$ ?

Many possibilities for  $D^n$  given  $(X^n, Y(X^n))$ , e.g.



⋮

What is conditional entropy rate  $\lim_{n \rightarrow \infty} H(D^n | X^n, Y)/n$ ?

# Key Lemma

## Lemma

$$\lim_{n \rightarrow \infty} \frac{1}{n} H(D^n | X^n, Y(X^n)) = d \frac{\mathbb{E}[L \log L]}{\mathbb{E}[L]} + O(d^{1.9})$$

*provided  $\mathbb{E}[L^2 \log L] < d^{-0.05}$ .*

- Lemma holds uniformly for all processes  $\mathbb{X}$ .
- Key to showing both achievability and upper bound.

# Key Lemma

## Lemma

$$\lim_{n \rightarrow \infty} \frac{1}{n} H(D^n | X^n, Y(X^n)) = d \frac{\mathbb{E}[L \log L]}{\mathbb{E}[L]} + O(d^{1.9})$$

*provided  $\mathbb{E}[L^2 \log L] < d^{-0.05}$ .*

- Lemma holds uniformly for all processes  $\mathbb{X}$ .
- Key to showing both achievability and upper bound.

# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*



# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

# Proof of Key Lemma: sparsity of deletions

## Simple case: no runs of length 1.

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

- No run disappears completely

$$\begin{array}{cccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & & \\
 X^n & : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & & & \dots \\
 & & \mathcal{S}_1 & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & & \\
 & & \mathcal{S}_i & \leftarrow & \mathcal{R}_i & & & & & & & & & 
 \end{array}$$

Given  $(X^n, Y(X^n))$ :

- We know which runs have (one) deletion
- Each bit of run equally likely candidate for deletion

# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

- No run disappears completely

$$\begin{array}{cccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & & \\
 X^n & : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & \\
 & & \mathcal{S}_i & \leftarrow & \mathcal{R}_i & & & & & & & & 
 \end{array}$$

Given  $(X^n, Y(X^n))$ :

- We know which runs have (one) deletion
- Each bit of run equally likely candidate for deletion

# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

- No run disappears completely

$$\begin{array}{cccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & & \\
 X^n & : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & \\
 & & \mathcal{S}_i & \leftarrow & \mathcal{R}_i & & & & & & & & 
 \end{array}$$

Given  $(X^n, Y(X^n))$ :

- We know which runs have (one) deletion
- Each bit of run equally likely candidate for deletion

# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

- No run disappears completely

$$\begin{array}{cccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & & \\
 X^n & : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & \\
 & & \mathcal{S}_i & \leftarrow & \mathcal{R}_i & & & & & & & & 
 \end{array}$$

Given  $(X^n, Y(X^n))$ :

- We know which runs have (one) deletion
- Each bit of run equally likely candidate for deletion

# Proof of Key Lemma: sparsity of deletions

**Simple case: no runs of length 1.**

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

- No run disappears completely

$$\begin{array}{cccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & & \\
 X^n & : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & \\
 & & \mathcal{S}_i \leftarrow \mathcal{R}_i & & & & & & & & & & 
 \end{array}$$

Given  $(X^n, Y(X^n))$ :

- We know which runs have (one) deletion
- Each bit of run equally likely candidate for deletion

# Proof of Key Lemma: sparsity of deletions

## Simple case: no runs of length 1.

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

Run of length  $\ell$  has:

- probability  $p_L(\ell)$  of occurring.
- probability  $\approx \ell d$  of suffering a deletion.
- Contribution  $\log \ell$  to  $H(D^n|X^n, Y)$  if deletion

$$\lim_{n \rightarrow \infty} \frac{1}{n} H(D^n|X^n, Y) \approx \frac{d}{\mathbb{E}[L]} \sum_{\ell=2}^{\infty} p_L(\ell) \ell \log \ell = d \frac{\mathbb{E}[L \log L]}{\mathbb{E}[L]}$$

# Proof of Key Lemma: sparsity of deletions

## Simple case: no runs of length 1.

Imagine *no run in  $X^n$  has  $> 1$  deletion in total.*

Run of length  $\ell$  has:

- probability  $p_L(\ell)$  of occurring.
- probability  $\approx \ell d$  of suffering a deletion.
- Contribution  $\log \ell$  to  $H(D^n|X^n, Y)$  if deletion

$$\lim_{n \rightarrow \infty} \frac{1}{n} H(D^n|X^n, Y) \approx \frac{d}{\mathbb{E}[L]} \sum_{\ell=2}^{\infty} p_L(\ell) \ell \log \ell = d \frac{\mathbb{E}[L \log L]}{\mathbb{E}[L]}$$



# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{ccccccc}
 R_1 & R_2 & R_3 & R_4 & R_5 & R_6 & \\
 x^n & = & 0011011101110 \dots \\
 x(x^n) & = & 001111110110 \dots \\
 S_1 & & S_2 & & S_3 & & S_4
 \end{array}$$

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & \dots \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & & & 
 \end{array}$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & \mathcal{R}_5 & & \mathcal{R}_6 & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & & \dots \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & & 
 \end{array}$$

$$|\mathcal{S}_1| = |\mathcal{R}_1|$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & \mathcal{R}_5 & & \mathcal{R}_6 & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & & \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & & & \mathcal{S}_3 & & \mathcal{S}_4 & & & 
 \end{array}$$

$$|\mathcal{S}_1| = |\mathcal{R}_1| \quad \Rightarrow \quad \mathcal{S}_1 \leftarrow \mathcal{R}_1$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n | X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & \mathcal{R}_5 & & \mathcal{R}_6 & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & & \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & & \mathcal{S}_3 & & \mathcal{S}_4 & & & & 
 \end{array}$$

$$|\mathcal{S}_2| > |\mathcal{R}_2|$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{rcccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & 
 \end{array}$$

$$|\mathcal{S}_2| > |\mathcal{R}_2| \quad \Rightarrow \quad \mathcal{S}_2 \leftarrow \mathcal{R}_2 + \mathcal{R}_3 + \mathcal{R}_4$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total*.

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{rccccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots & \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & \dots & & & \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & & & 
 \end{array}$$

$$|\mathcal{S}_3| = |\mathcal{R}_5|$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & \mathcal{R}_5 & & \mathcal{R}_6 & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & & \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & & & \mathcal{S}_3 & & \mathcal{S}_4 & & & 
 \end{array}$$

$$|\mathcal{S}_3| = |\mathcal{R}_5| \quad \Rightarrow \quad \mathcal{S}_3 \leftarrow \mathcal{R}_5$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .



# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{rcccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & & 
 \end{array}$$

$$|\mathcal{S}_4| < |\mathcal{R}_6|$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{rcccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & & 
 \end{array}$$

$$|\mathcal{S}_4| < |\mathcal{R}_6| \quad \Rightarrow \quad \mathcal{S}_4 \leftarrow \mathcal{R}_6$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots & \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & \dots & & & \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & 
 \end{array}$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & & \mathcal{R}_2 & & \mathcal{R}_3 & & \mathcal{R}_4 & & \mathcal{R}_5 & & \mathcal{R}_6 & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & & \dots \\
 & & \mathcal{S}_1 & & & & \mathcal{S}_2 & & & & \mathcal{S}_3 & & \mathcal{S}_4 & & 
 \end{array}$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Proof of Key Lemma: sparsity of deletions

## General case

Imagine *no three consecutive runs in  $X^n$  have  $> 1$  deletion in total.*

- Only runs of length 1 can disappear
- Runs in  $Y$  unambiguously associated with runs in  $X^n$

$$\begin{array}{cccccccccccccccc}
 & & \mathcal{R}_1 & \mathcal{R}_2 & \mathcal{R}_3 & \mathcal{R}_4 & \mathcal{R}_5 & \mathcal{R}_6 & & & & & & & & & \\
 X^n & : & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 & \dots & \\
 Y(X^n) & : & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & \dots & & & \\
 & & \mathcal{S}_1 & & & \mathcal{S}_2 & & \mathcal{S}_3 & \mathcal{S}_4 & & & & & & & & 
 \end{array}$$

- We know  $\mathcal{R}_3$  and  $\mathcal{R}_6$  have deletion.
- $\mathcal{R}_3$  has no contribution to  $H(D^n|X^n, Y)$ .

# Upper bound

Achievability of  $1 - d \log(1/d)$

$\Downarrow$

$$H(\mathbb{X}_{\text{opt}}) > 1 - d^{1-\epsilon}.$$

$\Downarrow$

$\mathbb{X}_{\text{opt}}$  is 'close' to Bernoulli(1/2) process.

# Conclusion

## **We obtained for deletion channel with small $d$ :**

- Asymptotic expansion of capacity upto order  $d^2$ .
- Optimal coding scheme.

## **Further directions:**

- Explicit upper and lower bounds.
- Next terms in expansion.
- Is the series convergent?
- Series expansion for other 'hard' channels ... ?

# Conclusion

## **We obtained for deletion channel with small $d$ :**

- Asymptotic expansion of capacity upto order  $d^2$ .
- Optimal coding scheme.

## **Further directions:**

- Explicit upper and lower bounds.
- Next terms in expansion.
- Is the series convergent?
- Series expansion for other 'hard' channels ... ?



# Conclusion

## We obtained for deletion channel with small $d$ :

- Asymptotic expansion of capacity upto order  $d^2$ .
- Optimal coding scheme.

## Further directions:

- Explicit upper and lower bounds.
- Next terms in expansion.
- Is the series convergent?
- Series expansion for other 'hard' channels ... ?

THANK YOU!