

Universal algebra over locally presentable categories

Yuto Kawase

RIMS, Kyoto University

November 24, 2024.

Categories in Tokyo 1st



← Today's slides

This work is supported by JSPS KAKENHI Grant Numbers JP24KJ1462.

1 Relativization of universal algebra

2 Birkhoff's variety theorem

3 Filtered colimit elimination

Relativization via monads

Theorem ([Lin69])

There is an equivalence

$$\mathbf{Th}^S \simeq \mathbf{Mnd}_f(\mathbf{Set}^S).$$

Here,

\mathbf{Th}^S : the category of S -sorted equational theories,

$\mathbf{Mnd}_f(\mathbf{Set}^S)$: the category of finitary monads on \mathbf{Set}^S .

S -sorted equational theory = finitary monad on \mathbf{Set}^S

↓ generalize

???

\mathcal{A} -relative algebraic theory = κ -ary monad on \mathcal{A}
(\mathcal{A} : a locally κ -presentable category)

Relative algebraic theories

Informal definition [Kaw23]

\mathcal{A} : a (locally presentable) category

An \mathcal{A} -relative algebraic theory consists of:

- a set Ω of partial operators;
- a set E of implications $\dots (\underbrace{\text{YYY}}_{\text{postcondition}} \text{ whenever } \underbrace{\text{XXX}}_{\text{precondition}})$

such that

- For each operator $\omega \in \Omega$, its domain must be defined by “ \mathcal{A} ’s language.”
- For each implication in E , its precondition must be written in “ \mathcal{A} ’s language.”

\mathbf{Set}^S -relative algebraic theories = S -sorted equational theories

A generalized Linton theorem

Theorem ([Kaw23; Kaw24])

For a locally κ -presentable category \mathcal{A} , there is an equivalence

$$\mathbf{Th}_{\kappa}^{\mathcal{A}} \simeq \mathbf{Mnd}_{\kappa}(\mathcal{A}).$$

Here,

$\mathbf{Th}_{\kappa}^{\mathcal{A}}$: the category of \mathcal{A} -relative (κ -ary) algebraic theories,

$\mathbf{Mnd}_{\kappa}(\mathcal{A})$: the category of κ -ary monads on \mathcal{A} .

Example: small categories

Example

A **small category** consists of:

- a base quiver $\text{mor}\mathcal{C} \xrightleftharpoons[c]{d} \text{ob}\mathcal{C}$;
- a total operator $\text{id}: \text{ob}\mathcal{C} \rightarrow \text{mor}\mathcal{C}$;
- a **partial** operator $\circ: \text{mor}\mathcal{C} \times \text{mor}\mathcal{C} \rightarrow \text{mor}\mathcal{C}$ such that

$$g \circ f \text{ is defined iff } d(g) = c(f)$$

which satisfy the following:

- $d(\text{id}(x)) = x$ and $c(\text{id}(x)) = x$;
- $d(g \circ f) = d(f)$ and $c(g \circ f) = c(g)$ whenever $d(g) = c(f)$;
- $f \circ \text{id}(d(f)) = f$ and $\text{id}(c(f)) \circ f = f$;
- $(h \circ g) \circ f = h \circ (g \circ f)$ whenever $d(h) = c(g)$ and $d(g) = c(f)$.

Small categories are algebras over quivers.

Further examples

Example

| | | algebras over \sim |
|------------------------------------|--------------------|---------------------------|
| small categories | \rightsquigarrow | quivers |
| UDO semirings | \rightsquigarrow | posets |
| partial Boolean algebras | \rightsquigarrow | graphs |
| monoid-graded rings | \rightsquigarrow | monoid-graded sets |
| generalized complete metric spaces | \rightsquigarrow | generalized metric spaces |
| Banach spaces | \rightsquigarrow | pointed metric spaces |

A technical remark

Definition ([PV07])

- (κ -ary) **partial Horn theory** \cdots a logical theory based on multi-sorts, partial functions, relations, and (partial) Horn implications.
- **Mod \mathbb{S}** \cdots the category of models of a partial Horn theory \mathbb{S} .

Theorem ([PV07])

TFAE for a category \mathcal{A} :

- 1 \mathcal{A} is locally κ -presentable.
- 2 $\mathcal{A} \simeq \mathbf{Mod} \mathbb{S}$ for some κ -ary partial Horn theory \mathbb{S} .

We actually define **\mathbb{S} -relative algebraic theories** for partial Horn theories \mathbb{S} .

$$\rightsquigarrow \mathcal{A}\text{-rel. alg. theory} = \mathbb{S}\text{-rel. alg. theory where } \mathcal{A} \simeq \mathbf{Mod} \mathbb{S}.$$

1 Relativization of universal algebra

2 Birkhoff's variety theorem

3 Filtered colimit elimination

Birkhoff's variety theorem

Birkhoff's variety theorem [Bir35]

(Ω, E) : a single-sorted algebraic theory. $\mathcal{C} \subseteq \mathbf{Alg}(\Omega, E)$: fullsub.

TFAE:

- ① $\mathcal{C} \subseteq \mathbf{Alg}(\Omega, E)$ is definable by equations.
- ② $\mathcal{C} \subseteq \mathbf{Alg}(\Omega, E)$ is closed under products, subobjects, and quotients.

closed under products: $A_i \in \mathcal{C} \implies \prod_i A_i \in \mathcal{C}$.

closed under subobjects: $B \subseteq A$: sub, $A \in \mathcal{C} \implies B \in \mathcal{C}$.

closed under quotients: $A \twoheadrightarrow B$: surj, $A \in \mathcal{C} \implies B \in \mathcal{C}$.

A generalized Birkhoff's theorem

Theorem ([Kaw23; Kaw24])

(Ω, E) : an \mathcal{A} -relative (κ -ary) algebraic theory. $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$: fullsub.
TFAE:

- ① $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$ is definable.
- ② $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$ is closed under products, closed subobjects, (U, κ) -pure quotients, and κ -filtered colimits.

| single-sorted alg. (Set-relative alg.) | | \mathcal{A} -relative alg. |
|---|--------------------|--|
| products | \rightsquigarrow | products |
| subobjects | \rightsquigarrow | <i>closed subobjects</i> |
| quotients | \rightsquigarrow | <i>(U, κ)-pure quotients</i> |
| | \rightsquigarrow | <i>κ-filtered colimits (new)</i> |

What are closed subobjects and (U, κ) -pure quotients?

\mathcal{A} ... a locally κ -presentable category
 (Ω, E) ... an \mathcal{A} -rel. alg. theory
 $\mathbf{Alg}(\Omega, E) \xrightarrow{U} \mathcal{A}$... the forgetful functor

Informal definition

① A subalg. $B \subseteq A$ in $\mathbf{Alg}(\Omega, E)$ is **closed** if:

- ▶ For every relation R in “the language of \mathcal{A} ,”

$$R(\vec{b}) \text{ holds in } UA \quad \Rightarrow \quad R(\vec{b}) \text{ holds in } UB.$$

② $A \xrightarrow{p} B$ in $\mathbf{Alg}(\Omega, E)$ is a **(U, κ) -pure quotient** if:

- ▶ For every κ -ary formula in “the language of \mathcal{A} ,”

$$\varphi(\vec{b}) \text{ holds in } UB \quad \Rightarrow \quad \exists \vec{a} \xrightarrow{Up} \vec{b} \text{ s.t. } \varphi(\vec{a}) \text{ holds in } UA.$$

Example

Pos ... the category of posets.

(Ω, \emptyset) ... a **Pos**-rel. alg. theory defined by
 $\Omega := \{\ominus\}, \quad x \ominus y \text{ is defined iff } x \geq y.$

$\mathbf{Alg}(\Omega, \emptyset) \xrightarrow{U} \mathbf{Pos}$... the forgetful functor.

In $\mathbf{Alg}(\Omega, \emptyset)$, under $x \ominus y := x - y$ in \mathbb{N} ,

• $\{0 \leq 2 \leq 3\} \subseteq \{0 < 1 < 2 < \dots\}$... subalgebra, but **not** closed.

• $\{0 \leq 2 \leq 4\} \subseteq \{0 < 1 < 2 < \dots\}$... closed subalgebra.

• $\left\{ \begin{array}{cc} 0 & 0 \\ \wedge & \\ 1 & 1 \\ & \wedge \\ & 2 \end{array} \right\} \rightarrow \left\{ \begin{array}{c} 0 \\ \wedge \\ 1 \\ \wedge \\ 2 \end{array} \right\}$... surjection, but **not** a (U, \aleph_0) -pure quotient.

• $\left\{ \begin{array}{cccc} 0 & 0 & 0 & \dots \\ & \wedge & \wedge & \\ & 1 & 1 & \dots \\ & & \wedge & \\ & & 2 & \dots \\ & & & \ddots \end{array} \right\} \rightarrow \left\{ \begin{array}{c} 0 \\ \wedge \\ 1 \\ \wedge \\ 2 \\ \wedge \\ \vdots \end{array} \right\}$... (U, \aleph_0) -pure quotient,
 but **not** (U, \aleph_1) -pure quotient.

Filtered colimits are necessary

Example ($\mathbf{Set}^{\mathbb{N}}$ -relative algebra [ARV12])

$$\mathcal{E} := \{1\} \cup \{A \in \mathbf{Set}^{\mathbb{N}} \mid \exists m \in \mathbb{N}. A_m = \emptyset\}.$$

$\mathcal{E} \subseteq \mathbf{Set}^{\mathbb{N}}$ is closed under...

- ✓ products
- ✓ closed subobjects = sort-wise injections
- ✓ pure quotients = sort-wise surjections
- ✗ **filtered colimits**

$$\begin{array}{llllll} A_0 & := & (\emptyset, & \emptyset, & \emptyset, & \emptyset, & \dots) & \in \mathcal{E} \\ \cap & & \cap & \cap & \cap & \cap & & \\ A_1 & := & (2, & \emptyset, & \emptyset, & \emptyset, & \dots) & \in \mathcal{E} \\ \cap & & \cap & \cap & \cap & \cap & & \\ A_2 & := & (2, & 2, & \emptyset, & \emptyset, & \dots) & \in \mathcal{E} \\ \cap & & \cap & \cap & \cap & \cap & & \\ \vdots & & \vdots & \vdots & \vdots & \vdots & & \\ \text{Colim}_{n \in \omega} A_n & = & (2, & 2, & 2, & 2, & \dots) & \notin \mathcal{E} \end{array}$$

Example (\mathbf{Set}_ω -relative algebra [Kaw25])

\mathbf{Set}_ω ... the category of sets with countably many constants $(c_n)_n$.

$$\mathcal{E} := \{1\} \cup \{A \in \mathbf{Set}_\omega \mid \exists i, j \text{ s.t. } c_i \neq c_j \text{ in } A\}.$$

$\mathcal{E} \subseteq \mathbf{Set}_\omega$ is closed under...

- ✓ products
- ✓ closed subobjects = subalgebras
- ✓ pure quotients = surjections that do not merge any constants
- ✗ **filtered colimits**

$$\begin{array}{rclcl}
 A_0 & := & \{c_0, & c_1, & c_2, & c_3, & \dots, & \infty\} & \in \mathcal{E} \\
 \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & & \downarrow & \\
 A_1 & := & \{c_0 = c_1, & c_2, & c_3, & \dots, & \infty\} & \in \mathcal{E} \\
 \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & & \downarrow & \\
 A_2 & := & \{c_0 = c_1 = c_2, & c_3, & \dots, & \infty\} & \in \mathcal{E} \\
 \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & & \downarrow & \\
 \vdots & & \vdots & \vdots & \vdots & \vdots & & \vdots & \\
 \operatorname{Colim}_{n \in \omega} A_n & := & \{c_0 = c_1 = c_2 = c_3 = \dots, & \infty\} & \notin \mathcal{E}
 \end{array}$$

The filtered colimit elimination problem

However, filtered colimits are **not** required for **Set**-rel. alg. in Birkhoff's theorem.

Question

Why can filtered colimits be eliminated in the case of **Set**-relative algebras?

Answer

The category **Set** satisfies a “noetherian” condition.

1 Relativization of universal algebra

2 Birkhoff's variety theorem

3 Filtered colimit elimination

A noetherian condition for categories

Definition ([Kaw25])

A category \mathcal{A} satisfies the **ascending chain condition (ACC)** if it has no chain $A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \cdots$ of objects such that there is no morphism $A_n \leftarrow A_{n+1}$ for all n .

Example

Set satisfies ACC.

Proof.

Let $A_0 \rightarrow A_1 \rightarrow \cdots$ be an ω -chain of sets. If there is no map $A_0 \leftarrow A_1$, then $A_0 = \emptyset$ and $A_1 \neq \emptyset$. Thus, a map $A_1 \leftarrow A_2$ exists. □

More generally,

Proposition

\mathbf{Set}^S satisfies ACC \Leftrightarrow the set S is finite.

Filtered colimit elimination

Theorem ([Kaw25; Kaw24])

(Ω, E) : an \mathcal{A} -relative (κ -ary) algebraic theory. $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$: fullsub.

If \mathcal{A} satisfies ACC,

TFAE:

- ① $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$ is definable.
- ② $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$ is closed under products, closed subobjects, (U, κ) -pure quotients, and ~~κ -filtered colimits~~.

Some applications of filtered colimit elimination

Corollary

- **Set** satisfies ACC.
 - ↪ fil.colim.elim. holds for **single-sorted alg.**
 - ↪ The classical Birkhoff theorem [Bir35]
- **Setⁿ** satisfied ACC.
 - ↪ fil.colim.elim. holds for **finite-sorted alg.**
 - ↪ This subsumes a result in [ARV12].
- **Pos** satisfied ACC.
 - ↪ fil.colim.elim. holds for **ordered alg.**
 - ↪ This subsumes a result in [Blo76].
- **Met_∞**, the category of generalized metric spaces, satisfied ACC.
 - ↪ fil.colim.elim. holds for **metric alg.**
 - ↪ This subsumes a result in [Hin16].

Filtered colimit elimination: sketch of proof

fullsub $\mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$: closed under products, closed sub, (U, κ) -pure quo.
 $(A_J)_{J \in \mathbb{I}}$: a κ -filtered diagram s.t. $A_J \in \mathcal{E}$.

For each $J \in \mathbb{I}$, we can construct a “nice” wide sub-diagram $J \in \mathbb{I}_J \subseteq \mathbb{I}$.

$$\begin{array}{ccccc}
 \prod_{I \in \mathbb{I}} A_I & & & & \\
 \uparrow m_J & \nwarrow m & & \xrightarrow{\text{Colim}_{J \in \mathbb{I}} \pi_J} & \\
 & \text{Colim}_{J \in \mathbb{I}} \text{Lim}_{I \in \mathbb{I}_J} A_I & & & \text{Colim}_{J \in \mathbb{I}} A_J \quad \text{in } \mathbf{Alg}(\Omega, E) \\
 & \nearrow & & \nearrow & \\
 \text{Lim}_{I \in \mathbb{I}_J} A_I & \xrightarrow{\pi_J} & A_J & &
 \end{array}$$

$\leadsto \mathcal{E} \subseteq \mathbf{Alg}(\Omega, E)$ is closed under κ -filtered colimits.

□

Weak converse

Theorem ([Kaw25])

\mathcal{A} : a l.f.p. category. Assume that, for every fullsub. of \mathcal{A} , closure under filtered colimits follows from the others: **P**(products), **S**(closed sub), **H**(\aleph_0 -pure quo).

Then,

- 1 The fullsub $\mathcal{A}_{\text{fp},c} := \{\text{finitely presentable connected objs}\} \subseteq \mathcal{A}$ satisfies ACC.
- 2 If $\emptyset \xrightarrow{!} 1$ in \mathcal{A} is strongly monic, the fullsub $\mathcal{A}_{\text{fp}} := \{\text{finitely presentable objs}\} \subseteq \mathcal{A}$ satisfies ACC.

Sketch of proof:

Let $A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$ in \mathcal{A}_{fp} . Consider

$$\mathcal{C} := \{X \mid \exists n. X \xrightarrow{\exists} A_n\} \subseteq \mathcal{A}.$$

Using finite presentability, its **HSP**-closure can be computed as

$$\mathbf{HSP}(\mathcal{C}) = \mathbf{S}(1) \cup \mathbf{H}(\mathcal{C}).$$

Since $A_n + A_n \in \mathcal{C}$ ($\forall n$), $B := \text{Colim}_{n \in \omega} (A_n + A_n) \in \mathbf{HSP}(\mathcal{C})$.

The additional conditions ensure that $B \notin \mathbf{S}(1)$. $\therefore B \in \mathbf{H}(\mathcal{C})$

$$\begin{aligned}
& A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \cdots \text{ in } \mathcal{A}_{\text{fp}} \\
& \mathcal{E} := \{X \mid \exists n. X \xrightarrow{\exists} A_n\} \subseteq \mathcal{A} \\
& B := \operatorname{Colim}_{n \in \omega} (A_n + A_n) \in \mathbf{H}(\mathcal{E})
\end{aligned}$$

Thus, we have:

$$\begin{array}{ccccc}
& & & X \xrightarrow{\in \mathcal{E}} A_N \\
& \nearrow & & \downarrow \text{pure quo} \\
A_n & \xrightarrow{\text{coproj}} & A_n + A_n & \xrightarrow{\text{coproj}} & B
\end{array}
\quad (\forall n \geq N)$$

Thus, $A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \cdots$ eventually “stabilizes.”



Open problems

Open problem 1

Is there any locally presentable category that satisfies filtered colimit elimination but not ACC?

More precisely, is there any κ -ary partial Horn theory \mathbb{S} for some κ that satisfies the following conditions?

- Every full subcategory of $\mathbf{Mod}\,\mathbb{S}$ is closed under κ -filtered colimits whenever it is closed under products, \mathbb{S} -closed subobjects, and κ -pure quotients.
- The category $\mathbf{Mod}\,\mathbb{S}$ does not satisfy ACC.

The next one is weaker than 1 and independent of partial Horn theories:

Open problem 2

Is there any locally finitely presentable category that does not satisfy ACC but satisfies it for the full subcategory of finitely presentable objects?

Thank you!



Today's slides



My homepage

References I

- [AR04] J. Adámek and J. Rosický. “On pure quotients and pure subobjects”. In: *Czechoslovak Math. J.* 54(129).3 (2004), pp. 623–636.
- [AR94] J. Adámek and J. Rosický. *Locally Presentable and Accessible Categories*. Vol. 189. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1994.
- [ARV12] J. Adámek, J. Rosický, and E. M. Vitale. “Birkhoff’s variety theorem in many sorts”. In: *Algebra Universalis* 68.1-2 (2012), pp. 39–42.
- [Adá+21] J. Adámek et al. “Finitary monads on the category of posets”. In: *Math. Structures Comput. Sci.* 31.7 (2021), pp. 799–821.
- [BH12] B. van den Berg and C. Heunen. “Noncommutativity as a colimit”. In: *Appl. Categ. Structures* 20.4 (2012), pp. 393–414.
- [BMW12] C. Berger, P.-A. Melliès, and M. Weber. “Monads with arities and their associated theories”. In: *J. Pure Appl. Algebra* 216.8-9 (2012), pp. 2029–2048.
- [Bir35] G. Birkhoff. “On the structure of abstract algebras”. In: *Math. Proc. Cambridge Philos. Soc.* 31.4 (1935), pp. 433–454.
- [Blo76] S. L. Bloom. “Varieties of ordered algebras”. In: *J. Comput. System Sci.* 13.2 (1976), pp. 200–212.
- [BG19] J. Bourke and R. Garner. “Monads and theories”. In: *Adv. Math.* 351 (2019), pp. 1024–1071.
- [CV98] A. Carboni and E. M. Vitale. “Regular and exact completions”. In: *J. Pure Appl. Algebra* 125.1-3 (1998), pp. 79–116.
- [Gol03] J. S. Golan. *Semirings and Affine Equations over Them: Theory and Applications*. Vol. 556. Mathematics and its Applications. Kluwer Academic Publishers Group, Dordrecht, 2003, pp. xiv+241.

References II

- [Hin16] W. Hino. *Varieties of metric and quantitative algebras*. 2016. [arXiv: 1612.06054 \[cs.LO\]](#).
- [Kaw23] Y. Kawase. *Birkhoff's variety theorem for relative algebraic theories*. 2023. [arXiv: 2304.04382 \[math.CT\]](#).
- [Kaw24] Y. Kawase. *Relativized universal algebra via partial Horn logic*. 2024. [arXiv: 2403.19661 \[math.CT\]](#).
- [Kaw25] Y. Kawase. "Filtered colimit elimination from Birkhoff's variety theorem". In: *J. Pure Appl. Algebra* 229.1 (2025), Paper No. 107794, 25.
- [Lin69] F. E. J. Linton. "An outline of functorial semantics". In: *Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*. Springer, Berlin, 1969, pp. 7–52.
- [LP23] R. B. B. Lucyshyn-Wright and J. Parker. "Diagrammatic presentations of enriched monads and varieties for a subcategory of arities". In: *Appl. Categ. Structures* 31.5 (2023), Paper No. 40, 39.
- [PV07] E. Palmgren and S. J. Vickers. "Partial Horn logic and cartesian categories". In: *Ann. Pure Appl. Logic* 145.3 (2007), pp. 314–353.
- [Ros21] J. Rosický. "Metric monads". In: *Math. Structures Comput. Sci.* 31.5 (2021), pp. 535–552.
- [Wea95] N. Weaver. "Quasi-varieties of metric algebras". In: *Algebra Universalis* 33.1 (1995), pp. 1–9.