

Collective motion in large deviations of active particles

Yann-Edwin Keta

in collaboration with E. Fodor, F. van Wijland, M.E. Cates, and R.L. Jack

APS March Meeting 2021

19/03/2021

Phys. Rev. E **103**, 022603 (2021)


 [yketa/DAMTP_MSC_2019_Wiki](#)



SIMONS
FOUNDATION



- 1 Model of active Brownian particles.
- 2 Active work and its large deviations.
- 3 Description of dynamical phase transition.
- 4 Mechanism for collective motion.



standard ABP model

$$\dot{\mathbf{r}}_i = -D\nabla_i U(\{\mathbf{r}_j\}) + v_0 \mathbf{u}(\theta_i) + \sqrt{2D}\boldsymbol{\eta}_i \quad (1)$$

$$\dot{\theta}_i = \sqrt{2D_r}\xi_i \quad (2)$$

$$\tilde{l}_p = v_0 D_r^{-1} / \sigma \quad (3)$$

T. Nemoto *et al.*, *Physical Review E* **99**, 022605 (2019).

G. S. Redner *et al.*, *Physical Review Letters* **110**, 055701 (2013).

How far does the active forcing translate into real motion?

How far does the active forcing translate into real motion?

$$\text{active work } w_{\tau} = \frac{1}{v_0 N_{\tau}} \sum_{i=1}^N \int_0^{\tau} \mathbf{u}(\theta_i) \circ \dot{\mathbf{r}}_i \, dt \quad (4)$$

How far does the active forcing translate into real motion?


active work $w_{\tau} = \frac{1}{v_0 N_{\tau}} \sum_{i=1}^N \int_0^{\tau} \mathbf{u}(\theta_i) \circ \dot{\mathbf{r}}_i dt$ (4)

flocking



$$\dot{\mathbf{r}}_i \approx v_0 \mathbf{u}(\theta_i) \Rightarrow \langle w_{\tau} \rangle = 1$$

jamming



$$\dot{\mathbf{r}}_i \approx 0 \Rightarrow \langle w_{\tau} \rangle = 0$$

How does the active work control emerging behaviours?

-
- H. Touchette, *Physics Reports* **478**, 1–69 (2009).
 - R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).
 - T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).
 - F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN\tau w_\tau} \quad (5)$$

H. Touchette, *Physics Reports* **478**, 1–69 (2009).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN_\tau w_\tau} \quad (5)$$

- 1 Cloning algorithm \Rightarrow trajectories biased with respect to w_τ .

H. Touchette, *Physics Reports* **478**, 1–69 (2009).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN\tau w_\tau} \quad (5)$$

➊ Cloning algorithm \Rightarrow trajectories biased with respect to w_τ .

$$\langle \mathcal{A} \rangle_s = \frac{\langle \mathcal{A} e^{-sN\tau w_\tau} \rangle}{\langle e^{-sN\tau w_\tau} \rangle} \quad (6)$$

H. Touchette, *Physics Reports* **478**, 1–69 (2009).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN_\tau w_\tau} \quad (5)$$

1 Cloning algorithm \Rightarrow trajectories biased with respect to w_τ .

$$\langle \mathcal{A} \rangle_s = \frac{\langle \mathcal{A} e^{-sN_\tau w_\tau} \rangle}{\langle e^{-sN_\tau w_\tau} \rangle} \quad (6)$$

$$\psi(s) = \lim_{\tau \rightarrow \infty} \frac{1}{N_\tau} \log \left\langle e^{-sN_\tau w_\tau} \right\rangle \quad (7)$$

H. Touchette, *Physics Reports* **478**, 1–69 (2009).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN_\tau w_\tau} \quad (5)$$

- ❶ Cloning algorithm \Rightarrow trajectories biased with respect to w_τ .

$$\langle \mathcal{A} \rangle_s = \frac{\langle \mathcal{A} e^{-sN_\tau w_\tau} \rangle}{\langle e^{-sN_\tau w_\tau} \rangle} \quad (6)$$

$$\psi(s) = \lim_{\tau \rightarrow \infty} \frac{1}{N_\tau} \log \left\langle e^{-sN_\tau w_\tau} \right\rangle \quad (7)$$

- ❷ Singularities in $\psi \Rightarrow$ qualitative changes at dynamical transition.

H. Touchette, *Physics Reports* **478**, 1–69 (2009).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

How does the active work control emerging behaviours?

$$P_s[\{\mathbf{r}_i, \theta_i\}_0^\tau] \propto P_0[\{\mathbf{r}_i, \theta_i\}_0^\tau] e^{-sN\tau \mathbf{w}_\tau} \quad (5)$$

- ❶ Cloning algorithm \Rightarrow trajectories biased with respect to \mathbf{w}_τ .

$$\langle \mathcal{A} \rangle_s = \frac{\langle \mathcal{A} e^{-sN\tau \mathbf{w}_\tau} \rangle}{\langle e^{-sN\tau \mathbf{w}_\tau} \rangle} \quad (6)$$

$$\psi(s) = \lim_{\tau \rightarrow \infty} \frac{1}{N\tau} \log \langle e^{-sN\tau \mathbf{w}_\tau} \rangle \quad (7)$$

- ❷ Singularities in $\psi \Rightarrow$ qualitative changes at dynamical transition.
 ❸ Compute probabilities of these fluctuations.

$$P(\mathbf{w}_\tau) \asymp \exp(-N\tau I(\mathbf{w}_\tau)) \quad (8)$$

$$I(\mathbf{w}_\tau) = \sup_s \{-s\mathbf{w}_\tau - \psi(s)\} \quad (9)$$

H. Touchette, *Physics Reports* **478**, 1–69 (2009).


R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

T. Nemoto *et al.*, *Physical Review E* **93**, 062123 (2016).

F. Cagnetta *et al.*, *Physical Review Letters* **119**, 158002 (2017).

unbiased steady state


$$s = 0$$



MIPS

biased to large w_T

$$s < 0$$



CM

T. Nemoto *et al.*, *Physical Review E* 99, 022605 (2019).



How much bias is needed for the transition to occur?

How much bias is needed for the transition to occur?

$$\boldsymbol{\nu} = \frac{1}{N} \sum_{i=1}^N \mathbf{u}(\theta_i), \quad \bar{\nu}_\tau = \frac{1}{\tau} \int_0^\tau |\boldsymbol{\nu}(t)| \, dt \quad (10)$$


How much bias is needed for the transition to occur?

$$\boldsymbol{\nu} = \frac{1}{N} \sum_{i=1}^N \mathbf{u}(\theta_i), \quad \bar{\nu}_\tau = \frac{1}{\tau} \int_0^\tau |\boldsymbol{\nu}(t)| dt \quad (10)$$



How much bias is needed for the transition to occur?

$$\boldsymbol{\nu} = \frac{1}{N} \sum_{i=1}^N \mathbf{u}(\theta_i), \quad \bar{\nu}_\tau = \frac{1}{\tau} \int_0^\tau |\boldsymbol{\nu}(t)| dt \quad (10)$$



separation between steady state physics and symmetry breaking physics

“Large deviations occur according to the least unlikely mechanism.”

F. den Hollander, *Large deviations*, Vol. 14 (American Mathematical Soc., 2008).

R. L. Jack, *The European Physical Journal B* 93, 1–22 (2020).

S. Whitelam *et al.*, *The Journal of Chemical Physics* 148, 154902 (2018).

T. GrandPre *et al.*, *Physical Review E* 103, 012613 (2021).

“Large deviations occur according to the least unlikely mechanism.”

$$\dot{\mathbf{r}}_i = -D\nabla_i U + v_0 \mathbf{u}(\theta_i) + \sqrt{2D}\boldsymbol{\eta}_i \quad (12)$$

$$\dot{\theta}_i = -D_r \frac{\partial}{\partial \theta_i} U_g^{\text{mod}} + \sqrt{2D_r}\xi_i \quad (13)$$

F. den Hollander, *Large deviations*, Vol. 14 (American Mathematical Soc., 2008).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

S. Whitelam *et al.*, *The Journal of Chemical Physics* **148**, 154902 (2018).

T. GrandPre *et al.*, *Physical Review E* **103**, 012613 (2021).

“Large deviations occur according to the least unlikely mechanism.”

$$\dot{\mathbf{r}}_i = -D\nabla_i U + v_0 \mathbf{u}(\theta_i) + \sqrt{2D}\boldsymbol{\eta}_i \quad (12)$$

$$\dot{\theta}_i = -D_r \frac{\partial}{\partial \theta_i} U_g^{\text{mod}} + \sqrt{2D_r} \xi_i \quad (13)$$

$$U_g^{\text{mod}} = -g \frac{N|\boldsymbol{\nu}|^2}{D_r} \quad (14)$$

F. den Hollander, *Large deviations*, Vol. 14 (American Mathematical Soc., 2008).

R. L. Jack, *The European Physical Journal B* **93**, 1–22 (2020).

S. Whitelam *et al.*, *The Journal of Chemical Physics* **148**, 154902 (2018).

T. GrandPre *et al.*, *Physical Review E* **103**, 012613 (2021).

“Large deviations occur according to the least unlikely mechanism.”

$$\dot{\mathbf{r}}_i = -D\nabla_i U + v_0 \mathbf{u}(\theta_i) + \sqrt{2D}\boldsymbol{\eta}_i \quad (12)$$

$$\dot{\theta}_i = -D_r \frac{\partial}{\partial \theta_i} U_g^{\text{mod}} + \sqrt{2D_r} \xi_i \quad (13)$$

$$U_g^{\text{mod}} = -g \frac{N|\boldsymbol{\nu}|^2}{D_r} \quad (14)$$



$$I(w) \leq \lim_{\tau \rightarrow \infty} \mathcal{D}_{\text{KL}}(P_{g(w)}^{\text{mod}} || P) \quad (15)$$



F. den Hollander, *Large deviations*, Vol. 14 (American Mathematical Soc., 2008).

R. L. Jack, *The European Physical Journal B* 93, 1–22 (2020).



S. Whitelam et al., *The Journal of Chemical Physics* 148, 154902 (2018).

T. GrandPre et al., *Physical Review E* 103, 012613 (2021).





CM: fluctuations of w_τ strongly coupled to those of \bar{v}_τ



CM: fluctuations of w_τ strongly coupled to those of \bar{v}_τ

CM: response to biasing $s \leftrightarrow$ response to aligning interaction g

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).






- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).
- This transition to CM happens at **finite biasing** $s^* \sim -D_r$ implying that these fluctuations are very distinct from the unbiased behaviour.

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).
- This transition to CM happens at **finite biasing** $s^* \sim -D_r$ implying that these fluctuations are very distinct from the unbiased behaviour.
- It is qualitatively understood through the introduction of **effective aligning interaction**.

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).
- This transition to CM happens at **finite biasing** $s^* \sim -D_r$ implying that these fluctuations are very distinct from the unbiased behaviour.
- It is qualitatively understood through the introduction of **effective aligning interaction**.

PHYSICAL REVIEW E **103**, 022603 (2021)

Collective motion in large deviations of active particles

Yann-Edwin Keta ^{1,2,3} Étienne Fodor ^{1,4} Frédéric van Wijland ² Michael E. Cates ¹ and Robert L. Jack ^{1,5}

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).
- This transition to CM happens at **finite biasing** $s^* \sim -D_r$ implying that these fluctuations are very distinct from the unbiased behaviour.
- It is qualitatively understood through the introduction of **effective aligning interaction**.

PHYSICAL REVIEW E **103**, 022603 (2021)

Collective motion in large deviations of active particles

Yann-Edwin Keta^{1,2,3} Étienne Fodor^{1,4} Frédéric van Wijland² Michael E. Cates¹ and Robert L. Jack^{1,5}

- We compute exactly for a system of **2 particles on a ring** the relation between w_T and particle alignment.

- We have introduced a model of **active Brownian particles** biased with respect to a measure of the **active work** w_T .
- This model of isotropic active matter displays **collective motion** (CM) when w_T is increased ($s < 0$).
- This transition to CM happens at **finite biasing** $s^* \sim -D_r$ implying that these fluctuations are very distinct from the unbiased behaviour.
- It is qualitatively understood through the introduction of **effective aligning interaction**.

PHYSICAL REVIEW E **103**, 022603 (2021)

Collective motion in large deviations of active particles

Yann-Edwin Keta^{1,2,3} Étienne Fodor^{1,4} Frédéric van Wijland^{1,2} Michael E. Cates¹ and Robert L. Jack^{1,5}

- We compute exactly for a system of **2 particles on a ring** the relation between w_T and particle alignment.
- We propose a **fluctuating hydrodynamic theory** which captures the emergence of polar order in the biased state.

Thank you!

- At $\mathbf{P} = 0$, biasing w.r.t. w_τ is equivalent to biasing w.r.t. $|\tilde{\rho}_\mathbf{q}|^2$.

$$S_s(\mathbf{q}) = \langle |\tilde{\rho}_\mathbf{q}|^2 \rangle_s = \begin{cases} \chi_0, & s = 0 \\ b_s q, & s < 0 \end{cases} \quad (16)$$

⇒ We expect hyperuniformity in the isotropic $s < 0$ phase for $N \gg 1$.




Figure: Biased structure factor S_s .

→ Finite system shows suppression of density fluctuations for $s < 0$.