

Lattice Models for Quantum Superalgebras

Colors and Supercolors

Presenters: Joey (Yizhou) Chen, Kenny Deng, Sam Stagner, Phillip Yoon

Collaborators: Hadi Abu-Alfa, Yuekun Feng, Daniel Kumm, Junseo Ko, Aaron Fidelis Lukman,
Ioana Milea, Timothy Niels Ruslim, David Saing, Hanzhang Wang, Arisu

Mentors: Ben Brubaker, Judy Chiang, Karlee Westrem

Polymath Jr 2025 Lattice Model and Representation Theory Group

August 28, 2025

Overview


1. Introduction and Motivations
2. Strategy
3. Results and Conjecture

Lattice Model


Definition

A lattice model \mathcal{L} is an $n \times m$ grid with its edges filled according to vertex table.

- We generally think of them as paths going from the top and exiting to the right
- They are indexed by external edges. Typically, we fix edge colorings at the top row with a partition $\lambda = (\lambda_1, \dots, \lambda_k)$.
- The example below uses this vertex table:



Example. Take $\lambda = (3)$








Partition Functions

Definition (Partition Function)

Given a lattice model with fixed boundary conditions, the partition function $\mathcal{Z}(\mathcal{L})$ of the lattice is the sum of all admissible states, which are paths with non-zero weight.

Note: Non-specified vertices have weight 0.




In this case: $\mathcal{Z}(\mathcal{L}) = x_1^2 + x_1 x_2 + x_2^2$

Boundary Conditions

For models with multiple colors, we may also fix a permutation $w \in S_m$ on the side to indicate the order of colors from top to bottom.

Example. Take $\lambda = (3, 2)$ and $w = (12)$




Super-Lattice Model

Definition


A super-lattice model is a lattice model indexed by one partition λ and two permutations $w, w' \in S_m$ such that it has both colored and dotted colored (supercolor) paths going in opposite directions.

Example. Below is an admissible state with $\lambda = (5, 3, 1)$, $w = (312)$, and $w' = (132)$



Weights for Super-Lattice Model

We are working with these weights [2]:



Research Questions

Motivating Questions

- What do the partition functions of these lattice $\mathcal{L}_{\lambda,w,w'}$ models look like?
- What combinatorial objects represent them?

Past work:

- Previous Polymath projects have "solved" models with one partition λ and one permutation w . [2]

For experts: The weights of the given model were chosen based on quantum superalgebra modules.

- Further Question: How can changing the weights in accordance with these superalgebras affect the partition functions?
- Connected to the supersymmetries between bosons and fermions


Research Strategy

- **Goal.** Compute the partition function of $\mathcal{L}_{\lambda,w,w'}$ for all $\lambda \vdash m$ and $w, w' \in S_n$.
- **First steps.**
 1. Identify w, w' with $\mathcal{L}_{\lambda,w,w'} = 0$.
 2. Identify w, w' such that $\mathcal{L}_{\lambda,w,w'}$ has a unique admissible state.
 3. Compute remaining partition functions recursively by relating permutation index pairs (train argument).
- **Dream.** Understand the quantum group module for super-lattice models

Question (rephrased).

- What is the minimal set of states we need to compute to know all partition functions?
- And what do their partition functions look like?

An Inadmissible (Vanishing) State



Vanishing Conjecture

Vanishing Conjecture

For boundary conditions $(w, w_0 u)$, if $u = w$ then there is only one state and if $u < w$ then there are no states, where $<$ indicates strong Bruhat order and w_0 indicates the longest word.

Strong (full) Bruhat order on S_3 [1]

For $1 \leq i < j \leq 3$ let (ij) be the transposition exchanging i and j . Given $u \in S_3$ we declare

$$u < (ij)u \iff \ell((ij)u) = \ell(u) + 1. \quad (*)$$

The *strong Bruhat order* is the reflexive-transitive closure of this relation $(*)$; i.e. for $u, v \in S_3$

$$u \leq v \iff \text{there exists a chain } u = w_0 < w_1 < \dots < w_k = v \text{ each step satisfying } (*).$$

There are $3! = 6$ elements, which we list by *length* $\ell(w) = \#\{(i < j) \mid w(i) > w(j)\}$ (the number of inversions).

Length Comparison

length	elements
0	$e = 123$
1	$213 = s_1, 132 = s_2$
2	$231 = s_1 s_2, 312 = s_2 s_1$
3	$321 = s_1 s_2 s_1 = s_2 s_1 s_2 = w_0.$

$$e < s_1, s_2,$$

$$s_1 < s_1 s_2, s_2 s_1 < w_0, \quad \text{with } s_1, s_2 \text{ incomparable and likewise } s_1 s_2, s_2 s_1.$$

$$s_2 < s_1 s_2, s_2 s_1 < w_0.$$

Vanishing / one-state table


	w_0	$s_2 s_1$	$s_1 s_2$	s_1	s_2	e
e	1	*	*	*	*	*
s_2	0	1	*	*	*	*
s_1	0	*	1	*	*	*
$s_1 s_2$	0	0	0	1	*	*
$s_2 s_1$	0	0	0	*	1	*
w_0	0	0	0	0	0	1

- 1 exactly one state ($u = w$),
- 0 vanishes ($u < w$ in strong Bruhat order),
- * conjecture does not constrain this pair.

Observations

1. Anti-diagonal of ones.
2. Zeros lie strictly to the left of that anti-diagonal.
3. Row counts reflect the poset.
4. Almost-unitriangular shape.


Train argument for single colored lattice model



Train argument for color/scolor model


Notation:

- w_i represents the **color** decorated at row i under permutation of w , and w'_i represents the **scolor** decorated at row i under permutation of w' .
- z_i and z_j marks the row number before and after the run-through of R -vertex.



Example of train argument on 3×3 Super-lattice model

For simplification, it seems like for a 3×3 lattice model, but one can image the arbitrary rows above and below with arbitrary column in the middle. The lattice models would be, from left to right, $\mathcal{L}_{\lambda, w_0, s_1}$, $\mathcal{L}_{\lambda, w_0, e}$, $\mathcal{L}_{\lambda, s_1 s_2, e}$, $\mathcal{L}_{\lambda, w_0, e}$.



The relations from train argument on the super-lattice model

Case ($w = w_0, w' = e$)

$$q(z_1^3 - z_2^3) Z(\mathcal{L}_{\lambda, w_0, s_1}) + (1 - q^2) z_2^2 z_1 Z(\mathcal{L}_{\lambda, w_0, e}) \\ = s_1 \left[(z_1^3 - z_2^3) Z(\mathcal{L}_{\lambda, s_1 s_2, e}) + (1 - q^2) z_1^3 Z(\mathcal{L}_{\lambda, w_0, e}) \right],$$

$$Z(\mathcal{L}_{\lambda, w_0, s_1}) = 0, \quad Z(\mathcal{L}_{\lambda, w_0, e}) = z_1^{\lambda_1 - 2} z_2^{\lambda_2 - 1} z_3^{\lambda_3}.$$

Upshot

By applying the vanishing partition function value and the already known partition function formula, we are only left with $Z(\mathcal{L}_{\lambda, s_1 s_2, e})$ as a variable of the equation, which means then we can solve for $Z(\mathcal{L}_{\lambda, s_1 s_2, e})$, one of the unknown partition functions according to the table.

Note: s_1 simply means to flip z_1 and z_2 in the context of spectral parameters. For example, if $f = z_1^2 + z_2$, then $s_1 f = z_2^2 + z_1$.

In the context of the table

	w_0	$s_2 s_1$	$s_1 s_2$	s_1	s_2	e
e	1	*	*	*	*	*
s_2	0	1	*	*	*	*
s_1	0	*	1	*	*	*
$s_1 s_2$	0	0	0	1	*	*
$s_2 s_1$	0	0	0	*	1	*
w_0	0	0	0	0	0	1

$Z(\mathcal{L}_{\lambda, s_1 s_2, e})$ corresponds to the circled entry, which is one of the partition functions that we do not know yet (we only know the partition functions corresponding with entries of either 0 or 1).

Gelfand-Tsetlin Patterns

Definition.


A **strict GT-pattern** is a triangular arrangement of non-negative integers

$$\begin{array}{ccccccc}
 x_{n,1} & & x_{n,2} & & \cdots & & x_{n,n} \\
 & \ddots & & \ddots & & \ddots & \\
 & & x_{2,1} & & x_{2,2} & & \\
 & & & x_{1,1} & & &
 \end{array}$$

with the constraint that $x_{i+1,j} \leq x_{i,j} \leq x_{i+1,j+1}$ and $x_{i,j-1} < x_{i,j} < x_{i,j+1}$

Bijection. The numbers in each row record the columns with a color descending path

Example.



Alternating Sign Matrices


Definition.

An **alternate sign matrix** is an $n \times n$ matrix with entries of $-1, 0, 1$ such that each column and row sum to 1, with the non-zero alternating sign entries.

Bijection between super-lattice vertices and ASM entries:

0	0	0	0	-1	1

Example.




Simple Permutation Pairs

Proposition

For $w \in S_n$, $\mathcal{L}_{\lambda, w, w_0 w}$ have a unique non-zero admissible state.

- These lattice models are in bijection with the permutation matrices.

Example.



\Leftrightarrow

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$


$(n - 1) \times n$ and $n \times n$ lattices

Idea.

The color/scolor exiting at the first row of a square lattice model need to be coming from the same column.

Proposition.

Given top boundary λ , the number of admissible states of $n \times n$ lattice model \mathcal{S}_λ equals to the number of admissible states of $(n - 1) \times n$ lattice model $\mathcal{L}_{\lambda'}$ such that λ' has one pair of color and scolor decoration less than λ .



Criteria of non-zero admissible states

We use the previous ideas to make a final conjecture:

$$\left\{ \begin{array}{l} \text{Alternating sign matrix} \implies \text{Uniqueness representation of each path} \\ \text{Simple permutation pairs} \implies \text{Foundation for train argument} \\ (n-1) \times n \text{ and } n \times n \text{ lattice} \implies \text{Enabled analysis on reduced dimension} \end{array} \right.$$

Remark: More specifically, we can take the idea from reducing n rows to $n-1$, and applied it more drastically and inductively, from n rows to lattice i rows with $i = n-1, \dots, 2$; as given a $i \times n$ lattice model with an admissible states, we can always find a $j \times n$ lattice model such that $i \leq j$ which the states is contained by some admissible state of the bigger lattice model.

Thus with this in mind, we want to introduce our last result.

Criteria of non-zero admissible states

Idea: Given any lattice model with a fixed boundary condition (λ, w', w) , we want to instantly justify whether it will or will not have any non-zero admissible state.

Define $s := \prod_{l=1}^k \sigma_l$ and $S_l := \{x \in \mathbb{Z}_n : \sigma_l(x) \neq x\}$, where σ_l are all disjoint permutations in S_n .


Conjecture

Given $\mathcal{L}_{(\lambda, w, w_0 w)}$ has a non-zero admissible state, then $\mathcal{L}_{(\lambda, w, s w_0 w)}$ has a non-zero admissible state if and only if

- (i) $w_0 w >_B s w_0 w$ in strong Bruhat order; and
- (ii) for all l , there exists $y \in S_l$, $y \neq \max(S_l)$ such that $y + 1 \notin S_l$ and $y + 1$ has exited.

Demonstration of the conjecture

	Simple permutation pair	Extra transposition
λ	$(5, 4, 3, 2, 1)$	$(5, 4, 3, 2, 1)$
$w'; sw$	$(\bar{1}\bar{5}\bar{4})(\bar{2}\bar{3}), e(1432)$	$(\bar{1}\bar{5}\bar{4})(\bar{2}\bar{3}), (13)(1432) = (12)(34)$



Recap

Questions.

- What is the partition function of the super-lattice Model?
- Are there any known combinatorial objects in bijection to these lattice models?

Results.

- Special cases with monostate
 - Gelfand-Tsetlin patterns, Alternating Sign Matrices
- Conjecture for non-vanishing states

Next Steps.

- Determine all boundary conditions with unique and multiple states in the square lattice model.
- Find the operator between two arbitrary partition functions for super-lattice models.

References

- [1] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter Groups*. Vol. 231. Graduate Texts in Mathematics. Springer, 2005. DOI: 10.1007/3-540-27596-7.
- [2] Ben Brubaker et al. *Kirillov's conjecture on Hecke-Grothendieck polynomials*. 2024. arXiv: 2410.07960 [math.CO]. URL: <https://arxiv.org/abs/2410.07960>.

Thank you!