

(a) Six resources, six native tasks
ASD-Kinect
ASD / typical
provenance caveat
ATAXIA
Normal / imitated
healthy volunteers
CARE-PD
UPDRS gait score 0-3
Parkinson's cohort
GMFCS
GMFCS I-IV
cerebral-palsy cohort
PARKINSONISM
Mild / severe
Parkinson's cohort
PATHOGAIT
Six gait patterns
healthy, simulated gait
(b) One evidence view per run
Raw coordinates
-0.21 +0.84 +1.37
+0.14 +0.73 +1.39
-0.10 +0.78 +0.86
124-frame source
Normalized coordinates
Center + scale + align
32 frames / 12 joints
Coordinate subset probes
Motion summary
Amplitude / asymmetry
Stability / smoothness
Fixed, label-free template
Named features + z
Supported measurements
Train-only scaling: f → z
Shared columns: LLM / LR / SVM
(c) Is signal recoverable?
Same numerical evidence
Supervised LR / linear SVM
Feature signal
vs. LLM evidence use
Label-prior references anchor chance.
(d) Does evidence matter?
No / shuffled evidence
Same task, altered evidence
A B C D
D C B A
Reverse / shuffle candidate order
Keep the test records and labels fixed.
(e) What does adaptation learn?
LoRA-SFT
All six: in-distribution summary SFT
ATAXIA
PATHOGAIT
Transfer probe: normal vs. abnormal
Six-dataset shared suite: 6 × 7 conditions
Accuracy • fixed-label macro-F1 • κ • per-class / participant checks