Numerical Semigroups, Lattice Ideals, and Markov Bases

Jon Allen

with Trevor McGuire Department of Mathematics North Dakota State University Fargo, ND

Capstone Presentation, North Dakota State University,
December 2015



A numerical semigroup is a nonempty subset S of $\mathbb N$ under addition, which contains the zero element, and whose complement in $\mathbb N$ is finite.

A numerical semigroup is a nonempty subset S of $\mathbb N$ under addition, which contains the zero element, and whose complement in $\mathbb N$ is finite.

It is closed under addition

A numerical semigroup is a nonempty subset S of $\mathbb N$ under addition, which contains the zero element, and whose complement in $\mathbb N$ is finite.

- It is closed under addition
- It is generated from positive (nonzero) integers

A numerical semigroup is a nonempty subset S of $\mathbb N$ under addition, which contains the zero element, and whose complement in $\mathbb N$ is finite.

- It is closed under addition
- It is generated from positive (nonzero) integers
- The greatest common divisor of its generators is 1

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1n_1 + \ldots a_kn_k$ for all $a_i \in \mathbb{N}$.

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1n_1 + \ldots a_kn_k$ for all $a_i \in \mathbb{N}$.

Example

• The numerical semigroup generated by $\{5,7,9\}$ is $\{0,5,7,9,10,12,14,15,16,17,18,\dots\}$

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1n_1 + \ldots a_kn_k$ for all $a_i \in \mathbb{N}$.

- The numerical semigroup generated by $\{5,7,9\}$ is $\{0,5,7,9,10,12,14,15,16,17,18,\dots\}$
- The complement of (5,7,9) in \mathbb{N} is $\{1,2,3,4,6,8,11,13\}$

Dot product

Each element of $\langle 5, 7, 9 \rangle$ is the dot product of the vector (5, 7, 9) and an element of \mathbb{N}^3 .

$$(5,7,9)\cdot(1,0,0)=5$$

$$(5,7,9)\cdot(1,1,0)=12$$

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5,7,9\rangle$.

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5,7,9\rangle$.

Example | 5 7 9 | | 5 1 0 0

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5,7,9\rangle$.

| | 5 | 7 | 9 |
|---|---|---|---|
| 5 | 1 | 0 | 0 |
| 7 | 0 | 1 | 0 |

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5,7,9\rangle$.

Example | 5 7 9 | | 5 1 0 0 | | 7 0 1 0 | | 9 0 0 1 | |

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example | 5 7 9 | | 5 1 0 0 | | 7 0 1 0 | | 9 0 0 1 | | 10 | 2 0 0 | | |

We can make a table where each row is the vector in \mathbb{N}^3 that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example | 5 7 9 | | 5 1 0 0 | | 7 0 1 0 | | 9 0 0 1 | | 10 2 0 0 | | 12 1 1 0

Overview of Numerical Semigroup

Making Markov

Integer Lattice

Smith Normal Form

These vectors are not necessarily unique.

These vectors are not necessarily unique.

| | 5 | 7 | 9 |
|----|---|---|---|
| 5 | 1 | 0 | 0 |
| 7 | 0 | 1 | 0 |
| 9 | 0 | 0 | 1 |
| 10 | 2 | 0 | 0 |
| 12 | 1 | 1 | 0 |
| 14 | 1 | 0 | 1 |
| | | | |

These vectors are not necessarily unique.

| | 5 | 7 | 9 |
|----|---|---|---|
| 5 | 1 | 0 | 0 |
| 7 | 0 | 1 | 0 |
| 9 | 0 | 0 | 1 |
| 10 | 2 | 0 | 0 |
| 12 | 1 | 1 | 0 |
| 14 | 1 | 0 | 1 |
| 14 | 0 | 2 | 0 |

Overview of Numerical Semigroup

Making Markov

Integer Lattice

Smith Normal Form

A fiber is the set of vectors which forms the preimage of each element of our NSG.

A fiber is the set of vectors which forms the preimage of each element of our NSG.

$$\mathcal{F}(12) = \{(1,1,0)\}$$
$$\mathcal{F}(14) = \{(1,0,1), (0,2,0)\}$$

Overview of Numerical Semigroup Making Markov Integer Lattice Smith Normal Form

Fibers can be disconnected or connected.

Fibers can be disconnected or connected.

| Example | | | | | | | | | |
|---------|----|---|---|---|----|---|---|---|--|
| | | 5 | 7 | 9 | | 5 | 7 | 9 | |
| | 5 | 1 | 0 | 0 | 16 | 0 | 1 | 1 | |
| | 7 | 0 | 1 | 0 | 17 | 2 | 1 | 0 | |
| | 9 | 0 | 0 | 1 | 19 | 2 | 0 | 1 | |
| | 10 | 2 | 0 | 0 | 19 | 1 | 2 | 0 | |
| | 12 | 1 | 1 | 0 | 20 | 4 | 0 | 0 | |
| | 14 | 1 | 0 | 1 | 21 | 1 | 1 | 1 | |
| | 14 | 0 | 2 | 0 | 21 | 0 | 3 | 0 | |
| | 15 | 3 | 0 | 0 | 22 | 3 | 1 | 0 | |

It is useful to think of a fiber as a graph.

$$\mathcal{F}(5) = \{(1,0,0)\}$$

$$\mathcal{F}(12) = \{(1,1,0)\}$$

$$\mathcal{F}(14) = \{(1,0,1), (0,2,0)\} \quad \mathcal{F}(19) = \{(2,0,1), (1,2,0)\}$$

$$\boxed{5} \quad \boxed{7}$$

$$\boxed{9} \quad \boxed{9}$$

• Moves happen when fibers are disconnected.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.
- Moves are the difference between disconnected elements of fibers.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.
- Moves are the difference between disconnected elements of fibers.

| | 5 | 7 | 9 | Markov Basis | | | | |
|----|---|---|---|--------------|----|----|--|--|
| 14 | 1 | 0 | 1 | | | | | |
| 14 | 0 | 2 | 0 | 1 | -2 | 1 | | |
| 25 | 5 | 0 | 0 | | | | | |
| 25 | 0 | 1 | 2 | 5 | -1 | -2 | | |
| 27 | 0 | 0 | 3 | | | | | |
| 27 | 4 | 1 | 0 | -4 | -1 | 3 | | |

Overview of Numerical Semigroup Making Markov Integer Lattice Smith Normal Form

We have an easy bijection between our Markov basis and a lattice ideal.

We have an easy bijection between our Markov basis and a lattice ideal.

$$\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \Leftrightarrow \begin{cases} xz - y^2 \\ x^5 - yz^2 \\ z^3 - x^4y \end{cases}$$

 We have actually explicitly built our Markov basis to be the null space of the numerical semigroup basis.

- We have actually explicitly built our Markov basis to be the null space of the numerical semigroup basis.
- If we can find some vector \vec{x} such that

$$\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \vec{x} = \vec{0}$$

Then we will have found our semigroup!

What we need is the Smith Normal Form.

$$UAV = B = \left[\begin{array}{ccc} b_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & b_r \end{array} \right]$$

We start with identity matrices on either side of our Markov matrix. The procedure is similar to finding an inverse matrix, (except the Markov matrix is singular).

We start with identity matrices on either side of our Markov matrix. The procedure is similar to finding an inverse matrix, (except the Markov matrix is singular).

We reduce our Markov matrix, mirroring column and row operations in the adjacent matrices.

We start with identity matrices on either side of our Markov matrix.

The procedure is similar to finding an inverse matrix, (except the Markov matrix is singular).

We reduce our Markov matrix, mirroring column and row operations in the adjacent matrices.

We can't use anything but integers for our row and column operations!

$$\left[\begin{array}{cccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array}\right]
\left[\begin{array}{cccc}
1 & -2 & 1 \\
5 & -1 & -2 \\
-4 & -1 & 3
\end{array}\right]
\left[\begin{array}{cccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array}\right]$$

Row operations on the left

$$\left[\begin{array}{ccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array}\right]
\left[\begin{array}{ccc}
1 & -2 & 1 \\
5 & -1 & -2 \\
-4 & -1 & 3
\end{array}\right]
\left[\begin{array}{ccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array}\right]$$

Row operations on the left

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & -9 & 7 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Row operations on the left

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & -9 & 7 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$


$$\left[\begin{array}{ccc}
1 & 0 & 0 \\
-5 & 1 & 0 \\
-1 & 1 & 1
\end{array}\right]
\left[\begin{array}{ccc}
1 & 0 & 0 \\
0 & 9 & 2 \\
0 & 0 & 0
\end{array}\right]
\left[\begin{array}{ccc}
1 & 2 & 1 \\
0 & 1 & 1 \\
0 & 0 & 1
\end{array}\right]$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & -3 & 1 \\ 0 & -4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & -3 & 1 \\ 0 & -4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 5 \\ 0 & -3 & 7 \\ 0 & -4 & 9 \end{bmatrix}$$



Thank You! Questions?