Numerical Semigroups, Lattice Ideals, and Markov Bases

Jon Allen

with Trevor McGuire
Department of Mathematics
North Dakota State University
Fargo, ND

Capstone Presentation, North Dakota State University, December 2015

Overview

A numerical semigroup is a nonempty subset S of $\mathbb{N}$ under addition, which contains the zero element, and whose complement in $\mathbb{N}$ is finite.

Overview

A numerical semigroup is a nonempty subset S of $\mathbb{N}$ under addition, which contains the zero element, and whose complement in $\mathbb{N}$ is finite.

- It is closed under addition

Overview

A numerical semigroup is a nonempty subset S of $\mathbb{N}$ under addition, which contains the zero element, and whose complement in $\mathbb{N}$ is finite.

- It is closed under addition
- It is generated from positive (nonzero) integers

Overview

A numerical semigroup is a nonempty subset S of $\mathbb{N}$ under addition, which contains the zero element, and whose complement in $\mathbb{N}$ is finite.

- It is closed under addition
- It is generated from positive (nonzero) integers
- The greatest common divisor of its generators is 1

Example

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1 n_1 + \ldots a_k n_k$ for all $a_i \in \mathbb{N}$.

Example

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1 n_1 + \ldots a_k n_k$ for all $a_i \in \mathbb{N}$.

Example

- *The numerical semigroup generated by $\{5, 7, 9\}$ is $\{0, 5, 7, 9, 10, 12, 14, 15, 16, 17, 18, \ldots\}$*

Example

Let S be the numerical semigroup generated by $\{n_1, \ldots, n_k\}$ with $n_i \in \mathbb{N} \setminus \{0\}$. Then the elements of S are $a_1 n_1 + \ldots a_k n_k$ for all $a_i \in \mathbb{N}$.

Example

- *The numerical semigroup generated by $\{5, 7, 9\}$ is $\{0, 5, 7, 9, 10, 12, 14, 15, 16, 17, 18, \ldots\}$*
- *The complement of $\langle 5, 7, 9 \rangle$ in $\mathbb{N}$ is $\{1, 2, 3, 4, 6, 8, 11, 13\}$*

Dot product

Each element of $\langle 5, 7, 9 \rangle$ is the dot product of the vector $(5, 7, 9)$ and an element of $\mathbb{N}^3$.

Example

$$(5, 7, 9) \cdot (1, 0, 0) = 5$$
$$(5, 7, 9) \cdot (1, 1, 0) = 12$$

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example

	5	7	9
5	1	0	0

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example

	5	7	9
5	1	0	0
7	0	1	0

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example

	5	7	9
5	1	0	0
7	0	1	0
9	0	0	1

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example

	5	7	9
5	1	0	0
7	0	1	0
9	0	0	1
10	2	0	0

Table

We can make a table where each row is the vector in $\mathbb{N}^3$ that corresponds to an element in $\langle 5, 7, 9 \rangle$.

Example

	5	7	9
5	1	0	0
7	0	1	0
9	0	0	1
10	2	0	0
12	1	1	0

These vectors are not necessarily unique.

These vectors are not necessarily unique.

Example

	5	7	9
5	1	0	0
7	0	1	0
9	0	0	1
10	2	0	0
12	1	1	0
14	1	0	1

These vectors are not necessarily unique.

Example

	5	7	9
5	1	0	0
7	0	1	0
9	0	0	1
10	2	0	0
12	1	1	0
14	1	0	1
14	0	2	0

A fiber is the set of vectors which forms the preimage of each element of our NSG.

A fiber is the set of vectors which forms the preimage of each element of our NSG.

Example

$$\mathcal{F}(12) = \{(1,1,0)\}$$
$$\mathcal{F}(14) = \{(1,0,1),(0,2,0)\}$$

Fibers can be disconnected or connected.

Fibers can be disconnected or connected.

Example

	5	7	9		5	7	9
5	1	0	0	16	0	1	1
7	0	1	0	17	2	1	0
9	0	0	1	19	2	0	1
10	2	0	0	19	1	2	0
12	1	1	0	20	4	0	0
14	1	0	1	21	1	1	1
14	0	2	0	21	0	3	0
15	3	0	0	22	3	1	0

It is useful to think of a fiber as a graph.

Example

$$\mathcal{F}(5) = \{(1,0,0)\} \qquad\qquad \mathcal{F}(12) = \{(1,1,0)\}$$

$$\mathcal{F}(14) = \{(1,0,1),(0,2,0)\} \quad \mathcal{F}(19) = \{(2,0,1),(1,2,0)\}$$

- Moves happen when fibers are disconnected.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.
- Moves are the difference between disconnected elements of fibers.

- Moves happen when fibers are disconnected.
- Moves are the elements of the Markov basis.
- Moves are the difference between disconnected elements of fibers.

Example

	5	7	9	Markov Basis		
14	1	0	1			
14	0	2	0	1	-2	1
25	5	0	0			
25	0	1	2	5	-1	-2
27	0	0	3			
27	4	1	0	-4	-1	3

We have an easy map between our Markov basis and a lattice ideal.

We have an easy map between our Markov basis and a lattice ideal.

Example

$$\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \Leftrightarrow \begin{cases} xz - y^2 \\ x^5 - yz^2 \\ z^3 - x^4y \end{cases}$$

- We have actually explicitly built our Markov basis to be the null space of the numerical semigroup basis.

- We have actually explicitly built our Markov basis to be the null space of the numerical semigroup basis.
- If we can find some vector $\vec{x}$ such that

$$\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \vec{x} = \vec{0}$$

Then we will have found our semigroup!

What we need is the Smith Normal Form.

$$UAV = B = \begin{bmatrix} b_1 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & b_r \end{bmatrix}$$

We start with identity matrices on either side of our Markov matrix.
The procedure is similar to finding an inverse matrix, (except the
Markov matrix is singular).

We start with identity matrices on either side of our Markov matrix.
The procedure is similar to finding an inverse matrix, (except the
Markov matrix is singular).
We reduce our Markov matrix, mirroring column and row
operations in the adjacent matrices.

We start with identity matrices on either side of our Markov matrix.
The procedure is similar to finding an inverse matrix, (except the
Markov matrix is singular).
We reduce our Markov matrix, mirroring column and row
operations in the adjacent matrices.
We can't use anything but integers for our row and column
operations!

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Row operations on the left

$$
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix}
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
$$

Row operations on the left

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right] \left[\begin{array}{ccc} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{array}\right] \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right]$$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ -5 & 1 & 0 \\ 4 & 0 & 1 \end{array}\right] \left[\begin{array}{ccc} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & -9 & 7 \end{array}\right] \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right]$$

Row operations on the left

$$
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
\begin{bmatrix} 1 & -2 & 1 \\ 5 & -1 & -2 \\ -4 & -1 & 3 \end{bmatrix}
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
$$

$$
\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix}
\begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & -9 & 7 \end{bmatrix}
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
$$

$$
\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix}
\begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix}
\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}
$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 9 & -7 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & -3 & 1 \\ 0 & -4 & 1 \end{bmatrix}$$

Column operations on the right

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 9 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & -3 & 1 \\ 0 & -4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ -5 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 5 \\ 0 & -3 & 7 \\ 0 & -4 & 9 \end{bmatrix}$$

Thank You!
Questions?