

Notes

January 16, 2015

$$\begin{aligned} \text{irrationals} &\subseteq \bigcup_{i=1}^{\infty} (-i, i) = \mathbb{R} \\ \{0\} &\subseteq \end{aligned}$$

outer measure of a sum

$$E \subseteq \mathbb{R} \text{ then } m^*(E) = \inf \left\{ \sum_{i=1}^{\infty} b_i - a_i : E \subseteq \bigcup_{i=1}^{\infty} (a_i, b_i) \right\}$$

example

$$m^*(\{x\}) = 0$$

$$\{x\} \subseteq (x - \epsilon, x + \epsilon) \text{ for any } \epsilon > 0, m^*(t) \leq 2\epsilon \text{ for any } \epsilon > 0 \rightarrow 0$$

1. note that $m^*(E) \subseteq [0, \infty]$, no negative

2. the empty set has measure zero

$$m^*(\emptyset) = 0$$

$$\emptyset \subseteq \bigcup_{i=1}^{\infty} (a_i, b_i)$$

3. $A \subseteq B \Rightarrow m^*(A) \leq m^*(B)$

proof

$$\text{if } B \subseteq \bigcup_{i=1}^{\infty} (a_i, b_i) \text{ then } A \subseteq \bigcup_{i=1}^{\infty} (a_i, b_i)$$

$$\inf \{ \sum b_i - a_i \mid A \subseteq \bigcup (a_i, b_i) \} \leq \inf \{ \sum c_i - d_i \mid B \subseteq \bigcup (c_i, d_i) \}$$

cant do this with contradiction, that's the "actual way"

example

$$m^*([a, b]) = b - a?$$

$$\inf \{ \sum (b_i - a_i) : [a, b] \subseteq \bigcup (a_i, b_i) \}$$

proof

$$[a, b] \subseteq (a - \epsilon, b + \epsilon) \text{ for all } \epsilon > 0$$

$$\begin{aligned} m^*([a, b]) &\subseteq (b + \epsilon) - (a - \epsilon) \text{ for all } \epsilon > 0 \\ &\subseteq (b - a) + 2\epsilon \end{aligned}$$

so $m * ([a, b]) \subseteq b - a$

heine-borel theorem (HBT) compact set is closed and bounded

compact: every open cover has finite subcovers? check this

if $[a, b] \subseteq \bigcup (a_i, b_i)$

wlog $[a, b] \subseteq \bigcup^n (a_i, b_i)$

$$a_1 < a$$

$$a_2 < b_1$$

$$a_3 < b_2$$

$$a_4 < b_3$$

$$\vdots a_n$$

$$b < b_n$$

$$< b_{n-1}$$

these are overlapping covers, they make a sequence, it is important that they overlap

sum to infinity is bigger than sum to n . $\sum (b_1 - a_i) + (b_2 - a_2) + \dots + (b_n - a_n) \geq (a_2 - a_1) + (a_3 - a_2) \dots (a_n - a_{n-1}) + (b_n - a_n) = b_n - a_1 \geq b - a$

and so $m * ([a, b]) \geq b - a$ and we already have less than so it's equal

example 2

$m * ((a, b)) = b - a$ second part of limhof? thm, uniqueness

notice that the different sets are the same size

example 3

$m * ([a, \infty)) = \infty$ notice that the measure of any subset is greater than or equal and so $[a, a+k] \subseteq [a, \infty)$ for all k and so $k = m * [a, a+k] \leq m * [a, \infty) \forall k$

example 4,5

$m * \mathbb{Q}, m * \mathbb{C} - \mathbb{R}$

$$4. m * (\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} m * (A_i)$$

proof

let $\epsilon > 0$ then for all i there is $\{(a_i^j, b_i^j)\}_{j=1}^{\infty}$ such that $m * A_i \leq \sum b_i^j - a_i^j \leq m * A_i + \frac{\epsilon}{2^j}$

notice that $\{\{a_i^j, b_i^j\}_{i=1}^{\infty}\}_{j=1}^{\infty}$ is countable collection of intervals with $\bigcup A_j \subseteq \bigcup_{j=1}^{\infty} (\bigcup_{i=1}^{\infty} (a_i^j, b_i^j))$ $m * A_j \leq \sum_{j=1}^{\infty} (\sum_{i=1}^{\infty} (a_i^j, b_i^j))$

this is called countable subadditivity