

# ***Coronary Obstruction Following Transcatheter Aortic Valve Replacement Using CoreValve or Evolut in Native Aortic Stenosis: A Systematic Review***

Aoi Omori, MD

Department of Cardiovascular Medicine,  
*Kobe City Medical Center General Hospital*

---



**TCT**<sup>®</sup>

TRANSCATHETER  
CARDIOVASCULAR  
THERAPEUTICS<sup>®</sup>

# Disclosure of Relevant Financial Relationships

Drs. Taniguchi, Hayashi, Ishizu, and Tabata have received lecture fee from Edwards Lifesciences, Medtronic, and Abbott.

Shinichi Shirai is a proctor of Edwards Lifesciences, Medtronic, and Abbott. Hiroshi Takiguchi has been an employee of Eli Lilly Japan since April 1, 2024.

I, Toshiaki Toyota, Kenji Ando, and Yutaka Furukawa do not have any financial relationships to disclose.

# Introduction

- ✓ Coronary obstruction following transcatheter aortic valve replacement (TAVR) in native aortic stenosis is a rare ( $< 1.0\%$ ) but fatal complication with high mortality.

Ribeiro HB et al. J Am Coll Cardiol. 2013 22;62:1552-62.

Jabbour RJ et al. J Am Coll Cardiol. 2018 10;71:1513-1524.


- ✓ The mechanisms and managements about coronary obstruction following TAVR remain unclear, especially with CoreValve or Evolut.

# Methods

## ✓ *a comprehensive systematic review*

reports about coronary obstruction following TAVR (CoreValve or Evolut)  
published from Mar. 2007 to Jan. 2025

## ✓ *Classification*



## ✓ *Data analysis*

baseline data: clinical, echocardiographic, CT characteristics


procedural data: valve type/size, clinical presentation, management

Jabbour RJ et al. J Am Coll Cardiol. 2018 10;71:1513-1524.

# Results: inclusion

✓ Out of 2601 reports, **33** reports (**39** patients) were included.

exclusion : other valves, valve-in-valve, prophylactic wire or stenting, obstruction due to embolism or migration, etc.



Followed the PRISMA 2020 statement

BMJ 2021;372:n71

# Results: characteristics and management

	Number of available data	Overall (N=39)	Acute or early obstruction (N=19)	Late obstruction (N=20)	P value
<u>Clinical characteristics</u>					
Age, years	37	82.6 ± 6.4	83.3 ± 7.5	82.1 ± 5.5	0.58
<b>Female</b>	37	<b>34 (92%)</b>	<b>14 (82%)</b>	<b>20 (100%)</b>	0.050
<u>CT characteristics</u>					
Perimeter	28	68.6 ± 5.7	69.6 ± 6.8	67.3 ± 3.5	0.28
Obstructed coronary artery height <12 mm	30	17 (57%)	10 (63%)	7 (50%)	0.49
<b>Mean SOV diameter below the manufacturer's recommended threshold</b>	24	<b>10 (42%)</b>	<b>4 (40%)</b>	<b>6 (43%)</b>	0.89
<u>Valve size</u>					
	39				
23 mm		1 (3%)	0	1 (5%)	-
<b>26 mm</b>		<b>32 (82%)</b>	<b>14 (74%)</b>	<b>18 (90%)</b>	-
29 mm		6 (15%)	5 (26%)	1 (5%)	-
31 mm CoreValve		0	0	0	-
34 mm Evolut		0	0	0	-

✓ **Females and the 26-mm valve** were predominant.

✓ In **40%** of the cases, the mean SOV diameter was below the threshold.

# Results: characteristics and management

	Number of available data	Overall (N=39)	Acute or early obstruction (N=19)	Late obstruction (N=20)	P value
<b><u>Timing</u></b>	39				
Acute		11 (28%)	11 (58%)	-	-
Early		8 (21%)	8 (21%)	-	
Late		20 (51%)	-	20 (51%)	-
<b><u>Clinical presentation</u></b>					
<b>ACS</b>	39	<b>32 (82%)</b>	<b>18 (95%)</b>	<b>14 (70%)</b>	0.04
<b>Culprit artery</b>					
<b>LCA</b>	39	<b>26 (67%)</b>	<b>15 (79%)</b>	<b>11 (55%)</b>	
RCA	39	6 (15%)	3 (16%)	3 (15%)	0.12
Both	39	7 (18%)	1 (5%)	6 (30%)	
<b><u>Treatment</u></b>					
Emergency treatment	39	32 (82%)	19 (100%)	13 (65%)	0.004
Percutaneous intervention (coronary stenting or snaring)	39	28 (72%)	<b>17 (89%)</b>	<b>11 (55%)</b>	0.02
<b>Unsuccessful PCI</b>	30	7 (23%)	2 (13%)	<b>5 (33%)</b>	0.20
Surgical intervention	39	11 (28%)	2 (11%)	<b>9 (45%)</b>	0.02

- ✓ **ACS** was more common in acute or early and **LCA** obstruction was predominant.
- ✓ In late cases, **unsuccessful PCI** tended to be more frequent .


# Results: etiologies

## Acute or early


*Displacement of  
calcified native valve leaflet*




*Commissural misalignment*



*female*




*Narrow SOV*




*Low coronary  
height*

## late

*High implantation*



*Endothelialization*





# From this systematic review

- ✓ **The 26-mm valve** was most frequently implicated (82%).
  - ➔ This suggests a device-specific relationship with coronary obstruction.
- ✓ The most cases were **female** (92%), and presented as **ACS** (82%).
  - ➔ Females generally have smaller anatomical features.

Jabbour RJ et al. J Am Coll Cardiol. 2018 10;71:1513-1524.  
Ribeiro HB et al. Eur Heart J. 2018;39:687-695.

Coronary obstruction after TAVR is life-threatening, yet PCI is sometimes unsuccessful.

- ✓ The mechanisms of coronary obstruction varied depending on **timing**.

# Why 26-mm ?

Q. Is the 26-mm valve most frequently implanted ?

A. Maybe 'No'.

## Three Generations of Self-Expanding Transcatheter Aortic Valves


A Report From the STS/ACC TVT Registry

John K. Forrest, MD,<sup>a</sup> Ryan K. Kaple, MD,<sup>a</sup> Gilbert H.L. Tang, MD, MSc, MBA,<sup>b</sup> Steven J. Yakubov, MD,<sup>c</sup>  
Tamim M. Nazif, MD,<sup>d</sup> Mathew R. Williams, MD,<sup>a</sup> Angie Zhang, MS,<sup>f</sup> Jeffrey J. Popma, MD,<sup>g</sup> Michael J. Reardon, MD<sup>h</sup>

	CoreValve (n = 5,514)	Evolut R (n = 11,295)	Evolut PRO (n = 2,065)	p Value*
Valve size implanted				
23 mm	271 (5.0)	444 (4.0)	79 (3.9)	0.82
26 mm	1,874 (34.5)	3,869 (34.5)	695 (33.9)	0.61
29 mm	3,294 (60.6)	6,913 (61.6)	1,277 (62.3)	0.56

39 cases in this study


15



10



14



The specific structural feature of the 26-mm valve may be the key factor.

# The 26-mm's feature and multiple factors

CoreValve  
Evolut

23 mm

26 mm

29 mm

34 mm (31mm\*)



Acute (intraprocedural  
or Early ( $\leq 7$  days)

0

14 cases

5 cases

0


Late ( $> 7$  days)

1 case

18 cases

1 case

0



The 26-mm was cylindrical shape  
in the lower half

**anatomical**

**procedural**

*narrow SOV*

*low coronary height*

*displacement the  
calcified valve*

*high implantation*

*commissural  
misalignment*

*small gap between  
the valve and SOV/STJ*

**coronary obstruction**

# Limitations

- ✓ This study is descriptive and cannot robustly assess the incidence of coronary obstruction or conclude a higher obstruction risk with 26-mm valves.
- ✓ Publication bias may overestimate procedural success.
- ✓ Incomplete CT data limited the assessment of the relationship between the implanted valve size and SOV diameter.

# Simultaneously published in *JACC: Advances*



Journal Pre-proof



Coronary Obstruction Following Transcatheter Aortic Valve Replacement With Supra-Annular Self-Expanding Valve: Review of Case Reports

Tomohiko Taniguchi, MD, PhD, Aoi Omori, MD, Toshiaki Toyota, MD, PhD, Shinichi Shirai, MD, Masaomi Hayashi, MD, Hiroshi Takiguchi, MD, PhD, Kenichi Ishizu, MD, Hiroyuki Tabata, MD, Kenji Ando, MD, Yutaka Furukawa, MD, PhD

PII: S2772-963X(25)00745-8

DOI: <https://doi.org/10.1016/j.jacadv.2025.102316>

Reference: JACADV 102316

To appear in: *JACC: Advances*

Received Date: 23 July 2025

Revised Date: 23 September 2025

Accepted Date: 3 October 2025

Please cite this article as: Taniguchi T, Omori A, Toyota T, Shirai S, Hayashi M, Takiguchi H, Ishizu K, Tabata H, Ando K, Furukawa Y, Coronary Obstruction Following Transcatheter Aortic Valve Replacement With Supra-Annular Self-Expanding Valve: Review of Case Reports, *JACC: Advances* (2025), doi: <https://doi.org/10.1016/j.jacadv.2025.102316>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation.

# Conclusions

- ✓ Coronary obstruction following TAVR with CoreValve/Evolut occurred most frequently with the 26-mm valve.  
The cylindrical shape may be associated with this fatal complication.
- ✓ Coronary obstruction was influenced by both anatomical and procedural factors. The mechanisms and managements differed according to timing.