

An Algorithmic Approach to Managing Catastrophic Hypotension During TAVR

Jamie McCabe

Section Head, Structural Heart
Beth Israel Deaconess Medical Center



Beth Israel Lahey Health 

Beth Israel Deaconess
Medical Center



HARVARD MEDICAL SCHOOL
TEACHING HOSPITAL



Disclosures

Honoraria / Consulting / Ad Boards:

- Abbott
- Edwards
- Medtronic
- Jena Valve
- Gore

Equity:


- Excision Medical
- ConKay Medical
- SESAME LLC
- Arcos Medical
- Sparrow Medical
- Nimble Surgical



Timing is everything



I. Sedation




Differential	Initial Steps
<ol style="list-style-type: none">1. Med-induced hypotension2. Hypoxia	<ol style="list-style-type: none">1. Call for anesthesia if RN-led sedation2. Anticipate intubation3. Consider ECMO candidacy if hypotension not imminently reversible



Timing is everything



2. Access




Differential	Initial Steps
<ol style="list-style-type: none">1. Access bleeding2. Vessel disruption / perforation	<ol style="list-style-type: none">1. Peripheral angio from alternative access2. Call for peripheral equipment & covered stents to be in the room



Timing is everything



3. Pacing




Differential	Initial Steps
<ol style="list-style-type: none">1. Effusion from RV perf2. Pacing for too long leading to LV spiral3. Venous bleeding (typically manifests later)	<ol style="list-style-type: none">1. Stop pacing!2. Call for echo



Timing is everything



4. Aortography / Angles




Differential	Initial Steps
<ol style="list-style-type: none">1. Contrast reaction2. Growing effusion3. Coronary embolus / air	<ol style="list-style-type: none">1. Call for echo2. Check skin for rash3. Quick check of tele



Timing is everything



5. Crossing Valve




Differential	Initial Steps
<ol style="list-style-type: none">1. Pinning aortic leaflet open / severe AR2. LV wire perf3. Severe MR from wire entanglement4. Occult effusion (temp wire / RV)5. Carabello's sign	<ol style="list-style-type: none">1. Review hemos for suggestion of AR2. Take tension off wire / catheter system3. Call for echo



Timing is everything



6. Valve Deployment




Differential	Initial Steps
<ol style="list-style-type: none">1. Effusion until proven otherwise following BEV or pre- / post-dilation SEV2. Coronary obstruction3. Valve malplacement / migration4. Complete heart block5. Acute stroke6. Aortic dissection7. Valve upside down (incorrect load)8. Contrast / protamine reaction	<ol style="list-style-type: none">1. Call for echo2. Aortogram to look for coronary obstruction or valve migration3. Review telemetry for CHB, ST Elevation, VT/VF vs PEA4. Consider ECMO candidacy



Timing is everything



7. Access Closure





Differential	Initial Steps
<ol style="list-style-type: none">1. Massive access bleeding2. RP Bleed no longer covered by sheath3. Growing effusion4. Vagal5. Contrast / protamine reaction	<ol style="list-style-type: none">1. Hold pressure if overt bleeding2. Peripheral angio3. Call for Coda / peripheral balloon equipment4. Consider quick follow up echo5. ECMO is <i>NOT</i> a solution for bleeding!



Conclusions



- Hypotension timing is fundamental to considering mechanism, diagnostics and management
- In almost every circumstance consider pericardial effusion, LV dysfunction & bleeding / vascular injury as the 3 most likely culprits and assess accordingly

  @jamiemccabeMD
@jamiemccabe.bsky.social
jmccabe4@bidmc.Harvard.edu

