

Ultra Low Contrast Computed Tomography (CT) for Transcatheter Aortic Valve Replacement (TAVR) Planning

Lesley Field MPH, PA-C

Structural Heart: Providence Santa Rosa Memorial Hospital



Acknowledgements

Robert Feiwell, MD and Vishal Patel, MD

Disclosure of Relevant Financial Relationships

I, Lesley Field DO NOT have any financial relationships to disclose over the past 24 months.

Severe aortic stenosis and CKD

High risk subset of patients at increased risk of morbidity and mortality post TAVR

Major risk factor:

Exposure to a significant dose of iodinated contrast (iohexol) during TAVR CT evaluation

Can result in acute kidney injury from contrast induced nephropathy which for some patients means ESRD requiring dialysis

Current Strategies to Reduce Contrast Nephropathy During TAVR Workup for Patients with CKD

Pre and Post procedure hydration with normal saline

- Difficult in patients with tenuous volume status
- Additional resources required
- Insufficient protection in many cases

Use of 3D Transesophageal echocardiography for Sizing

- Likely requires general anesthesia
- Not as reliable/reproducible as CT
- Does not evaluate vascular access

Current Strategies to Reduce Contrast Nephropathy During TAVR Workup for Patients with CKD

Stage CT Scan, Catheterization and TAVR procedures

- Can lengthen time from work-up to procedure
- May not be possible in hospitalized patients with acute aortic valve syndrome (acute CHF, syncope, arrest)

Non-Contrast CT Scan

- Annulus sizing less accurate/reproducible

Low Dose Contrast Pilot Study

- Collaboration with our chief of radiology to help with the technical aspects of CT, as well as helping to determine contrast administration and timing details
- 9 patients were selected for either low dose gadobutrol or low dose iohexol between the dates of 11/08/2024 and 06/06/2025
- Images were sent to the structural heart interventional cardiologist and the aortic valve company for 3 mensio analysis to determine diagnostic image quality and valve selection

Low dose gadolinium (gadobutrol) vs ultra low dose iodine (iohexol) TAVR CT

Low dose gadobutrol

- Initial 10cc dose iohexol for timing
- Timing of gadobutrol injection achieved using a cursor on ascending aorta with 3 second delay for scan onset + timing of early rise of contrast density in ascending aorta
- 20cc gadobutrol administered via right arm IV, CT set 100kV

Ultra low dose iodine

- Initial 10cc dose iohexol for timing
- Timing of second iohexol injection achieved using a cursor on ascending aorta with 3 second delay for scan onset + timing of early rise of contrast density in ascending aorta
- 20cc iohexol administered via right arm IV, CT set 100kV

Image quality and Impact on Renal Function

- Image quality determined by measuring Hounsfield units at the aortic annulus for both protocols
 - Both protocols provided diagnostic pre TAVR CT images
 - Low dose iohexol generated better enhancement of the aortic annulus
- Renal function: pre and post CT GFRs were obtained for all patients
 - None of the patients experienced a significant change in GFR post CT with either protocol

Low Dose Iohexol vs Gadobutrol Protocol Results

Patient Number	Protocol	Annulus Hounsfield Units	Pre-CT GFR	Post CT GFR
1	Gadobutrol+Iohexol	106	23	27
2	Gadobutrol+Iohexol	62	18	18
3	Gadobutrol+Iohexol	116	28	29
4	Gadobutrol+Iohexol	87	12	21
5	Iohexol Only	267	60	60
6	Iohexol Only	119	34	36
7	Iohexol Only	130	38	40
8	Iohexol Only	225	35	33
9	Iohexol Only	254	22	20


Standard Full Dose Iohexol TAVR CT



Full Dose Iohexol vs Low Dose Iohexol TAVR CT



Full dose iohexol 130cc



Low dose iohexol 30cc

Low Dose Iohexol vs Low Dose Gadobutrol TAVR CT



Low dose iohexol



Low dose gadobutrol

Take-home Message

- Patients with CKD can undergo successful TAVR CT utilizing a total dose of 30 cc of Iohexol
- This pilot study of 9 patients showed no evidence of a change in GFR after ultra-low dose iodine contrast CT

References

Cavallo, A. U., Patterson, A. J., Thomas, R., Alaiti, M. A., Attizzani, G. F., Laukamp, K., Große Hokamp, N., Bezerra, H., Gilkeson, R., & Rajagopalan, S. (2020). Low dose contrast CT for transcatheter aortic valve replacement assessment: Results from the prospective SPECTACULAR study (spectral CT assessment prior to TAVR). *Journal of Cardiovascular Computed Tomography*, 14(1), 68–74. <https://doi.org/10.1016/j.jcct.2019.06.015>

Cheong, B. Y. C., Wilson, J. M., Preventza, O. A., & Muthupillai, R. (2022). Gadolinium-Based Contrast Agents: Updates and Answers to Typical Questions Regarding Gadolinium Use. *Texas Heart Institute Journal*, 49(3). <https://doi.org/10.14503/thij-21-7680>

Chmielewski, M., Serafin, Z., Dorota Kamińska, Katarzyna Skrobisz, Kozak, O., Piotr Olczyk, Rutkowski, P., Adamczak, M., Edyta Szurowska, & Krajewska, M. (2024). The use of intravascular contrast media in patients with impaired kidney function – joint clinical practice position statement of the Polish Society of Nephrology and the Polish Medical Society of Radiology. *Polish Journal of Radiology*, 89(89), 161–171. <https://doi.org/10.5114/pjr.2024.136950>

Devarakonda, P., reddy, K., Gotlieb, V., & Dym, R. (2022). Can Gadolinium Be Used As An Alternative To Iodinated Contrast For CT Angiography During The Current Pandemic-Related Contrast Shortage? *Journal of Medical Case Reports and Case Series*, 5(5). <https://doi.org/10.38207/jmcrcs/2022/sep03090368>

Elison, D., Danek, B. A., Johnson, B. V., Chung, C. J., Oyetunji, S., Mackensen, G. B., Aldea, G., & McCabe, J. M. (2024). A Novel Protocol for Contrast-Free Transcatheter Aortic Valve Replacement Evaluation and Implantation. *Journal of the Society for Cardiovascular Angiography & Interventions*, 3(9), 102237. <https://doi.org/10.1016/j.jscvi.2024.102237>

Lacy, S. C., Benjamin, M. M., Osman, M., Syed, M. A., & Kinno, M. (2023). Low Contrast and Low kV CTA Before Transcatheter Aortic Valve Replacement: A Systematic Review. *Journal of Cardiovascular Imaging*, 31(2), 108. <https://doi.org/10.4250/jcvi.2022.0108>

References

Lippincott, E., Goel, K., Lima, B., Henry, C., Williams, J., Barker, C., Lindman, B., & Carr, J. (2024). Feasibility Of Gadolinium Enhanced Cardiac CT For TAVR Planning In Patients With Contraindication To Iodinated Contrast. *Journal of Cardiovascular Computed Tomography*, 18(4), S92–S92. <https://doi.org/10.1016/j.jcct.2024.05.215>

Oechsner, T., Soschynski, M., Schlett, C. L., Krauss, T., Schupppert, C., Müller-Peltzer, K., Milán Vecsey-Nagy, Kravchenko, D., Ákos Varga-Szemes, Emrich, T., Scheu, R., Taron, J., Bamberg, F., & Hagar, M. T. (2025). Feasibility of very low iodine dose aortoiliac CT angiography using dual-source photon-counting detector CT. *European Journal of Radiology*, 183, 111919–111919. <https://doi.org/10.1016/j.ejrad.2025.111919>