

Cardiac Damage Staging in Moderate or Greater Aortic Regurgitation



Cleveland Clinic

A New Framework for Risk Stratification

Francisco Alexandrino, MD



TCT[®]

TRANSCATHETER
CARDIOVASCULAR
THERAPEUTICS[®]

Accepted to *Structural Heart*

Disclosure of Relevant Financial Relationships

I, Francisco Alexandrino, DO NOT have any financial relationships to disclose.

How do we assess RISK in AR?

- *Progressive volume and pressure overload*¹
- *Severe AR + symptoms or LVEF <55% or ventricle enlargement*²
- *Asymptomatic moderate AR is associated with adverse outcomes*³
- *LV remodeling in moderate and severe AR is similar*⁴


How can we BETTER assess risk?

- *AR is not frequently in isolation ¹*
- *Early LV dysfunction increases risk ²*
- *Ongoing trials in moderate AS and moderate FMR*


Risk is a continuum

What has been done?

Cardiac damage in severe aortic stenosis ¹




Cardiac damage in moderate to severe aortic regurgitation ²



**Cardiac MRI*

Methodology



	Stage 0 (n=28)	Stage 1 (n=101)	Stage 2 (n=210)	Stage 3 (n=93)
	No cardiac damage	LV damage	LA or Mitral valve damage	Pulmonary valve, Tricuspid valve, or RV damage
		GLS > -19.5%	LAVI > 34 mL/m ²	≥ moderate TR
Echocardiographic staging		EF < 55%	Atrial fibrillation	Any RV systolic dysfunction
		ILVESD > 2	≥ moderate MR	RVSP > 60 mmHg

- *AR moderate-severe*
- *2008-2020*
- *Exclusion criteria*
- *n=432*
- *Hierarchical staging*
- *Criteria based on validated thresholds* ^{1,2,3}

Results – rate of each individual component

	Stage 0 (n=28)	Stage 1 (n=101)	Stage 2 (n=210)	Stage 3 (n=93)
GLS > -19.5%	0 (0%)	94 (93.1%)	164 (78.1%)	82 (88.2%)
iLVESD > 2	0 (0%)	15 (14.9%)	43 (20.5%)	26 (28.0%)
LVEF < 55%	0 (0%)	23 (22.8%)	42 (20%)	40 (43.0%)
LAVI > 34	0 (0%)	0 (0%)	135 (64.3%)	71 (76.3%)
Atrial fibrillation	0 (0%)	0 (0%)	155 (73.8%)	72 (77.4%)
≥ moderate MR	0 (0%)	0 (0%)	37 (17.6%)	29 (31.2%)
RVSP > 60 mmHg	0 (0%)	0 (0%)	0 (0%)	21 (22.6%)
≥ moderate TR	0 (0%)	0 (0%)	0 (0%)	75 (80.6%)
RV systolic dysfunction	0 (0%)	0 (0%)	0 (0%)	36 (38.7%)


Results – baseline characteristics

	Stage 0 (n=28)	Stage 1 (n=101)	Stage 2 (n=210)	Stage 3 (n=93)	P-value
Female	11 (39.3%)	50 (49.5%)	85 (40.5%)	46 (49.5%)	0.30
Age	56.5 (43-68)	63 (50.5-74.0)	68.0 (58.0-76.0)	74.0 (63.0-82.5)	<0.0001
NT pro BNP	76 ± 11	1,293 ± 3,398	4,852 ± 1,755	8,934 ± 1,682	0.14
Atrial fibrillation	0 (0%)	0 (0%)	55 (26.2%)	21 (22.6%)	<0.0001
Severe AR	11 (39.3%)	36 (35.6%)	81 (38.6%)	50 (53.7%)	0.048
HF	9 (32.1%)	49 (48.5%)	140 (66.7%)	82 (88.2%)	<0.0001
CKD	3 (10.7%)	23 (22.8%)	64 (30.5%)	38 (40.9%)	0.005

Results – echocardiography


	Stage 0 (n=28)	Stage 1 (n=101)	Stage 2 (n=210)	Stage 3 (n=93)	P-value
LVEF (%)	61.9 (60.0-65.0)	60.0 (55.0-62.5)	59.1 (55.0-62.7)	55.0 (45.0-60.0)	<0.0001
RVSP (mmHg)	26.5 (23.0-30.1)	28.0 (24.0-33.9)	30.3 (27.1-37.7)	45.5 (35.8-56.9)	<0.0001
LAVI (mL/m ²)	24.7 (20.4-30.8)	25.7 (21.1-29.4)	36.7 (29.0-46.9)	46.6 (35.4-58.5)	<0.0001
GLS endocardium (%)	-20.7 (-22.8- -20.1)	-16 (-18.5- -13.7)	-15.7 (-19- -12.8)	-12.2 (-16.7- -8.0)	<0.0001
LVEDD (cm)	4.7 (4.4-5.0)	4.7 (4.2-5.3)	5.0 (4.4-5.5)	4.7 (4.3-5.4)	0.10
LVESD (cm)	2.8 (2.6-3.0)	3.0 (2.6-3.5)	3.2 (2.7-3.7)	3.3 (2.8-4.0)	<0.0001
>=mod MR	0 (0%)	0 (0%)	37 (17.6%)	29 (31.2%)	<0.0001
>=mod TR	0 (0%)	0 (0%)	0 (0%)	75 (80.7%)	<0.0001
MV E/e' septal	9.5 (8.5-12.4)	11.9 (9.0-15.9)	14.6 (11.0-20.6)	16.9 (13.4-36.0)	<0.0001

Results – outcomes



Results – outcomes


HF Hospitalizations



Number at risk

Stage 0	28	28	28	27	24	24
Stage 1	101	94	84	77	72	61
Stage 2	210	180	167	151	135	116
Stage 3	93	54	44	40	32	27


All-cause mortality



Number at risk


Stage 0	28	28	28	27	24	24
Stage 1	101	94	88	84	81	68
Stage 2	210	191	183	168	155	136
Stage 3	93	70	63	58	49	41

Results – sensitivity analysis



Number at risk

Stage 0	28	28	28	27	24	23
Stage 1	67	64	57	53	48	39
Stage 2	34	29	28	24	24	23
Stage 3	210	180	167	151	135	116
Stage 4	57	37	32	29	26	22
Stage 5	36	18	13	12	7	6



Results – multivariate model

All-cause mortality and HF hospitalizations

	HR (95% CI)	P-value
Stage (per each level increase)	2.1 (1.6-2.6)	<0.01
Age (per each year increase)	1.05 (1.03-1.07)	<0.01
HF	2.0 (1.3-3.1)	<0.01
Prior CABG	1.8 (1.2-2.6)	<0.01
DM	1.9 (1.4-2.7)	<0.01

Cardiac damage staging has the strongest association with the primary outcome

Limitations

- *Single center and retrospective study*
- *Qualitative assessment*
- *GLS is variable among different institutions*
- *Age and comorbidities are highly associated with staging*

Conclusions

- *Novel AR-specific cardiac damage staging system was developed and tested*
- *Staging incorporates GLS, structural and functional echocardiographic markers*
- *Progressive cardiac damage was strongly associated with adverse clinical outcomes*

Future

Cardiac damage captures continuum risk

- More advanced stages benefit from early AVR?
- What's the degree of cardiac damage in mild AR?
- Is there structural recovery after AVR?

Acknowledgements



Ian Persits, DO , Joseph Hajj, MD, Ziad Zalaquett, MD, V Karthik, MD, Travis Howard, MD, Vincent Chen, MD, Samir R. Kapadia, MD, Serge C. Harb, MD

Rishi Puri, MD, PhD