

# Dimensionless Index as a Prognostic Marker of Mortality in Aortic Stenosis with Preserved Ejection Fraction

*A Meta-analysis and Systematic Review*

Hriday Shah, MD

Internal Medicine Resident

New York Medical College St Mary's General Hospital and St Clare's Health



**TCT**<sup>®</sup>

TRANSCATHETER  
CARDIOVASCULAR  
THERAPEUTICS<sup>®</sup>



# Disclosure

- I, Hriday Shah, DO NOT have any financial relationships to disclose.



# Acknowledgement

## Co-Investigators

- Haider Kashif, MD
- Kaushal Patel, MD
- Nirmal Patel, MD
- Ajitha Ganesan, MD
- Nithin Kumar, MD
- Amna Zainab, MD
- Ramsha Jeoffrey, MD
- Saraschandra Vallabhajosyula, MD



## Background

- Aortic stenosis (AS) is a common valvular heart disease in aging populations.
- Causes: degenerative calcification or congenital bicuspid valve.
- Severity assessment: Aortic Valve Area (AVA), jet velocity, gradient—limited in preserved Ejection Fraction (EF).

### Pathophysiology:

- Increased LV afterload → hypertrophy, fibrosis, decreased compliance.
- Traditional parameters are flow-dependent and may misclassify severity.



# Limitations of Current Assessment

- AVA estimation depends on Left ventricular outflow tract (LVOT) geometry and flow.
- Echocardiography can underestimate AVA due to turbulence or shape distortion.



# Understanding the Dimensionless Index (DI)

The Dimensionless Index (DI) is a ratio used in echocardiography to assess Aortic Stenosis (AS) severity.

Formula:  $DI = \text{LVOT Velocity Time Integral} \div \text{Aortic Jet Velocity Time Integral}$ .

- **LVOT VTI (Left Ventricular Outflow Tract Velocity Time Integral)**
- Measures blood flow speed and distance in the LVOT during systole.
- Reflects how far blood moves out of the left ventricle with each beat (“stroke distance”).
- Used to estimate stroke volume and cardiac output.
  
- **Aortic VTI (Aortic Jet Velocity Time Integral)**
- Measures velocity and distance of blood flow through the aortic valve.
- Reflects how fast blood passes the aortic valve — higher in stenosis.
- Used with LVOT VTI to calculate the Dimensionless Index (DI) for aortic stenosis severity.



# Objective

- To assess mortality in AS patients with  $DI \leq 0.25$  vs  $\geq 0.25$  and preserved EF.
- Primary endpoint: Overall mortality.



# Methods

- Systematic review & meta-analysis per PRISMA guidelines.
- Databases: PubMed, Cochrane, EMBASE, Scopus, Google Scholar (till July 2025).
- Random-effects model using RevMan 5.4.



# Selection Criteria

- Inclusion: AS with preserved EF, clear DI values.
- Exclusion: Mitral disease, prosthetic valves, AVA > 2.0 cm<sup>2</sup>, pediatrics, <10 patients.



# Results Summary


- 5 studies (3 observational, 2 abstracts) included.
- $DI \leq 0.25$  associated with higher mortality (HR 1.69 [95% CI 1.46–1.97],  $p < 0.00001$ ).
- Low heterogeneity ( $I^2 = 26\%$ ).



# Forest Plot – Mortality

- Visual representation of pooled Hazard Ratios.
- HR consistently  $>1.5$  for  $DI \leq 0.25$  group.
- Indicates strong association with mortality.

Fig 1: Forest plot for **overall mortality**





# Sensitivity & Bias Assessment

**Figure 2:** Quality Assessment using ROBINS-I V2 tool

		Risk of bias domains						
		D1	D2	D3	D4	D5	D6	D7
Study	Leibowitz et al. (2024)	-	-	-	+	-	+	-
	Altes et al. (2020)	-	-	-	+	-	+	-
	Rusinaru et al. (2015)	X	-	X	+	-	+	-
		Overall						
		-	-	-	+	-	+	-
		-	-	-	+	-	+	-
		X	-	X	+	-	+	-

Domains:  
D1: Bias due to confounding.  
D2: Bias due to selection of participants.  
D3: Bias in classification of interventions.  
D4: Bias due to deviations from intended interventions.  
D5: Bias due to missing data.  
D6: Bias in measurement of outcomes.  
D7: Bias in selection of the reported result.

Judgement  
X Serious  
- Moderate  
+ Low



# Discussion

- $DI < 0.25$  is a strong predictor of mortality in AS with preserved EF.
- DI complements AVA and gradient in assessing severity.
- Independent of flow and geometry, improving accuracy.



# Clinical Implications

- Incorporate DI with standard Doppler parameters for SAS.
- Improves sensitivity and prognostic accuracy.
- May guide earlier valve replacement decisions.




# Limitations

- Small number of included studies and limited demographics.
- Abstract-only data reduced robustness of quality assessment.



# Conclusion

- $DI \leq 0.25$  predicts higher mortality in AS with preserved EF.
- Useful for risk stratification and guiding valve intervention.
- Future research needed for DI in low- vs normal-flow AS.





# References

1. Leibowitz D, Zwas D, Elbaz-Greener G, Planer D, Alcalai R, Gotsman I. Dimensionless index and outcome in patients with aortic valve disease. Echocardiography [Internet]. 2024 Jun 1 [cited 2025 Jul 10];41(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/38814015/>
2. Mantha Y, Futami S, Moriyama S, Hieda M. Valvulo-Arterial Impedance and Dimensionless Index for Risk Stratifying Patients With Severe Aortic Stenosis [Internet]. Vol. 8, Frontiers in Cardiovascular Medicine. Frontiers Media SA; 2021 [cited 2025 Jul 10]. Available from: <https://pubmed.ncbi.nlm.nih.gov/34926605/>
3. Joseph J, Naqvi SY, Giri J, Goldberg S. Aortic Stenosis: Pathophysiology, Diagnosis, and Therapy [Internet]. Vol. 130, American Journal of Medicine. Elsevier Inc.; 2017 [cited 2025 Sep 26]. p. 253–63. Available from: <https://pubmed.ncbi.nlm.nih.gov/27810479/>
4. Leibowitz D, Zwas D, Elbaz-Greener G, Planer D, Alcalai R, Gotsman I. Dimensionless index and outcome in patients with aortic valve disease. Echocardiography [Internet]. 2024 Jun 1 [cited 2025 Jul 10];41(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/38814015/>
5. Rusinaru D, Malaquin D, Maréchaux S, Debry N, Tribouilloy C. Relation of Dimensionless Index to Long-Term Outcome in Aortic Stenosis with Preserved LVEF. JACC Cardiovasc Imaging [Internet]. 2015 Jul 1 [cited 2025 Sep 26];8(7):766–75. Available from: <https://pubmed.ncbi.nlm.nih.gov/26093931/>
6. Altes A, Thellier N, Rusinaru D, Marsou W, Bohbot Y, Chadha G, et al. Dimensionless Index in Patients With Low-Gradient Severe Aortic Stenosis and Preserved Ejection Fraction. Circ Cardiovasc Imaging [Internet]. 2020 Oct 1 [cited 2025 Jul 10];13(10):E010925. Available from: <https://pubmed.ncbi.nlm.nih.gov/33076698/>
7. Altes A, Thellier N, Rusinaru D, Marsou W, Bohbot Y, Chadha G, et al. Clinical significance of dimensionless index in patients with low-gradient severe aortic stenosis and preserved ejection fraction. Arch Cardiovasc Dis Suppl [Internet]. 2021 Jan 1 [cited 2025 Jul 10];13(1):48. Available from: <https://www.sciencedirect.com/science/article/pii/S1878648020302317#abst0025>
8. Minners J, Allgeier M, Gohlke-Baerwolf C, Kienzle RP, Neumann FJ, Jander N. Inconsistencies of echocardiographic criteria for the grading of aortic valve stenosis. Eur Heart J [Internet]. 2008 Apr 1 [cited 2025 Sep 26];29(8):1043–8. Available from: <https://dx.doi.org/10.1093/eurheartj/ehm543>



# References

9. Rusinaru D, Malaquin D, Maréchaux S, Debry N, Tribouilloy C. Relation of Dimensionless Index to Long-Term Outcome in Aortic Stenosis with Preserved LVEF. JACC Cardiovasc Imaging [Internet]. 2015 Jul 1 [cited 2025 Jul 10];8(7):766–75. Available from: <https://pubmed.ncbi.nlm.nih.gov/26093931/>
10. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews [Internet]. Vol. 372, The BMJ. BMJ; 2021 [cited 2024 Jun 7]. Available from: <https://pubmed.ncbi.nlm.nih.gov/33781993/>
11. Cochrane Bias Methods Group and the Cochrane Non-Randomised Studies Methods Group. Risk of bias tools: ROBINS-I V2 tool [Internet]. 2025 [cited 2025 Sep 26]. Available from: <https://www.riskofbias.info/welcome/robins-i-v2>
12. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. In: Research Synthesis Methods [Internet]. Res Synth Methods; 2021 [cited 2025 Sep 26]. p. 55–61. Available from: <https://pubmed.ncbi.nlm.nih.gov/32336025/>
13. Namasivayam M, Churchill TW, Capoulade R, Pibarot P, Danik JS, Picard MH, et al. Combined Value of Dimensionless Index and Transvalvular Flow Rate in Risk Stratification of Aortic Stenosis. Am J Cardiol [Internet]. 2024 Feb 15 [cited 2025 Sep 26];213:69–71. Available from: <https://pubmed.ncbi.nlm.nih.gov/38103765/>
14. Rusinaru D, Malaquin D, Maréchaux S, Debry N, Tribouilloy C. Relation of Dimensionless Index to Long-Term Outcome in Aortic Stenosis with Preserved LVEF. JACC Cardiovasc Imaging [Internet]. 2015 Jul 1 [cited 2025 Sep 26];8(7):766–75. Available from: [/doi/pdf/10.1016/j.jcmg.2015.01.023?download=true](https://doi/pdf/10.1016/j.jcmg.2015.01.023?download=true)
15. Nishimura RA, Otto CM, Bonow RO, Carabello BA, Erwin JP, Guyton RA, et al. 2014 AHA/ACC guideline for the management of patients with valvular heart disease: A report of the American college of cardiology/American heart association task force on practice guidelines. J Am Coll Cardiol [Internet]. 2014 Jun 10 [cited 2025 Sep 26];63(22). Available from: [/doi/pdf/10.1016/j.jacc.2014.02.536?download=true](https://doi/pdf/10.1016/j.jacc.2014.02.536?download=true)
16. Otto CM, Pearlman AS, Comess KA, Reamer RP, Janko CL, Huntsman LL. Determination of the stenotic aortic valve area in adults using Doppler echocardiography. J Am Coll Cardiol [Internet]. 1986 [cited 2025 Sep 26];7(3):509–17. Available from: <https://pubmed.ncbi.nlm.nih.gov/3950230/>
17. Jander N, Hochholzer W, Kaufmann BA, Bahlmann E, Gerdtts E, Boman K, et al. Velocity ratio predicts outcomes in patients with low gradient severe aortic stenosis and preserved EF. Heart [Internet]. 2014 Dec 1 [cited 2025 Sep 26];100(24):1946–53. Available from: <https://pubmed.ncbi.nlm.nih.gov/25217488/>