

# Outcomes of Transcatheter Aortic Valve Replacement in Patients with Cardiac Amyloidosis: Impact of Disease Recognition and Treatment Response

***Cleveland Clinic Foundation***

Tamari Lomaia, MD

---



**TCT**®

TRANSCATHETER  
CARDIOVASCULAR  
THERAPEUTICS®

# Authors

Tamari Lomaia MD , Besir Besir MD, Judah Rajendran MD , Neha Sangani ,  
Vishalraj Bandari MD, Osamah Badwan MD, Judah Rajendran MD , Shivabalan  
Kathavarayan Ramu MD, Rhonda Miyasaka MD, Serge Harb MD, James Yun  
MD PhD, Grant Reed MD MSc, Rishi Puri MD PhD, Amar Krishnaswamy MD

Special thanks to our PI and mentor Samir Kapadia MD FACC

# Disclosure of Relevant Financial Relationships

I, [Tamari Lomaia](#) DO NOT have any financial relationships to disclose.


# Background

-prevalence of calcific aortic stenosis and amyloid transthyretin cardiomyopathy (ATTR-CM) increase with age, and they often coexist. Older patients with severe aortic stenosis (AS) are increasingly identified as having cardiac amyloidosis (CA)

-Studies have shown that TAVR improved outcomes in AS amyloid. However, cardiac amyloidosis itself is associated with worse outcomes.

*-No published study to date specifically compares patients with organ-limited cardiac amyloidosis to patients without amyloidosis, both groups having received transcatheter aortic valve replacement (TAVR), and reports 5-year mortality outcomes*

# Study design



# Methods

-Using TriNetX database, we identified 546 patients with cardiac amyloidosis who underwent TAVR and matched them 1:1 to 546 patients without documented cardiac amyloidosis using propensity score matching.

-Baseline demographics, comorbidities, and procedural variables were compared.

-Survival probabilities were estimated using Kaplan–Meier methods and compared using the log-rank test.

# Baseline Characteristics

Variables	Before Propensity Matching			After Propensity Matching		
	Cohort 1 N=546	Cohort 2 n=66572	P-value	Cohort 1 N=546	Cohort 2 N=546	P-value
Current age	82±7	81±8	0.01	82±7	83±7	0.06
Female sex	178(32.6%)	28460(40.7%)	<0.0001	178(32.6%)	178(32.6%)	1
Male sex	351(64.3%)	38193(54.6%)		351(64.3%)	351(64.3%)	1
BMI	27.7±6	29.3±6	<0.0001	27.7±6	28.1±6	0.33
White race	421(77.1%)	57630(82.4%)	0.001	421(77.1%)	417(76.3%)	0.77
Black or African American race	55(10%)	3099(4.4%)	<0.0001	55(10%)	56(10.2%)	0.92
Asian	29(5.3%)	1848(2.6%)	0.0001	29(5.3%)	28(5.1%)	0.89
Hypertensive diseases	521(95.4%)	58058(83.1%)	<0.0001	521(95.4%)	526(96.3%)	0.44
Ischemic heart disease	493(90.3%)	54502(77.9%)	<0.0001	493(90.3%)	500(91.6%)	0.46
Diseases of arteries, arterioles and capillaries	383(70.14%)	35588(50.9%)	<0.0001	383(70.14%)	386(70.7%)	0.84

Pulmonary heart disease and diseases of pulmonary circulation	296(54.2%)	20901(29.9%)	<0.0001	296(54.2%)	282(51.6%)	0.39
Chronic rheumatic heart disease	285(52.2%)	224157(24.6%)	<0.0001	285(52.2%)	279(51.1%)	0.71
Cerebrovascular diseases	269(49.3%)	22560(32.3%)	<0.0001	269(49.3%)	260(47.6%)	0.58
Unspecified disorders of circulatory system	222(40.6%)	14576(20.8%)	<0.0001	222(40.6%)	227(41.5%)	0.75
Diseases of veins, lymphatic vessels and lymph nodes	167(30.5%)	11881(16.9%)	<0.0001	167(30.5%)	149(27.3%)	0.22
Acute rheumatic fever	10(1.8%)	240(0.34%)	<0.0001	10(1.8%)	10(1.83%)	1
Other forms of heart disease	544(99.6%)	68414(97.8%)	0.004	544(99.6%)	546(100%)	0.15

Variables	Before propensity matching			After propensity matching		
	Cohort 1	Cohort 2	P-value	Cohort 1	Cohort 2	P-value
Heart rate	73±14	72±14	0.17	73±14	72±14	0.29
Systolic Blood pressure	127±21	128±22	0.26	127±21	127±21	0.65
LVEF	55.9±12	57±13	0.27	55.9±12	54.1±13.8	0.23

# 5-year survival


## Cohort Statistics

Variable	Patients in cohort	Patients with outcome	Risk of outcome
Cohort 1 (TAVR+Amyloid)	546	164	30%
Cohort 2 (TAVR without amyloid)	546	144	26.3%


## Log rank test

Variable	Patients in cohort	Survival probability at the end of time window	P-value
Cohort 1 (TAVR+Amyloid)	546	31.54%	0.0021
Cohort 2 (TAVR without amyloid)	546	44.58%	

Risk of Outcome



Kaplan Meier Survival Curve





# Conclusion

- TAVR offers a safe and effective treatment option for patients with or without cardiac amyloidosis. However, long-term survival may be adversely affected by disease progression and suboptimal response to therapies such as Tafamidis.
- These findings highlight the importance of routine screening for cardiac amyloidosis in TAVR candidates
- early detection and targeted treatment may significantly improve patient outcomes.

# References

- Nitsche, C., Dobner, S., Rosenblum, H. R., et al. (2025). Cardiac transthyretin amyloidosis treatment improves outcomes after aortic valve replacement for severe stenosis. *European Heart Journal*. Advance online publication. <https://doi.org/10.1093/eurheartj/ehaf362>
- Riley, J. M., Junarta, J., Ullah, W., et al. (2023). Transcatheter aortic valve implantation in cardiac amyloidosis and aortic stenosis. *The American Journal of Cardiology*, 198, 101–107. <https://doi.org/10.1016/j.amjcard.2023.04.003>
- Nitsche, C., Scully, P. R., Patel, K. P., et al. (2021). Prevalence and outcomes of concomitant aortic stenosis and cardiac amyloidosis. *Journal of the American College of Cardiology*, 77(2), 128–139. <https://doi.org/10.1016/j.jacc.2020.11.006>
- Scully, P. R., Patel, K. P., Treibel, T. A., et al. (2020). Prevalence and outcome of dual aortic stenosis and cardiac amyloid pathology in patients referred for transcatheter aortic valve implantation. *European Heart Journal*, 41(29), 2759–2767. <https://doi.org/10.1093/eurheartj/ehaa170>
- Fatima, K., Uddin, Q. S., Tharwani, Z. H., et al. (2024). Concomitant transthyretin cardiac amyloidosis in patients undergoing TAVR for aortic stenosis: A systemic review and meta-analysis. *International Journal of Cardiology*, 402, 131854. <https://doi.org/10.1016/j.ijcard.2024.131854>

**Thank you**