

Moderate Trans-Valvular Aortic Insufficiency Immediately After Implantation With Self Resolution One Month Post Procedure

Self resolving Intra-prosthetic Aortic Regurgitation

Marlena Lesniowska, MD, PGY3, Internal Medicine

Fadi Ghrair, MD, Interventional Cardiology/Structural PD



TCT[®]

TRANSCATHETER
CARDIOVASCULAR
THERAPEUTICS[®]

Bonsecours Mercy Health
St. Elizabeth Youngstown, Ohio

Disclosure of Relevant Financial Relationships


I, Marlena Lesniowska *DO NOT* have any financial relationships to disclose.

Background

- Despite the evolving technology of transcatheter valves, regurgitation post-TAVR is common, with a wide range of reported incidences
- In patients treated with transcatheter valve-in-valve implantation for failed surgical bio-prosthesis, the incidence of significant AR seems to be comparable to TAVI in native valves
- Patients with valvular regurgitation post TAVR experience higher risk of all-cause mortality, rehospitalization, and cardiovascular mortality.

Regurgitation mechanism



- Transvalvular
- Paravalvular
- Supra-skirtal



Case Summary



- 73-year-old female with prior SAVR with #19 mm Edwards Intuity valve in 2017
- Presented with severe symptomatic bioprosthetic aortic stenosis and moderate intra-valvular regurgitation
 - *AVA was 0.6 cm², Vmax of 4.8 m/s, and a MG of 56 mmHg.*
 - EF of 60 – 65%
- Underwent Transcatheter Valve-in-Valve implantation procedure
 - Right common femoral artery access, deploying a 20-mm SAPIEN 3 Ultra Resilia valve under rapid pacing.
 - Post dilatation was performed using 20 mm True balloon for valve frame optimization

Post implantation Echo imaging



- Moderate regurgitation
- Likely transvalvular rather than peri-valvular based on transthoracic imaging
- Mean gradient of 9 mmHg on TTE

TEE and fluoroscopy post deployment



What happened?

- Confirmation of leak etiology/location/type (paravalvular vs transvalvular vs supra-skirtal)
- Transvalvular leak pathology:
 - Damaged leaflet
 - Stuck leaflet



What to do next?

- Further balloon dilatation (risk of further damage to leaflets)
- Valve in valve in valve (risk of patient prosthesis mismatch and elevated gradients, especially with an already small 20 mm valve)
- Redo surgical procedure (High risk patient with high-risk procedure)
- Leaflet manipulation (via pigtail)
- Watchful waiting (since patient was hemodynamically stable)

Patient Update

- Patient was hemodynamically stable and compensated (likely due to a preprocedural moderate AR)
- Decision was made to clinically observe patient and assess her symptoms as an outpatient and if she becomes symptomatic or if regurgitation worsens/persists, we always have the option of bringing her back for Valve in valve in valve procedure.
- TTE POD#1 with MG 19 mmHg, with a persistent moderate AR.

TTE 1 month post procedure



Conclusion

- *Our case* is the first that shows Transvalvular AI (intra-prosthetic) with spontaneous resolution on the 1-month follow up imaging
- Proposed mechanism is a stuck leaflet with spontaneous restoration of normal movement.
- Possibly related to the new Resilia dry-storing technology
- Watchful waiting with periodic clinical and echocardiographic reassessment may be appropriate
- Recommended for hemodynamically stable patients
- Aims to avoid unnecessary repeat valve-in-valve procedures
- Prevents added procedural risk and hemodynamic compromise and helps maintain effective orifice area and reduce risk of prosthesis–patient mismatch.

Sources

1. Stähli BE, Maier W, Corti R, et al. Aortic regurgitation after transcatheter aortic valve implantation: mechanisms and implications. *Cardiovascular Diagnosis and Therapy*. 2013;3(1):15-22
2. Vahanian A, Beyersdorf F, Praz F, et al. 2021 ESC/EACTS guidelines for the management of valvular heart disease: developed by the task force for the management of valvular heart disease of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2022;43:561–632.34453165
3. Werner N, Sinning JM. Aortic Regurgitation After Transcatheter Aortic Valve Replacement — Nothing to Worry About Anymore? *Circulation Journal*. 2014; 78: 811–818
4. Villari B, Hess OM, Kaufmann P, et al. *Effect of aortic valve stenosis (pressure overload) and regurgitation (volume overload) on left ventricular systolic and diastolic function*. *American Journal of Cardiology*. 1992; 69(9):927-934
5. Sá MPBO, Sampaio RO, Cavalcanti LRP, et al. Impact of paravalvular leak on outcomes after transcatheter aortic valve implantation: meta-analysis of Kaplan–Meier–derived individual patient data. *Frontiers in Cardiovascular Medicine*. 2023;10:1165665. doi:10.3389/fcvm.2023.1165665