

Cerebral Embolic Protection Devices in Transcatheter Aortic Valve Replacement: A Meta-Analysis of Randomized Controlled Trials

Transcatheter Cardiovascular Therapeutics (TCT) 2025

Mostafa Ali, MD – Crestwood Medical Center, Huntsville, Alabama



TCT[®]

TRANSCATHETER
CARDIOVASCULAR
THERAPEUTICS[®]

Disclosure of Relevant Financial Relationships

- I, Mostafa Ali, DO NOT have any financial relationships to disclose.
- Faculty disclosure information can be found on the TCT App.

Background

- Stroke remains a serious complication of TAVR (2–6%).
- Cerebral Embolic Protection Devices (CEPDs) capture or deflect debris.
- Previous trials show mixed results.
- This meta-analysis evaluates clinical efficacy and safety outcomes.

Objective

- To evaluate the impact of CEPDs on:
 - Stroke (overall, disabling, and non-disabling)
 - Mortality, bleeding, and vascular complications
- Using data from randomized controlled trials only.


Methods

- Systematic search: PubMed, Scopus, Cochrane, Web of Science (June 2025).
- Inclusion: RCTs comparing CEPD vs Control during TAVR.
- Random-effects model; Risk Ratio (RR) [95% CI].
- Primary outcome: overall stroke at 2–5 and 30 days.


Included Studies

- 8 RCTs (n = 11,623):
- CLEAN-TAVI, MISTRAL-C, DEFLECT III, REFLECT I/II,
- SENTINEL, PROTECTED TAVR, BHF PROTECT-TAVR.
- Devices: Sentinel, TriGuard, EMBOL-X.


Results – Early Disabling Stroke



Results – Other Outcomes



Results – Other Outcomes



Discussion

- Findings align with Balata et al 2025 and Warraich et al 2025 meta-analyses.
- CEPDs reduce early disabling stroke but not overall stroke or mortality.
- Small absolute benefit may still be clinically meaningful.
- Routine use not justified; selective use may be appropriate.

Limitations

- Low event rates.
- Device heterogeneity (Sentinel vs TriGuard vs EMBOL-X).
- Limited follow-up and MRI standardization.

Conclusion

- CEPDs reduce early disabling stroke after TAVR.
- No significant difference in overall stroke or mortality.
- Supports selective, not routine, use in high-risk patients.
- Future: cost-effectiveness and long-term outcomes.

Clinical Implications

- Consider in prior stroke, heavy calcification, or high-risk anatomy.
- Tailored approach improves neuroprotection.
- Encourages ongoing innovation in device design.

Acknowledgments

- No conflicts of interest.
- Gratitude to co-authors and TCT 2025 organizers.
- Contact: ali.mostafa25md@gmail.com

Co-Authors & Collaborators

- **First Author:** Ahmed Emara, MD
- **Presenting Author:** Mostafa Ali, MD
- **Co-Authors:**
 - Ahmed Abdel Azim, MD
 - Ahmed Diaa, MD
 - Mohamed Emara, MD
 - Abdalkhakim Shubietah, MD
 - Mohamed Ellebedy, MD
 - Mohamed Elgendy, MD
 - Ahmed Mohamed Elsayed Rezk, MD
 - Michael Megaly, MD

References

- Haussig S et al. JAMA 2016;316:592–601.
- Lansky AJ et al. Eur Heart J 2015;36:2070–2078.
- Mieghem NM et al. EuroIntervention 2016;12:1290–1297.
- Kapadia SR et al. JACC 2017;69:367–377.
- Nazif TM et al. JACC Cardiovasc Interv 2021;14:1234–1245.
- Lansky AJ et al. Eur Heart J 2021;42:2562–2573.
- Kapadia SR et al. N Engl J Med 2022;387:1253–1263.
- Kharbanda RK et al. N Engl J Med 2025;392:112–121.
- Warraich HJ et al. BMC Cardiovasc Disord 2025;25:1128.