

Can We Improve TAVR Durability Today?

Imaging-based Procedural Solutions

João Cavalcante MD, FASE, FSCCT, FSCMR

Section Head, Cardiac Imaging

Allina Health Minneapolis Heart Institute

Scientific Director, Cardiovascular Imaging Core Lab and Research Center

Chair, Van Tassel Innovation Center



TCT[®]

TRANSCATHETER
CARDIOVASCULAR
THERAPEUTICS[®]

Disclosures

☒ **I have the following potential conflicts of interest to declare:**

Receipt of grants / research support: Abbott Structural, Allina Health Foundation, JenaValve, NIH/NHLBI, Scripps Research Institute.

Receipt of honoraria or consultation fees: 4C Medical, Abbott Structural, Alleviant, Anteris, Braile Medical, Boston Scientific, Edwards Lifesciences, JenaValve, JC Medical, Medtronic, Novo Nordisk, Nyra Medical, PIE Medical, VDYne, Zoll.

MHI Foundation Imaging Core Lab:

- Institutional Contracts: Abbott, Boston Scientific, Edwards Lifesciences, Medtronic, JC Medical, JenaValve, Products & Features.
- Research Support: PIE Medical
- Research Grants: Siemens Healthineers

Durability is a Multifactorial and Inter-Related Issue


- The Host AV Leaflet, Anatomy and Surrounding Tissue
- The Host Comorbidities (Age, CKD, Ca^{++} x Phosp, etc)
- Prosthetic valve type, valve size/PPM, Frame and Leaflet Design and Leaflet Tissue
- THV device preparation/crimping
- THV Frame Deformation / Frame Expansion
- How it is defined, *and some more we don't know...*



VARC-3 Definitions of BVD and BVF

Is the BVD related to intrinsic permanent changes to the prosthetic valve?

NO




Non-Structural BVD

Any abnormality, not intrinsic to the valve, resulting in BVD

- Prosthesis-patient mismatch
- Paravalvular regurgitation
- Other: mal-positioning, embolization, etc.


YES



Structural BVD

Intrinsic permanent structural changes to the prosthetic valve
Stage 1 SVD

YES/NO



Thrombosis
Endocarditis

Possible

May be reversible

No Hemodynamic Valve Deterioration during FU

Hemodynamic Valve Deterioration during FU
Stage 2 (Moderate); Stage 3 (Severe) SVD

Non-Structural BVF

Structural BVF


Bioprosthetic Valve Failure (BVF)

- i) Any BVD with clinically expressive criteria OR irreversible Stage 3 BVD
- ii) Re-intervention or indication for reintervention; iii) Valve-related death


Genereaux P et al. JACC 2021 Jun 1;77(21):2717-2746.

- Deformation at functional portion is crucial

- Deformation is 3D phenomenon





- Deformation could cause THV deterioration via fibrosis and calcification process




THV Shape Matters

HALT seen in 19% of patients at 30 days CT screening registry
No differences in THV gradients for HALT + vs. HALT -






HALT -	HALT +
1.04	1.21
THV deformation index = (a + c) / (2 × b)	
7.2	6.3
THV sinus volume (ml)	




565 patients at 30-days post TAVR

- 352 S3 (62%);
- 213 Evolut R/Pro+ (38%)



Blanke et al. JACC 2020 May 19;75(19):2430-2442.



Makkar, R.R. et al. J Am Coll Cardiol. 2020;75(24):3003-15.

How Often does HALT Occur and what is its Natural History (∅ AC)?

True finding? Likely to “resolve”

Harder to miss, unlikely to resolve spontaneously


SAPIEN 3

EVOLUT

SHORT AXIS


LONG AXIS

VOLUME RENDER



A= Diastolic Image B= Systolic Image



Treatment with warfarin was associated with resolution of HALT in 82% of patients undergoing serial imaging.



Anticoagulating everyone without HALT increases bleeding risk, and provides no gain...

Does HALT Matter?

CENTRAL ILLUSTRATION: A Brief Study Flowchart and Main Findings





No. at risk:						
— No HALT	689	617	552	461	310	189
— HALT	115	101	91	74	42	15
- - - No HALT	689	449	435	378	242	154
- - - HALT	115	76	70	60	29	13

Hein M, et al. J Am Coll Cardiol Interv. 2022;15(11):1113-1122.


3-year event rate of symptomatic HVD was 9.4% vs 1.5% (with vs without HALT, $P < 0.001$)
(HR: 6.10; 95% CI: 2.59-14.29; $P < 0.001$)

Atlantis-4D



Ischemic events are defined as the composite of death, myocardial infarction, stroke or peripheral embolism,

Montalescot et al. JACC Interv 2022 Sep 26;15(18):1794-1804.



“Thrombus was found on all valves, unlike the stated prevalence of 7% to 14% on CT”

THV degeneration appears to occur in a stepwise fashion: thrombus → fibrosis (after 60 days) → calcification after 4 years. Trend toward increased leaflet thickness throughout implantation.

Is HALT a Harmless Bystander?

Table 4. One-Year Clinical Outcomes

Variable	All patients (n=565)	HALT (n=108)	No HALT (n=457)	Unadjusted HR (95% CI)	P value	Adjusted HR (95% CI)	P value
All-cause death	40 (7%)	16 (15%)	24 (5%)	2.90 (1.54–5.46)	0.001	2.98 (1.57–5.63)*	0.001
Cardiac death	18 (3%)	9 (8%)	9 (2%)	4.29 (1.70–10.8)	0.002	4.58 (1.81–11.6)†	0.001
HF hospitalization	35 (6%)	10 (9%)	25 (6%)	1.77 (0.85–3.69)	0.13	1.91 (0.91–4.02)*	0.09
Composite (all cause death + HF hospitalization)	66 (12%)	21 (19%)	45 (10%)	2.08 (1.24–3.49)	0.006	1.94 (1.14–3.30)‡	0.02
Myocardial infarction	9 (2%)	6 (6%)	3 (1%)	4.10 (1.02–16.4)	<0.05		
Stroke/TIA	21 (4%)	8 (7%)	13 (3%)	1.29 (0.54–3.13)	0.57	1.27 (0.50–3.23)†	0.61
Bleeding event	56 (10%)	11 (10%)	45 (10%)	1.07 (0.55–2.07)	0.84	1.03 (0.53–2.00)*	0.92

HALT indicates hypoattenuating leaflet thickening; HF, heart failure; HR, hazard ratio; LV, left ventricular; STS-PROM, Society of Thoracic Surgeons Predicted Risk of Mortality; and TIA, transit ischemic attack.


*Adjusted by age, sex, and log STS-PROM score.

†Adjusted by log STS-PROM score.


‡Adjusted by age, sex, log STS-PROM score, LV ejection fraction at baseline, LV stroke volume index at 30 days after transcatheter aortic valve replacement; and not performed for myocardial infarction because of the small number of the event.

Perhaps not immediately harmful to the patient – but meaningful to the THV durability


What Factors Are Associated with Early Bioprosthetic Valve Deterioration,
And What Are The Clinical Implications?




306 Patients underwent TAVR
Early BVD = 44 (14.3%)
Without Early BVD = 262 (85.7%)



HALT
30% vs 5%







Valve Size
20-23 mm
57% vs 23%



BMI >30
45% vs 29%

Higher Incidence With Early BVD:

-  Heart Failure Hospitalization
-  Reintervention (Redo-TAVR)
-  Early Mortality




Anticoagulation = 4.3% with early BVD
No Anticoagulation = 20.6% with early BVD

Hypoattenuated Leaflet Thickening (HALT), small valve size, increased BMI were associated with higher rates of early BVD. Anticoagulation had a protective effect. Early BVD was associated with higher rates of reintervention, heart failure hospitalization, and early mortality

*BMI: Body Mass Index, HALT: Hypoattenuated Leaflet Thickening, TAVR: Transcatheter Aortic Valve Replacement

Central Illustration. Risk factors and clinical outcomes associated with early bioprosthetic valve deterioration.



Underexpanded Acurate neo2 Valves Tied to Worse Outcomes: ACURATE IDE

(UPDATED) The explanation for why the valve failed is too little, too late: Boston Scientific ended global sales of the device Wednesday.

by Michael O'Riordan | MAY 28, 2025



CENTRAL ILLUSTRATION Redo-TAVR Frame Expansion Analysis by MDCT and Fluoroscopy

A First-in-Human Study of MDCT and Fluoroscopy to Evaluate Valve Frame Expansion and Geometry of Index and Second TAV After Redo-TAVR

Study Flow

TAV-in-TAV Combinations

MDCT Findings


40 Redo-TAVR
(1/2023-4/2025)
30 Pre+Post Redo-TAVR MDCT
25 Fluoroscopic studies

= 3


% Underexpansion in Functional Zone



- Index TAV 100% underexpanded at baseline
- Expanded significantly after redo-TAVR




Symmetric TAVR Expansion



Asymmetric TAVR Expansion

High TAVR asymmetry index was associated with impaired prosthetic valve performance (mean residual gradient ≥ 20 mm Hg and/or \geq moderate paravalvular leak) on echocardiography before hospital discharge





Maznyczka et al. JACC Interv 2024 Sep 9;17(17):2011-2022

- Asymmetric prosthetic heart valve expansion occurred in 17% of the population
- TAVR asymmetry index was associated with impaired hemodynamic valve performance
- TAVR asymmetry index was not associated with clinical outcomes



TAVR in Native AS

1. Oversizing
2. Calcium distribution
3. Valve morphology (Bicuspid)
4. Underfilling (BEV)



Valve-in-Valve (Aortic, Mitral)

1. Oversizing
2. Implant depth



Final Thoughts

- Transcatheter valves durability will require long-term studies with standardized VARC-3 definitions
- HALT relates to stent frame deformation (underexpansion → pinwheeling → leaflet stress → thrombus, fibrosis, thickening)
- It is better to prevent HALT than to treat it. If so, how can we predict or avoid?
 - Should TAVR involve again pre/post-dilation to improve THV frame deformation?
 - Better sizing algorithms (intermediate sizing – MyVal, Braile, X4?)
 - Potential role of AV calcium leaflet modification techniques
 - Screening for HALT in those with frame asymmetry, high Ca^{++} and aggressive oversizing
- Stakes should be higher in asymptomatic severe AS → procedural optimization, proactive HALT screening and THV durability needs to be investigated.
- Such strategies, along with new THV stent and leaflet designs may help to achieve better THV frame expansion, improve achieving laminar flow, minimize leaflet stress and ultimately improve durability.

MHI Structural/ Advanced Imaging Group

- Victor Cheng
- Nadira Hamid
- Felipe Kazmirczak
- Thomas Knickelbine
- John Lesser
- Michael Miedema
- Marc Newell
- Erik Schelbert
- Jonathan Urbach

HealthCare Delivery & Innovation

- Steve Bradley
- Craig Strauss

MHIF

- Kris Fortman
- Ross Garberich
- Scott Sharkey
- Lisa Tindell
- Tamara O'Black
- Charles Zaugg

MHIF Imaging and Valve Research Center

- Maurice Enriquez-Sarano
- Miho Fukui
- Davide Margonato
- Oksana Marchenko
- Mohamed Abed
- Takahiro Nishihara (in memoriam)

MHI Leadership

- Bill Katsiyannis
- Raed Abdelhadi
- Sean Ging
- Marc Newell

MHIF Imaging Core Lab

- Miho Fukui
- Kristin Lambrecht
- Oksana Marchenko
- Mady Olson
- Michelle Montone Campbell
- Ruken Bastimar

Interventional/Structural Team

- Manos Brilakis
- Nicholas Burke
- Ivan Chavez
- Mario Gössl
- Michael Mooney
- Anil Poulouse
- Yader Sandoval
- Jay Traverse
- Kostis Voudris
- Yale Wang

CT Surgery

- Rizwan Attia
- **Vinnie Bapat**
- Erik Beckmann
- Carly Lodewyks
- Sarah Palmer
- Ben Sun
- Sebastian Iturra

Joao.Cavalcante@allina.com



@JoaoLCavalcante

Thank you!