

MCS-Supported Left Main Bifurcation PCI and Transfemoral TAVR in an Octagenarian Patient

Dhruvil A. Patel, MD

*Internal Medicine – Pediatrics, PGY-3
University of Chicago Medicine, Chicago, IL*

*Dhruvil Ashishkumar Patel MD, Akshat Rana BS, Stanton Greenstone BS,
Lauren Roark RN, Janet Friant APN, Jennifer Smazil APN, Moira McDowell
APN, Atman P. Shah MD, Sandeep Nathan MD, MSc., Jay Khambhati MD*



TCT®

TRANSCATHETER
CARDIOVASCULAR
THERAPEUTICS®



AT THE FOREFRONT

UChicago
Medicine

Disclosure of Relevant Financial Relationships

I, **Dhruvil Patel**, DO NOT have any financial relationships to disclose.


Background

- **Elderly patients with concomitant left main disease and severe aortic stenosis represent a high-risk patient population for MACCE/CV death**
- **There is significant variability of treatment patterns for this unique patient cohort, with variables affecting discussions including:**
 - Frailty
 - Comorbid burden
 - Procedure time
 - Heart team expertise & resources available


Case Presentation

- 90M with:
 - HFpEF
 - Prostate Cancer s/p XRT
 - Bladder Cancer, s/p TURBT
 - Type 2 Diabetes Mellitus c/b Peripheral Neuropathy

**Presents with CCS class
2 angina and NYHA class
2 DOE symptoms**



Case Presentation – Initial Workup





Case Presentation – Severe D3 Paradoxical LFLG Aortic Stenosis

LVEF 59%



- MG_{AV} 20mmHg
- PG_{AV} 38mmHg
- $V_{max(AV)}$ 3.1 m/s
- AVA 0.78cm²
- SV_i 30mL/m²

**Mild Aortic
Insufficiency
No other significant
valve disease**






Case Presentation – Severe D3 Paradoxical LFLG Aortic Stenosis


- AV Ca^{2+} Score: 2438




Case Presentation – Severe Paradoxical LFLG Aortic Stenosis



Femoral Access Site Appears Safe to Access





Planning An Approach?







Impella Supported LM PCI

- Impella-CP placement
- dLM-pLAD rotational atherectomy
- oLCX IVL with Shockwave C2+ (120 pulses delivered)



Per-PCI IVUS

- Prior to intervention, IVUS demonstrates proximal calcifications




Impella Supported LM PCI

**DK Crush 3.0 mm DES in the
Left Circumflex**



**Crush, Rewire, KBI, and LM-LAD stent
with 3.0 x 30 mm DES, POT with 4.5mm
balloon, Rewire, and KBI, re-POT**




Impella Supported LM PCI

**PCI of the left main was then attempted
with TIMI stage 3 Flow Observed**



8 Weeks Later – Transfemoral Transcatheter Aortic Valve Replacement via R CFA



Post-Valve Deployment & Recovery

- Post-valve deployment, the patient developed HGAVB. TVP left in place
- **Day #1 Post-Procedure:**
Dual-chamber PPM
- **Day #3 Post-Procedure:**
Discharged home
- **No readmissions**

30-Day Echo:

- MG_{AV} 8 mmHg
- PG_{AV} 16.6 mmHg
- $V_{max(AV)}$ 2.3 m/sec

**23 mm S3 Sapien Valve
is Well Seated with no
Peri-Valvular Leak**

Considerations

At the 30-day follow-up post-TAVR, no peri-valvular leak with resolution of the patient's chest pain on exertion was achieved.

- In older, higher-risk patients, the use of a staged Impella-supported bifurcation PCI (HR-PCI) as a bridge to TAVR is proven to be safe
- Multidisciplinary surgical and interventional decision-making was vital in establishing a curated treatment path that led to success.

Thank You For Your Time & Attention