

Name: _____

ECE 269: Linear Algebra and Applications, Midterm
Instructor: Behrouz Touri

- Policy:
 - *No electronics*
 - Closed book
 - All paper & scratch paper provided:
 - * feel free to use back of each page as a scratch paper,
 - * there is an extra page provided as a blank page at the back of the exam sheet,
 - * if you need extra papers, ask the proctors,
 - * write your name on top of each provided paper,
 - * all the papers need to be returned to the proctors (including cheat-sheets).
 - *You must show all work to receive credit for a solution...even if final answer is correct.*
 - The exam carries 6 extra points (out of 100).
- There are 3 equally weighted questions.
- *Please do not forget to write down your name at the top of each page.*
- In all of the following questions, $(\mathbb{F}, +, \cdot)$ is a field.

Name: _____

1. For each of the following statements, determine whether it is true or false. Explain your answer. Correct answers without explanation carry no point.

- a. There exists a 2×3 matrix $A \in \mathbb{R}^{2 \times 3}$, such that there are two matrices B and C with $BA = I_{3 \times 3}$ and $AC = I_{2 \times 2}$ (here, $I_{n \times n}$ is the $n \times n$ identity matrix).

- b. If columns of the matrix $A \in \mathbb{F}^{n \times n}$ are independent, then the columns of A^2 are also independent.

Name: _____

- c. For an inner-product space V with the inner product $\langle \cdot, \cdot \rangle$, if S is a basis and $\langle x, v \rangle = 0$ for all $v \in S$, then $x = 0$.

- d. For all matrices $A \in \mathbb{F}^{n \times n}$, $\text{rank}(A) = \text{rank}(A^T A)$.

Name: _____

2. Prove the following statements.

- (a) We say that a matrix $A \in \mathbb{F}^{n \times n}$ is a lower-triangular matrix if $A_{ij} = 0$ for $j > i$. Show that if A is an invertible lower-triangular matrix, then all its diagonal elements are non-zero.
- (b) Suppose that \mathbb{F} is a finite-field. Show that if A is invertible, then $A^k = I$ for some $k \geq 1$.
- (c) Is the statement in Part (b) still true for an infinite field? Justify your answer.

Name: _____

3. Consider the space of finite-energy real-valued functions $L_2([0, 1])$, i.e., the space of functions $f : [0, 1] \rightarrow \mathbb{R}$ with $\int_0^1 f^2(x)dx < \infty$.
- (a) Show that the mapping $\langle f(x), g(x) \rangle := \int_0^1 f(x)g(x)dx$ is an inner-product in this space.
- (b) Consider the three polynomials $p_1(x) = 1$, $p_2(x) = x$, and $p_3(x) = x^2$. Using the Gram-Schmidt procedure, find out three orthonormal polynomials $q_1(x), q_2(x), q_3(x)$, such that $\text{span}(\{p_1(x), \dots, p_i(x)\}) = \text{span}(\{q_1(x), \dots, q_i(x)\})$ for $i = 1, 2, 3$.

Name:

(blank page)