

Midterm 2 Answers
16 April, 2015

P1. (5 points) Ali arrives at the Bilkent bus stop to take a bus to Tunus. The chances that he finds a bus waiting upon arrival is 50%. If there is no bus, his waiting time is uniformly distributed between 0 and 10 mins. Let X denote Ali's waiting time (in minutes) from the time of arrival at the bus stop until a bus arrival (if there is a bus waiting $X = 0$).

(a) (3 pt) Determine and sketch the CDF of X .

$$F_X(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{2}, & x = 0, \\ \frac{1}{2} + \frac{1}{20}x, & 0 < x < 10, \\ 1, & x \geq 10 \end{cases}$$

(b) (2 pt) Determine the expectation of X .

$$E[X] = 2.5 \text{ min.}$$

P2. (8 points) Let X be a random variable with PDF

$$f_X(x) = \begin{cases} x^{-1}, & 1 \leq x \leq e, \\ 0, & \text{otherwise} \end{cases}$$

where $e \approx 2.718$ is the base of the natural logarithm.

(a) (2 pt) Let $A = \{X \geq \sqrt{e}\}$. Compute $P(A)$.

$$P(A) = \int_{\sqrt{e}}^e x^{-1} dx = \ln(x) \Big|_{\sqrt{e}}^e = \ln(e) - \ln(\sqrt{e}) = 1 - \frac{1}{2} = \frac{1}{2}.$$

$$P(A) = \frac{1}{2}$$

(b) (2 pt) Compute $f_{X|A}(x)$ as a function of x .

$$f_{X|A}(x) = \begin{cases} 2x^{-1}, & \sqrt{e} \leq x \leq e, \\ 0, & \text{otherwise} \end{cases}$$

(c) (2 pt) Compute $E[X|A]$. (Leave the result as a function of e .)

$$E[X|A] = \int_{-\infty}^{\infty} x f_{X|A}(x) dx = \int_{\sqrt{e}}^e x (2x^{-1}) dx = 2(e - \sqrt{e}).$$

$$E[X|A] = 2(e - \sqrt{e})$$

(d) (2 pt) Let $Y = X^2$. Calculate $E[Y]$. (Leave the result as a function of e .)

$$E[Y] = \int_{-\infty}^{\infty} x^2 f_X(x) dx = \int_1^e x dx = \frac{1}{2}(e^2 - 1).$$

$$E[Y] = \frac{1}{2}(e^2 - 1)$$

P3. (10 points)

Let (X, Y) be jointly distributed random variables with a joint PDF

$$f_{X,Y}(x, y) = \begin{cases} xy, & 0 < x < 1, 0 < y < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

(a) (4 pt) Find the marginal PDF $f_X(x)$ and $f_Y(y)$.

$$f_X(x) = \begin{cases} 2x, & 0 < x < 1, \\ 0, & \text{elsewhere.} \end{cases}$$

$$f_Y(y) = \begin{cases} \frac{1}{2}y, & 0 < y < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

(b) (4 pt) Let $Z = X + Y$. Compute the CDF and PDF of Z .

$$F_Z(z) = \begin{cases} 0, & x \leq 0, \\ \frac{z^4}{24}, & 0 < z \leq 1, \\ \frac{z^2}{4} - \frac{z}{3} + \frac{1}{8}, & 1 < z \leq 2 \\ -\frac{z^4}{24} + \frac{5z^2}{4} - 3z + \frac{17}{8}, & 2 < z \leq 3 \\ 1, & z > 3. \end{cases}$$

$$f_Z(z) = \begin{cases} \frac{z^3}{6}, & 0 < z \leq 1, \\ \frac{z}{2} - \frac{1}{3}, & 1 < z \leq 2 \\ -\frac{z^3}{6} + \frac{5z}{2} - 3, & 2 < z \leq 3 \\ 0, & \text{otherwise.} \end{cases}$$

(c) (2 pt) Compute $E[2X + 3Y + XY]$. The result must be a numerical value. Do not leave it as an integral.

$$E(2X + 3Y + XY) = \frac{56}{9}$$

P4. (8 points)

Let X, Y, Z be independent identically-distributed (i.i.d.) random variables drawn from the uniform distribution on $[0, 1]$. Compute $P(A)$ where A is defined as the event

$$A = \{|X - Y| > a\} \cap \{|X - Z| > a\} \cap \{|Z - Y| > a\}.$$

Express $P(A)$ as a function of a . Note that A is the event that the samples of X, Y, Z are spaced at least a apart from each other.

Let $B = \{Z \leq Y \leq X\}$.

$$P(A|B) = \int_{2a}^1 dx \int_a^{x-a} dy \int_0^{y-a} dz = \frac{(1-2a)^6}{6}.$$

There are 6 permutations of X, Y, Z and each has this probability. So, $P(A) = 6P(B|A) = (1-2a)^3$.

Note that the three $\{|X - Y| \geq a\}$, $\{|X - Z| \geq a\}$, $\{|Z - Y| \geq a\}$ are not independent. If you solved the problem by assuming that they were independent, you get the right answer by coincidence but the solution is not valid.

$P(A) = (1-2a)^3$
