

Dark Matter Heating vs. Rotochemical Heating in Old Neutron Stars

Keisuke Yanagi (University of Tokyo)

Based on Koichi Hamaguchi, Natsumi Nagata, KY [arXiv: 1904.04667, 1905.02991]

June 25th, 2019 @ KEK

Introduction/Motivation



Dark matter search

Weakly Interacting Massive Particle (WIMP)

- DM candidate which has standard model weak interaction
- Typical mass range: $m \sim 100 \text{ GeV} - 1 \text{ TeV}$

Limitation of DM direct detection


- DM + nucleus \rightarrow DM + nucleus
- **Neutrino floor limits ultimate sensitivity**
- **Insensitive to Inelastic scattering ($\Delta M < 100 \text{ keV}$)** $\longleftrightarrow \Delta M \sim O(100) \text{ MeV}$ for pure Higgsino/Wino



Dark matters accrete in neutron stars

- Consider weakly interacting massive particles (WIMPs)
- WIMPs scatter with nucleons and lose their kinetic energy
- Then they are trapped by a NS, and annihilate to SM particles

[Kouvaris, 0708.2362]




Energy injection

$$L_{\text{WIMP}} = (\text{Energy flux}) \times (\text{Capture probability})$$
$$\sim \rho_{\text{DM}} v_{\text{DM}} \pi b_{\text{max}}^2 \sim 1 \text{ for } \sigma_n \gtrsim 10^{-45} \text{ cm}^2$$

Dark matter kinetic/mass energy heats NS

DM scattering/annihilation deposits energy in NS


→ Late time heating!



- w/o WIMP : $T_s < 1000$ K @ $t > 10$ Myr
- w/ WIMP : $T_s \sim 3000$ K @ $t > 10$ Myr
- Sensitive to $\Delta M \lesssim 1$ GeV

Can we really see DM heating?


The observation suggests presence of **other heating mechanisms**



- Old NSs can be hotter than the cooling prediction or DM heating prediction
 - Several old ($t > 10$ Myr) pulsars have $T_s \sim 10^5$ K
 - WIMP cannot heat up a NS to $T_s \sim 10^5$ K
- An old NS is **not always warm**; it sometimes remains cold
 - PSR2144-3933: $T_s < 4 \times 10^4$ K @ $t \sim 100$ Myr

Can we really see DM heating?


The observation suggests presence of **other heating mechanisms**



- Old NSs can be hotter than the cooling prediction or DM heating prediction
 - Several old ($t > 10$ Myr) pulsars have $T_s \sim 10^5$ K
 - WIMP cannot heat up a NS to $T_s \sim 10^5$ K
- An old NS is **not always warm**; it sometimes remains cold
 - PSR2144-3933: $T_s < 4 \times 10^4$ K @ $t \sim 100$ Myr

Can we really see DM heating?

The observation suggests presence of **other heating mechanisms**



- Old NSs can be hotter than the cooling prediction or DM heating prediction
 - Several old ($t > 10$ Myr) pulsars have $T_s \sim 10^5$ K
 - WIMP cannot heat up a NS to $T_s \sim 10^5$ K
- An old NS is **not always warm**; it sometimes remains cold
 - PSR2144-3933: $T_s < 4 \times 10^4$ K @ $t \sim 100$ Myr

Can we really see DM heating?

Theoretically, several heating mechanisms are suggested

[Gonzalez & Reisenegger, 1005.5699]

- Non-equilibrium beta process (rotochemical heating)
 - Superfluid vortex heating
 - Decay of magnetic field
 - e.t.c...
- } Maybe responsible, but theoretically less clear...

If these mechanisms keep NS at $T_s \sim 10^5$ K, DM heating may be hidden...

Can we really see the DM heating? If so, we want to clarify the condition!

Can we really see DM heating?

Theoretically, several heating mechanisms are suggested

[Gonzalez & Reisenegger, 1005.5699]

- **Non-equilibrium beta process (rotochemical heating)**

← Inevitable for pulsars, our focus

- Superfluid vortex heating
- Decay of magnetic field
- e.t.c...

} Maybe responsible, but theoretically less clear...

If these mechanisms keep NS at $T_s \sim 10^5$ K, DM heating may be hidden...

Can we really see the DM heating? If so, we want to clarify the condition!

Outline

- Minimal cooling theory
- Rotochemical heating
- Results
 - We compare theory and observation including rotochemical heating [\[KY, Koichi Hamaguchi, Natsumi Nagata, arXiv: 1904.04667\]](#)
 - We discuss the possibility to search DM under the rotochemical heating [\[Koichi Hamaguchi, Natsumi Nagata, KY, arXiv: 1905.02991\]](#)

Minimal cooling of a neutron star

Basics of NS

- NS core consists of n, p, e, μ

- They are Fermi-degenerate


$$p_{F,n} \sim O(100) \text{ MeV}$$

$$p_{F,e,p,\mu} \sim O(10) \text{ MeV}$$

- Birth temperature $\sim 10^{11} \text{ K}$, and

quickly cools to $T < 10^{10} \text{ K}$

- **NS is cold system**



Nucleon superfluidity in NS

Cooper pairing occurs due to the attractive nuclear force



At $T < T_c^{(N)} \sim 10^{8-9}$ K

Superfluid in NS core

- Proton singlet pairing (1S_0)
- Neutron triplet pairing (3P_2)

Superfluid in NS crust (not important for thermal evolution)


- Neutron singlet pairing (1S_0)



[Figures from Page et al. (2013)]

Energy gap

Once Cooper pairing occurs, the **energy gap** appears in the spectrum



[Figures from Page et al. (2013)]

Energy spectrum near Fermi surface


$$\epsilon_N(\mathbf{p}) \simeq \mu_N + \text{sign}(p - p_{F,N}) \sqrt{\Delta_N^2 + v_{F,N}^2 (p - p_{F,N})^2}$$

Pairing gap models

The effects of superfluidity depends on momentum dependence of gap

$$\Delta_N = \Delta_N(\mathbf{k}_F, T = 0)$$


Proton 1S_0 pairing models



$$T_c^{(p)} = O(1) \times 10^9 \text{ K}$$

$$\Delta_N(k_F, T = 0) \simeq 1.764 k_B T_c^{(N)}$$

Neutron 3P_2 pairing models



$$T_c^{(n)} \sim 10^8 - 10^9 \text{ K}$$

$$\Delta_N(k_F, \cos \theta = 0, T = 0) \simeq 1.188 k_B T_c^{(N)}$$

[Figures from Page et al. (2013)]

Thermal evolution

Thermal evolution is governed by the energy conservation law


$$C \frac{dT}{dt} = -L_\nu - L_\gamma$$

Heat capacity (n, p, e, μ)

Neutrino luminosity


Photon luminosity:

$$L_\gamma = 4\pi R^2 \sigma_B T_s^4$$



Direct Urca process

Neutrino emission from beta decay and its inverse **on Fermi surface**



$$L_\nu^{\text{DU}} \propto T^6$$





Direct Urca does not operate unless the NS is very heavy

- Nucleons and leptons are strongly degenerate; $p_\nu \sim T \ll p_{F,n,p,\ell}$

- Momentum conservation requires

$$p_{F,p} + p_{F,\ell} > p_{F,n}$$

- Since $p_F^3 \propto n$, direct Urca requires **high p, e, μ density** ($M \gtrsim 2 M_\odot$ for APR EOS)




Modified Urca process


Threshold of direct Urca is relaxed by spectator nucleon

$$n + N \rightarrow p + N + \ell + \bar{\nu}_\ell$$

$$p + N + \ell \rightarrow n + N + \nu_\ell$$



- **Beta equilibrium** is usually assumed: $\mu_n = \mu_p + \mu_\ell$
- Before Cooper pairing: Luminosity = $L_\nu^{\text{MU}} \propto T^8$
- After Cooper pairing: modified Urca is highly suppressed



$$f \sim e^{-\Delta_N/T} \text{ for } Q_{M,N\ell} = \int \left[\prod_{j=1}^4 \frac{d^3 p_j}{(2\pi)^3} \right] \frac{d^3 p_\ell}{(2\pi)^3} \frac{d^3 p_\nu}{(2\pi)^3} (2\pi)^4 \delta^4(P_f - P_i) \cdot \epsilon_\nu \cdot \frac{1}{2} \sum_{\text{spin}} |\mathcal{M}_{M,N\ell}|^2$$


$$\times [f_1 f_2 (1 - f_3)(1 - f_4)(1 - f_\ell) + (1 - f_1)(1 - f_2) f_3 f_4 f_\ell],$$

Cooper pair-breaking and formation (PBF)



The Cooper pairing triggers rapid neutrino emission (called PBF)

[Flowers et al. (1976)]

- Pair-breaking**



- Pair-formation**





Pair breaking occurs by thermal disturbance
 —————→ efficient while $T \sim \Delta$

PBF dominates L_ν for $T < T_c$

Minimal cooling

Minimal cooling paradigm explains many NSs surface temperatures

[Page et al., astro-ph/0403657; Gusakov et al., astro-ph/0404002; Page et al., 0906.1621]




Different lines = Different gap/envelope model

- Direct Urca is not included
- $t < 10 - 100$ yr: Equilibrium modified urca $n + N \leftrightarrow p + N + \ell \pm \bar{\nu}_{\ell}$
- $10 - 100 \text{ yr} < t < 10^5 \text{ yr}$: PBF $[\tilde{N}\tilde{N}] \rightarrow \tilde{N}\tilde{N}$ $\tilde{N}\tilde{N} \rightarrow [\tilde{N}\tilde{N}] + \nu\bar{\nu}$
- $t > 10^5 \text{ yr}$: Photon emission $L_{\gamma} = 4\pi R^2 \sigma_B T_s^4$

Minimal cooling

Minimal cooling paradigm explains many NSs surface temperatures

[Page et al., astro-ph/0403657; Gusakov et al., astro-ph/0404002; Page et al., 0906.1621]



Different lines = Different gap/envelope model

- Direct Urca is not included
- $t < 10 - 100$ yr: Equilibrium modified urca $n + N \leftrightarrow p + N + \ell \pm \bar{\nu}_{\ell}$
- $10 - 100 \text{ yr} < t < 10^5 \text{ yr}$: PBF $[\tilde{N}\tilde{N}] \rightarrow \tilde{N}\tilde{N}$ $\tilde{N}\tilde{N} \rightarrow [\tilde{N}\tilde{N}] + \nu\bar{\nu}$
- $t > 10^5 \text{ yr}$: Photon emission $L_{\gamma} = 4\pi R^2 \sigma_B T_s^4$

Rotochemical heating

Pulsar spin-down

Spin-down: pulsar is rotating, and its rotation is gradually slowing down

$$P \sim 10^{-3} - 1 \text{ s}$$

$$\dot{P} \sim 10^{-20} - 10^{-13}$$

- Spin-down is caused by the **magnetic dipole radiation**

$$\frac{d\Omega}{dt} = -k\Omega^3 \quad \longrightarrow \quad \Omega(t) = \frac{2\pi}{\sqrt{P_0^2 + 2P\dot{P}t}}$$


$$k \propto B^2 \propto P\dot{P}$$

$$B \sim 3.2 \times 10^{19} (P\dot{P}/s)^{1/2} \text{ G}$$

- Centrifugal force is continuously decreasing
 - NS tries to change local pressure $P(r)$
 - Number density of each particle has to be rearranged
 - **(Hydrostatic) Equilibrium density is time-dependent**

$$n_i^{\text{eq}} = n_i^{\text{eq}}(t)$$

$$i = n, p, e, \mu$$




Hydrostatic equilibrium is not guaranteed

Each particle goes to new equilibrium $n_i^{\text{eq}}(t)$ by modified Urca process

If (modified) Urca is too slow, it cannot catch up with change of $n_i^{\text{eq}}(t)$

(Schematic picture)

$$n_i = n_i^{\text{eq}} + \delta n_i$$



Heat production through entropy production

- (Hydrostatic) equilibrium density is changing, so **chemical (or beta) equilibrium is also not guaranteed**

Measure of departure from beta equilibrium:

$$\eta_\ell = \mu_n - \mu_p - \mu_\ell = \delta\mu_n - \delta\mu_p - \delta\mu_\ell$$

\uparrow
 $\mu_i = \mu_i^{\text{eq}} + \delta\mu_i$

- Departure from chemical equilibrium generates heat

$$C \frac{dT^\infty}{dt} = -L_\nu^\infty - L_\gamma^\infty + L_H^\infty$$

$$L_H^\infty = \sum_{\ell=e,\mu} \sum_{N=n,p} \int dV \eta_\ell \cdot \Delta\Gamma_{M,N\ell} e^{2\Phi(r)}$$

$$dE^\infty = T^\infty dS + \sum_{i=n,p,e,\mu} \mu_i^\infty dN_i = -(L_\nu^\infty + L_\gamma^\infty) dt$$

Heat production through entropy production

- (Hydrostatic) equilibrium density is changing, so **chemical (or beta) equilibrium is also not guaranteed**

Measure of departure from beta equilibrium:

$$\eta_\ell = \mu_n - \mu_p - \mu_\ell = \delta\mu_n - \delta\mu_p - \delta\mu_\ell$$


$$\mu_i = \mu_i^{\text{eq}} + \delta\mu_i$$

- Departure from chemical equilibrium generates heat

$$C \frac{dT^\infty}{dt} = -L_\nu^\infty - L_\gamma^\infty + L_H^\infty$$

$$L_H^\infty = \sum_{\ell=e,\mu} \sum_{N=n,p} \int dV \eta_\ell \cdot \Delta\Gamma_{M,N\ell} e^{2\Phi(r)}$$

$$dE^\infty = T^\infty dS + \sum_{i=n,p,e,\mu} \mu_i^\infty dN_i = -(L_\nu^\infty + L_\gamma^\infty) dt$$



Effect of superfluidity



Nucleon superfluidity generates threshold

[Petrovich & Reisenegger, 0912.2564]

$$\Delta_{\text{th}} = \min\{3\Delta_n + \Delta_p, \Delta_n + 3\Delta_p\}$$

$\eta_\ell > \Delta_{\text{th}}$: heating begins

Larger $\Delta \sim$ larger $\eta \rightarrow$ hotter NS



Previous work incorporates only neutron triplet pairing

[González-Jiménez et al, 1411.6500]

We include both neutron and proton pairing

Rotochemical heating vs. observation


Two categories of observed pulsars

Ordinary pulsars and XDINSs $P \sim 1 - 10 \text{ s}, \dot{P} \sim 10^{-(15-13)}$

- **Ordinary pulsars** : most NSs belong to this class
- **XDINSs** (X-ray dim Isolated Neutrons Stars) : large magnetic field, thought to be remnants of magnetar

Millisecond pulsars $P \sim 1 \text{ ms}, \dot{P} \sim 10^{-20}$

- **Millisecond pulsars** : small rotational period and its derivative, formed by recycle of a binary system





$$\Omega(t) = \frac{2\pi}{\sqrt{P_0^2 + 2P\dot{P}t}}$$

$$B \sim 3.2 \times 10^{19} \left(\frac{P\dot{P}}{s} \right)^{1/2} \text{ G}$$

Gap models we use

The profile of pairing gap is one major source of uncertainty



$$\Delta_{\text{th}} = \min\{3\Delta_n + \Delta_p, \Delta_n + 3\Delta_p\}$$


- Large gap delays the beginning of rotochemical heating
- Heating power is stronger for larger gap

Results

Observed pulsars are explained for various choice of gap models

MSP


$M = 1.4 M_{\odot}$, $P_0 = 1$ ms




$$\Omega(t) = \frac{2\pi}{\sqrt{P_0^2 + 2P\dot{P}t}}$$

Ordinary pulsars & XDINSs

neutron: a, proton: CCDK, $P_0 = 1$ ms



neutron: a, proton: CCDK, $P_0 = 10$ ms




Results

Observed pulsars are explained for various choice of gap models

MSP

$M = 1.4 M_{\odot}$, $P_0 = 1$ ms




$$\Omega(t) = \frac{2\pi}{\sqrt{P_0^2 + 2P\dot{P}t}}$$


Increasing P_0 significantly suppresses rotochemical heating

Ordinary pulsars & XDINSs

neutron: a, proton: CCDK, $P_0 = 1$ ms




neutron: a, proton: CCDK, $P_0 = 10$ ms



DM heating vs. rotochemical heating

DM heating rate

DM accretion



Rate of DM hitting the NS

$$\dot{N} \simeq \pi b_{\text{max}}^2 v_{\text{DM}} (\rho_{\text{DM}}/m_{\text{DM}})$$

Heating luminosity

$$L_H^\infty = e^{2\Phi(R)} \dot{N} m_{\text{DM}} [\chi + (\gamma - 1)]$$

gravitational redshift factor \rightarrow

fraction of ann. energy into heat \rightarrow


$\frac{1}{\sqrt{1 - v_{\text{esc}}^2}}$

$\left\{ \begin{array}{l} = 1 \text{ for all annihilation into heat} \\ = 0 \text{ for no annihilation or all DM ann. into (e.g.) neutrinos} \end{array} \right.$

DM heating vs. rotochemical heating

DM heating effect is visible if the initial period is sufficiently large!

Ordinary pulsar: $P = 1 \text{ s}$ $\dot{P} = 10^{-15}$




$P_0 \lesssim 7 \text{ ms}$: DM heating < rotochemical heating

We can see the DM effect!

Uncertainty from superfluid gap models

- Critical P_0 depends on the choice of gap models
- **(DM heating) \gg (rotochemical heating) for $P_0 \gtrsim 100$ ms indep. of gap models**
- Recent studies of NS birth period suggest $P_0 = O(100)$ ms

[Popov & Turolla, 1204.0632; Noutsos et.al., 1301.1265; Igoshev & Popov, 1303.5258; Faucher-Giguere & Kaspi, astro-ph/0512585; Popov et al., 0910.2190; Gullo'n et al., 1406.6794, 1507.05452; Mu'ller et al., 1811.05483]



Summary


Summary

- It is known that DM heating can heat up a old NS
- We point out that DM heating may be hidden by other NS heating mechanisms
- Among proposed heating mechanisms, rotochemical heating is inevitable for any pulsar
- We compare the prediction of rotochemical heating to observations including both neutron and proton pairing gaps
- We then find that if the initial spin period is long enough, DM heating is stronger than rotochemical heating


Backup

Millisecond pulsars

Can we explain hot MSPs?



- $P = 5.8 \text{ ms.}$
- $\dot{P} = 5.7 \times 10^{-20}.$




- Two old hot MSPs are explained for various choice of gap models
- **Including both proton and neutron gap enhances heating**

Ordinary pulsars and XDINSs

Can the same setup explain other NS temperatures?



- $P = 1$ s.
- $\dot{P} = 1 \times 10^{-15}$.



- Many ordinary pulsars and XDINSs are also explained
- XDINSs are warmer, but may be explained by systematic uncertainties or heating caused by strong magnetic field

Initial spin period is a key parameter

$$P_0 = 10 \text{ ms}$$




[KY, Koichi Hamaguchi, Natsumi Nagata, arXiv: 1904.04667]

- Heating is weakened for longer initial period
- Old and cold NS is explained by assuming they had long initial period

Threshold of heating

Superfluidity makes threshold for rotochemical heating

For simplicity, consider direct Urca: $n \rightarrow p + e + \bar{\nu}_e$ $p + e \rightarrow n + \nu_e$



For modified Urca $\Delta_{\text{th}} = \min\{3\Delta_n + \Delta_p, \Delta_n + 3\Delta_p\}$

Neutron star envelope


Envelope: composed of light elements (H, He, C,...) and heavy elements (Fe)

Large temperature gradient exists


$$\frac{T}{10^9 \text{ K}} \sim 0.1288 \times \left(\frac{(T_s/10^6 \text{ K})^4}{g_{14}} \right)^{0.455}$$

[Gudmundsson et al. (1983)]

surface gravity [$10^{14} \text{ cm s}^{-2}$]



More accurate relation is available [Potekhin et al. (1997)]



Characterized by

$$\eta = g_{14}^2 \frac{\Delta M}{M}$$

mass of light elements

[Figure from Page et al. (2004)]