

-Project 1-

A Computational Study of the  
Ground-State Energy for Bosonic  
N-Particle Systems



Yngve Mardal Moe

Norges Miljø- og Biovitenskapelige Universitet

Fabian Maximilian Faulstich

Universitetet i Oslo

Teacher: Prof. Dr. Morten Hjorth-Jensen



# Abstract

Here comes the abstract...



# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Hard Sphere Bose Gas' Quantum Mechanical Model</b>	<b>1</b>
2.1	The Trapped Hard Sphere Bose Gas . . . . .	2
2.2	The Trial Function . . . . .	2
<b>3</b>	<b>Variational Monte Carlo Method and Statistical Analysis</b>	<b>3</b>
3.1	Variational Monte Carlo Method . . . . .	3
3.1.1	Metropolis Sampling . . . . .	5
3.1.2	Importance Sampling . . . . .	8
3.2	Statistical Analysis . . . . .	11
3.2.1	Bootstrapping . . . . .	12
3.2.2	Blocking . . . . .	13
<b>4</b>	<b>Implementation</b>	<b>15</b>
4.1	Analytical Solutions . . . . .	16
4.1.1	Harmonic Oscillator Potential . . . . .	16
4.1.2	Quantum Force for the Harmonic Oscillator Potential . . . . .	18
4.1.3	The Local Energy in the full Potential . . . . .	18
4.2	The Repulsive Interaction . . . . .	20
4.2.1	Change units of Length . . . . .	20
<b>5</b>	<b>Numerical Results</b>	<b>21</b>
<b>6</b>	<b>Conclusion and Perspectives</b>	<b>21</b>

## List of Figures

## List of Tables

# 1 Introduction

It was more than 80 years ago that Paul Dirac made his famous statement:

*The underlying physical laws necessary for the mathematical theory of a large part of physics and the whole of chemistry are [...] completely known [1].*

However, the development of *approximate practical methods of applying quantum mechanics*, demanded by Dirac in the same breath, is still a highly active field of research intersecting physics, chemistry, applied mathematics and computer science. In order to obtain results that can be utilized in practice, an extremely high accuracy of such practical methods is needed. On the other hand, solving the governing equation of quantum mechanics, the molecular Schrödinger equation, is an exceedingly high-dimensional and thus computationally demanding problem. Due to this very unfortunate combination of demanded high accuracy on the one side and the computational bad scaling on the other, small to medium-sized quantum chemical problems are still pushing the limits of commonly available computational resources.

In this project we focus on  $N$ -particle bosonic systems, i.e., particles that follow *Bose–Einstein statistics*. We present a computational study of the system’s ground-state energy using statistical methods. This work is outlined as follows:

We start by introducing the considered bosonic system in a more detailed, and for our approaches suitable, way in Section 2. We describe the quantum mechanical model we use to quantify hard sphere Bose gas in an elliptic trap (this covers in particular the special case of a spherical trap). In Section 3 we present the Monte Carlo method and the used statistical tools in a formal and mathematical rigorous way. We introduce Markov processes, in discrete and continuous time, as the mathematical concept on which we build the Monte Carlo method and two sampling methods, namely, the metropolis sampling and the importance sampling. The later will make excessive use of the Fokker–Planck and the Langevin equation. As the focus of this work is on a computational study, we do not introduce the concept of stochastic differential equations, however, we present a *quick and dirty* derivation of the Fokker–Planck equation via the *Kramers–Moyal* expansion. To present our computed data with a proper evaluation of the statical errors we introduce the *Bootstrap* and the *Blocking* method. In Section 4 we present the implementation used in this project. We hereby follow the project description, starting with deriving analytic expressions for the *local energy* in different potentials. In Section 5 we present our numerical results.

## 2 The Hard Sphere Bose Gas’ Quantum Mechanical Model

The aim of this project is to evaluate the ground state energy of a trapped, hard sphere Bose gas for different numbers of particles considering a specific trial wave function  $\psi_T$ . This trial wave function is used to study the sensitivity of condensate and non-condensate properties to the hard sphere radius and the number of particles. We start this section by introducing the quantum mechanical model used to describe a trapped hard sphere Bose Gas. Note that this model involves a *formal error*, i.e., an error that follows from

not knowing the exact theory to describe the physical behavior of atomic and subatomic particles.

## 2.1 The Trapped Hard Sphere Bose Gas

In this work we study the ground state energy of a trapped, hard sphere Bose gas, i.e., a gas composed of bosons, which have an integer value of spin and obey *Bose–Einstein statistics*. The trap we use in this project is a *spherical* (S) or an *elliptical* (E) harmonic trap in one, two and finally three dimensions, with the latter given by

$$V_{ext}(r) = \begin{cases} \frac{1}{2}m\omega_{ho}^2 r^2 & (S) , \\ \frac{1}{2}m[\omega_{ho}^2(x^2 + y^2) + \omega_z^2 z^2] & (E) . \end{cases} \quad (2.1)$$

The considered two-body Hamiltonian is formally defined by

$$H\psi(r) = \sum_i^N \left( \frac{-\hbar^2}{2m} \nabla_i^2 + V_{ext}(r_i) + \sum_{i<j}^N V_{int}(r_i, r_j) \right) \psi(r) , \quad (2.2)$$

where  $\omega_{ho}^2$  defines the *trap potential strength* and  $r = (r_1, \dots, r_N)$ . In the case of the elliptical trap,  $\omega_{ho}$  is the trap frequency in the *perpendicular plane*, i.e., the *x-y*-plane, and  $\omega_z$  is the frequency in the *z*-direction. We represent the boson's interaction by a pairwise, repulsive potential

$$V_{int}(|r_i - r_j|) = \begin{cases} \infty & |r_i - r_j| \leq a , \\ 0 & |r_i - r_j| > a , \end{cases} \quad (2.3)$$

where  $a$  is the so-called *hard-core diameter* of the bosons. This potential describes non-interacting bosons modeled as hard spheres of radius  $a/2$  described as point mass at position  $r_i$ , i.e., the distance of two bosons positioned at  $r_i$  and  $r_j$  can not be smaller than  $a$ . Consequently,  $V_{int}(|r_i - r_j|)$  is zero if the bosons are separated by a distance  $|r_i - r_j|$  greater than  $a$  but infinite if they attempt to come within a distance  $|r_i - r_j| \leq a$ , i.e., their volumes overlap.

## 2.2 The Trial Function

The trial wave function is used to study the sensitivity of condensate and non-condensate properties to the hard sphere radius and the number of particles. For the ground state with  $N$  atoms it is given by

$$\psi_T(r) = \psi_T(r_1, r_2, \dots, r_N, \alpha, \beta) = \prod_i g(\alpha, \beta, r_i) \prod_{i<j} f(a, |r_i - r_j|) , \quad (2.4)$$

where  $\alpha$  and  $\beta$  are variational parameters. The single-particle wave function  $g$  is proportional to the harmonic oscillator function for the ground state, i.e.,

$$g(\alpha, \beta, r_i) = \exp(-\alpha(x_i^2 + y_i^2 + \beta z_i^2)) . \quad (2.5)$$

The mean square vibrational amplitude of a single boson at  $T = 0K$  in the trap (2.1) is  $\langle x^2 \rangle = \hbar/2m\omega_{ho}$  so that  $a_{ho} \equiv (\hbar/m\omega_{ho})^{\frac{1}{2}}$  defines the characteristic length of the trap. For spherical traps we set  $\beta = 1$  and for non-interacting bosons we have  $\alpha = 1/2a_{ho}^2$ , setting  $a = 0$ . The *correlation wave function* is given by

$$f(a, |r_i - r_j|) = \begin{cases} 0 & , \text{ for } |r_i - r_j| \leq a , \\ (1 - \frac{a}{|r_i - r_j|}) & , \text{ for } |r_i - r_j| > a . \end{cases} \quad (2.6)$$

### 3 Variational Monte Carlo Method and Statistical Analysis

One of the main challenges in computational physics lies in the high dimensionality of the considered problems. To illustrate this we consider a single water molecule reduced to the electronic problem by the Born–Oppenheimer approximation [2, 3]. The spacial discretization of this three-dimensional ten-electron problem is in  $\mathcal{O}(n^{30})$ , where  $n$  describes the discretization points. Hence, for a rough discretization of  $n = 10^3$  the storage of this simple molecule already exceeds the estimated number of particles in the universe. This exponential growth with respect to the dimensionality is also known as *curse of dimensionality*. Subsequently, we introduce computational methods breaking this fatal growth behavior.

#### 3.1 Variational Monte Carlo Method

The main object of study will be the system’s energy, by means of the *Copenhagen interpretation* [4], given by the Hamilton operator’s expectation value, i.e.,

$$\langle H \rangle_\psi = \frac{\int_\Omega \psi^*(R) H \psi(R) dR}{\|\psi\|} .$$

We emphasize that in most cases  $\Omega$  will be high dimensional, e.g., for nuclear problems  $\Omega = \mathbb{R}^{3N} \times \{\pm 1/2\}^N \times \{\pm 1/2\}^N$  describing the spacial-, spin- and isospin-coordinates of the nucleons. This high dimensionality prohibits a use of conventional numerical integration schemes like the Gaussian quadrature. There are various alternative many-body methods trying to break the curse of dimensionality, however, their success depends strongly on the system’s physical properties.

The method chosen for this work is the *variational Monte-Carlo method* (vmc). Before going into further detail, we notice that the system’s energy is defined by the Rayleigh–Ritz variational principle [5], i.e.,

$$E_0 = \min_{\|\psi\|=1} \langle \psi, H \psi \rangle = \langle H \rangle_{\psi_0} .$$

Hence, by choosing any trial wave function  $\psi_T$  with  $\|\psi_T\| = 1$  we get  $E_0 \leq \langle H \rangle_{\psi_T}$ . Using special properties of the Hamilton operator<sup>1</sup> we can expand any trial function in terms

---

<sup>1</sup>Careful, linear self-adjoint and bounded operators do not necessarily have an eigen-basis. However, in finite dimensional Hilbert spaces (as relevant for computations) this is always true.



of eigenstates, i.e.,

$$\psi_T = \sum_i a_i \psi_i ,$$

with  $H\psi_i = E_i\psi_i$ . Without loss of generality we assume  $\|\psi\| = 1$  which implies

$$E_0 \leq \frac{\sum_i |a_i|^2 E_i}{\sum_i |a_i|^2} .$$

We see that by varying the trial function's parameters  $\{a_i\}$ , the Rayleigh–Ritz variational principle yields the ground state solution. Note, that in most cases the wave function has small extension coefficients in large parts of the configuration space, i.e., a straightforward computation using homogeneously distributed points in the configuration space will most likely not be efficient. Consequently, we will use the *importance sampling* combined with the *Metropolis* algorithm.

Subsequently we describe the underlying theory of Monte Carlo methods, which will be used in this project. Monte-Carlo methods are based on the *strong law of large numbers*:

**Etemadi, 1981 [6]:** *Let  $(X_n)_{n \in \mathbb{N}} \subseteq L^1(\Omega, \mathcal{A}, \mathbb{P})$  be a sequence of pairwise independent, identically distributed random variables. Then*

$$\frac{1}{N} \sum_{n=1}^N X_n \xrightarrow{a.s.} \mathbb{E}(X_1) ,$$

as  $n \rightarrow \infty$ .

**Remark 1.** *In the above theorem, we used the standard notation for probability theory: The function space  $L^1(\Omega, \mathcal{A}, \mathbb{P})$  is the quotient space with respect to the kernel of the  $L^1$ -norm of real-valued  $\mathbb{P}$ -integrable functions  $f : \Omega \rightarrow \mathbb{R}$  with the underlying sigma-algebra  $\mathcal{A}$ .*

*Almost sure convergence or strong convergence, denoted by  $\xrightarrow{a.s.}$ , means that*

$$\mathbb{P} \left( \lim_{N \rightarrow \infty} X_n - \mathbb{E}(X_1) \right) = 0 ,$$

*i.e., the set on which  $\lim_{N \rightarrow \infty} X_n$  differs from  $\mathbb{E}(X_1)$  is of  $\mathbb{P}$ -measure zero.*

*The above cited version is a stronger version of the strong law of large numbers proven by Kolmogorov in 1930 [7] as it only presumes pairwise independence of the random variables.*

For a chosen trial function  $\psi_T(R; a) = \sum_i a_i \psi_i(R)$  we define the probability density function

$$P(R; a) = \frac{|\psi_T(R; a)|^2}{\|\psi_T(\cdot; a)\|}$$

and observe that

$$\begin{aligned}\langle H \rangle_{\psi_T(\cdot; a)} &= \int_{\Omega} \frac{\psi_T(R; a)^* H \psi_T(R; a)}{\|\psi_T(\cdot; a)\|} dR = \int_{\Omega} \frac{|\psi_T(R; a)|^2}{\|\psi_T(\cdot; a)\|} \frac{H \psi_T(R; a)}{\psi_T(R; a)} dR \\ &= \int_{\Omega} P(R; a) E_L(R; a) dR ,\end{aligned}$$

where  $E_L(R; a) = H \psi_T(R; a) / \psi_T(R; a)$  denotes the *local energy*. We see that the last term corresponds to an integration with respect to the probability measure  $\mathbb{P}$  induced by the density  $P$ . By **Etemadi** this yields

$$\frac{1}{N} \sum_{n=1}^N P(R_i; a) E_L(R_i; a) \xrightarrow{a.s} \langle H \rangle_{\psi_T(\cdot; a)} .$$

We emphasize, that in the setting of Monte–Carlo methods, the number of pairwise identically distributed, independent variables  $N$  corresponds to the number of Monte–Carlo samples.

GO INTO DETAILS, HOW OUR COMPUTATIONS WORK!!!  
MENTION THE VARIANCE = 0

### 3.1.1 Metropolis Sampling

We start with a so-called brute-force sampling using the *metropolis sampling*, i.e., sampling based on the Metropolis–Hastings algorithm [8, 9]. As workhorse of *Markov-Chain Monte-Carlos methods*, the Metropolis–Hastings algorithm (also called Metropolis algorithm) is based on the following described theory of *Markov chains*. Given the unknown ground-state solution density  $p = |\psi_0|^2 / \|\psi_0\|$  (also called *the target* of the Metropolis algorithm), which is defined on the sample space  $\Omega$ , and computable up to a multiplying constant (due to the Rayleigh–Ritz principle  $p \propto \tilde{p}$ ). The Metropolis–Hastings algorithm then proposes a generic way to construct a Markov chain on  $\Omega$  that is *ergodic* and *stationary* with respect to  $p$  and therefore converges by the ergodic theorem in distribution to  $p$  [10, 11]. Due to its simplicity and versatility, the Metropolis–Hastings algorithm is the most used Markov-Chain Monte–Carlos method. This makes it the first solution to consider for intractable situations after homogeneously distributed sampling algorithms (cf. [12]). The main motivation for using Markov chains is that they provide *shortcuts* in cases where generic sampling requires too much effort from the experimenter. Rather than aiming at the big picture, like homogeneously distributed sampling algorithm, Markov chains construct a progressive picture of the target distribution, proceeding by local exploration of the state space  $\Omega$  until all the regions of interest have been uncovered.

We start this section by briefly introducing the necessary concept of Markov-chains:

Markov chains are one of the most important classes of stochastic processes, as they appear in various time-evolution models. In the following, we restrict the introduction to discrete time-homogeneous Markov chains on a finite dimensional sample set. A sequence of  $\Omega'$ -valued random variable  $(X_n)_{n \in \mathbb{N}}$  is said to fulfill the *Markov property* if for all  $k \in \mathbb{N}$  and all  $\omega_1, \omega_2, \dots, \omega_{k+1} \in \Omega'$  the following holds:

$$\mathbb{P}(X_{k+1} = \omega_{k+1} | X_k = \omega_k, X_{k-1} = \omega_{k-1}, \dots, X_1 = \omega_1) = \mathbb{P}(X_{k+1} = \omega_{k+1} | X_k = \omega_k) ,$$

where  $\mathbb{P}(\cdot | \cdot)$  describes the *conditional probability*. Hence, if a stochastic process fulfills the Markov property, the  $k + 1$ -step is only depending on the  $k$  step, i.e., the Markov chain has no memory<sup>2</sup>. As we only consider Markov chains on a finite dimensional sample set, we use the class of stochastic matrices  $\mathcal{S}$  to describe a Markov process, i.e.,

$$\mathcal{S} = \left\{ S \in \mathbb{R}^{m \times m} | S_{i,j} \in [0, 1] \forall i, j \in \{1, \dots, m\}, \sum_{i=j}^m S_{i,j} = 1 \forall i \in \{1, \dots, m\} \right\} .$$

We then define a Markov chain as follows:

*Let  $S \in \mathcal{S}$  be a stochastic matrix. A sequence of  $\Omega'$ -valued random variable  $(X_n)_{n \in \mathbb{N}}$  is called a time-homogeneous Markov chain with transition matrix  $S$  if for all  $k \in \mathbb{N}$  and all  $\omega_1, \omega_2, \dots, \omega_{k+1} \in \Omega'$  with  $\mathbb{P}(X_k = \omega_k, X_{k-1} = \omega_{k-1}, \dots, X_1 = \omega_1) > 0$  the following holds:*

$$\mathbb{P}(X_{k+1} = \omega_{k+1} | X_k = \omega_k, X_{k-1} = \omega_{k-1}, \dots, X_1 = \omega_1) = S_{\omega_k, \omega_{k+1}} .$$

*The matrix entries  $S_{i,j}$  are called transition probabilities and the initial distribution  $\mu$  is defined by  $\mu(\omega) = \mathbb{P}(X_1 = \omega)$  for  $\omega \in \Omega'$ .*

An important feature of Markov chain is a *stationary distribution*, i.e., a distribution  $\nu$  that stays invariant under the Markov chain and therewith fulfilling the matrix equation  $\nu S = \nu$ . The existence and uniqueness of such stationary distributions depends strongly on the Markov chain. The for the Metropolis algorithm central theorem, is the *ergodic theorem for time-homogeneous Markov chains on a finite dimensional sample space* [10, 11]. It states that for an *aperiodic* and *irreducible* Markov chain with transition matrix  $S$ ,  $\nu S^n \xrightarrow{\mathbb{P}} \mu$  for  $n \rightarrow \infty$  and any initial distribution  $\nu$ , with  $\mu$  being the stationary distribution. We again used the standard notation from probability theory, where  $\nu S^n \xrightarrow{\mathbb{P}} \mu$  denotes the convergence in probability, i.e., for all  $\varepsilon > 0$  we have

$$\lim_{n \rightarrow \infty} \mathbb{P}(|\nu S^n - \mu| \geq \varepsilon) = 0 .$$

Note that demanding the Markov chain with transition matrix  $S$  to be *aperiodic* and *irreducible* is the same as demanding the existence of  $n \in \mathbb{N}$  such that  $S_{i,j}^n > 0$  for all  $i, j$ ,

---

<sup>2</sup>This is the major and most important difference between Markov chains and Martingales – another class of important stochastic processes.

i.e., the Markov chain is ergodic. The aim is now to construct an ergodic Markov chain that converges to the ground-state solution density  $p$ , we follow hereby the reasoning presented in [13]:

The Markov chain  $(X_n)_{n \in \mathbb{N}}$  returned by the Metropolis-Hastings method is indeed such that it converges to  $p$  [8]. This means that the chain can be considered as a sample, albeit a dependent sample, and approximately distributed from  $p$ . Due to the Markovian nature of the simulation, the first values are highly dependent on the initial distribution and therewith from  $X_1$ . Consequently, these are removed from the sample as *warm-up*, also called *burn-in*. While there are very few settings where the time when the chain reaches stationarity can be determined [14], there is no need to look for such an instant since the empirical average

$$\frac{1}{N} \sum_{n=1}^N P(X_i; a) E_L(X_i; a)$$

converges almost surely if the Markov chain is ergodic. Comparing this to (3.1) it implies that, in theory, simulating a Markov chain is intrinsically equivalent to the pairwise independent identically distributed simulation from the target. Note, that the difference being in a loss of efficiency, i.e., it is necessary to simulate more terms to achieve a given variance for the above Monte Carlo estimator.

The Metropolis-Hastings algorithm associated with the target density  $p$  requires the choice of a conditional density  $q$  also called *proposal* or *candidate kernel*. The transition  $t \rightarrow t + 1$  is then given by Algorithm 1.

---

**Algorithm 1** Metropolis–Hasting

---

- 1: **procedure** CREATE METROPOLIS-HASTING MARKOV CHAIN
- 2:   Given  $X_t = x_t$
- 3:   Generate  $Y_t \sim q(y|x_t)$
- 4:   Take

$$X_{t+1} = \begin{cases} Y_t, & \text{with probability } \rho(x_t, Y_t), \\ x_t, & \text{with probability } 1 - \rho(x_t, Y_t), \end{cases}$$

where

$$\rho(x, y) = \min \left\{ 1, \frac{\tilde{p}(y)q(x|y)}{\tilde{p}(x)q(y|x)} \right\}.$$


---

Then, as shown in [8], this transition preserves the stationary density  $p$  if the chain is irreducible, i.e.,  $q$  has a wide enough support to eventually reach any region of the state space  $\Omega'$  with positive mass under  $p$ . As mentioned before, a sufficient condition is that  $q$  is positive everywhere. The very nature of accept-reject step introduced by Metropolis et al. [8] is therefore sufficient to turn a simulation from an almost arbitrary proposal

density  $q$  into a generation that preserves  $p$  as the stationary distribution. This appears somewhat surprising but works indeed, as shown in theoretical description above. In practice, the performances of the algorithm are highly dependent on the choice of the transition  $q$ , since some choices see the chain unable to converge in a manageable time. In this work we use  $X_t$  perturbed by a normal distribution, i.e.,  $Y_t \sim X_t + \varepsilon_t$  with  $\varepsilon_t \sim \mathcal{N}(0, 1)$ .

### 3.1.2 Importance Sampling

Despite the Metropolis–Hasting sampling’s simplicity, we replace this brute force algorithm with a walk in coordinate space biased by the trial wave function. This *importance sampling* is based on a stochastic differential equation, namely the *Langevin equation*, i.e.,

$$\frac{d^2x}{dt^2} = -\lambda \frac{dx}{dt} + \eta(t),$$

with the position  $x$  of a particle of mass one. The noise term  $\eta(t)$  follows a Gaussian probability distribution, where the correlation function depends on the physical problem. For this work, however, we are not interested in a solution of a Langevin equation for a particular realization of the fluctuating force. Rather, we are interested in correlation functions of the slow variables after averaging over the fluctuating force. As shown in [15, 16, 17], such correlation functions can be described by the *Fokker–Planck equation*, a deterministic differential equation. Unlike Langevin equations, the Fokker–Planck equation describes a time dependent probability density of a random variable and not the random variable itself. We shall see that the important sampling is not as far from the previously described Metropolis–Hasting sampling, which was derived from the theory of discrete time-homogeneous Markov chains on a finite dimensional sample set.

Subsequently, we present a quick and dirty derivation of the Fokker–Planck equation, by shortly dropping the mathematical rigor of this work. We now consider time-continuous Markov chains, i.e.,

*The stochastic process  $(X_t)_{t \geq 0}$  is called a time continuous Markov chain, if for all  $0 \leq s_0 < s_1 < \dots < s_n < s$  and all possible events  $\omega_0, \omega_1, \dots, \omega_n, \omega_i, \omega_j \in \Omega'$  the following holds*

$$\mathbb{P}(X_{t+s} = \omega_j | X_s = \omega_i, X_{s_n} = \omega_n, \dots, X_{s_0} = \omega_0) = \mathbb{P}(X_t = \omega_j | X_0 = \omega_i)$$

The Markov property and the definition of the conditional probability we derive straightforwardly the Chapman-Kolmogorow equality. From the Markov property we obtain

$$\begin{aligned} \mathbb{P}(X_{t_3} = \omega_3, X_{t_2} = \omega_2, X_{t_1} = \omega_1) \\ = \mathbb{P}(X_{t_1} = \omega_1) \mathbb{P}(X_{t_2} = \omega_2 | X_{t_1} = \omega_1) \mathbb{P}(X_{t_3} = \omega_3 | X_{t_2} = \omega_2) \end{aligned}$$

which is equivalent to

$$\mathbb{P}(X_{t_3} = \omega_3, X_{t_1} = \omega_1) = \mathbb{P}(X_{t_1} = \omega_1) \int_{\Omega'} \mathbb{P}(X_{t_2} = \omega_2 | X_{t_1} = \omega_1) \mathbb{P}(X_{t_3} = \omega_3 | X_{t_2} = \omega_2) d\omega_2 .$$

Division by  $\mathbb{P}(X_{t_1} = \omega_1)$  yields the *Chapman-Kolmogorow* equality

$$\mathbb{P}(X_{t_3} = \omega_3 | X_{t_1} = \omega_1) = \int_{\Omega'} \mathbb{P}(X_{t_2} = \omega_2 | X_{t_1} = \omega_1) \mathbb{P}(X_{t_3} = \omega_3 | X_{t_2} = \omega_2) d\omega_2$$

and considering the times  $t' \leq t' + \tau \leq t$  this reads

$$\mathbb{P}(X_t = \omega | X_{t'} = \omega') = \int_{\Omega'} \mathbb{P}(X_t = \omega | X_{t'+\tau} = \omega'') \mathbb{P}(X_{t'+\tau} = \omega'' | X_{t'} = \omega') d\omega'' . \quad (3.1)$$

By construction of the delta distribution, we further have

$$\mathbb{P}(X_{t'+\tau} = \omega'' | X_{t'} = \omega') = \int_{\Omega'} \mathbb{P}(X_{t'+\tau} = y | X_{t'} = \omega') \delta(y - \omega'') dy . \quad (3.2)$$

Inserting the Taylor expansion of the delta distribution, i.e.,

$$\delta(y - \omega'') = \sum_{n=0}^{\infty} \frac{(y - \omega')^n}{n!} \left( \frac{\partial}{\partial \omega'} \right)^n \delta(\omega' - \omega'')$$

into Eq. (3.2), we obtain

$$\mathbb{P}(X_{t'+\tau} = \omega'' | X_{t'} = \omega') = \left( 1 + \sum_{n=0}^{\infty} \frac{1}{n!} \underbrace{\int_{\Omega'} (y - \omega')^n \mathbb{P}(X_{t'+\tau} = y | X_{t'} = \omega') dy}_{=M_n(\omega', t', \tau)} \left( \frac{\partial}{\partial \omega'} \right)^n \right) \delta(\omega' - \omega'') .$$

Inserting this into Eq. (3.1) yields

$$\mathbb{P}(X_t = \omega | X_{t'} = \omega') = \mathbb{P}(X_t = \omega | X_{t'+\tau} = \omega') + \sum_{n=0}^{\infty} \frac{1}{n!} M_n(\omega', t', \tau) \left( \frac{\partial}{\partial \omega'} \right)^n \mathbb{P}(X_t = \omega | X_{t'+\tau} = \omega') .$$

As mentioned above, the Fokker-Planck equation is a deterministic differential equation describing the probability density of a random variable. To meet that end, we take a closer look at a differential quotient

$$\frac{\partial \mathbb{P}(X_t = \omega | X_{t'} = \omega')}{\partial t'} = - \lim_{\tau \rightarrow 0} \frac{\mathbb{P}(X_t = \omega | X_{t'} = \omega') - \mathbb{P}(X_t = \omega | X_{t'+\tau} = \omega')}{\tau} ,$$

together with Eq. (3.1.2) this can be written as

$$\frac{\partial \mathbb{P}(X_t = \omega | X_{t'} = \omega')}{\partial t'} = - \lim_{\tau \rightarrow 0} \frac{1}{\tau} \sum_{n=0}^{\infty} \frac{1}{n!} M_n(\omega', t', \tau) \left( \frac{\partial}{\partial \omega'} \right)^n \mathbb{P}(X_t = \omega | X_{t'+\tau} = \omega') .$$

Using the Taylor expansion of  $M_n(\omega', t', \tau)$ , i.e.,

$$\frac{M_n(\omega', t', \tau)}{n!} = 0 + D^{(n)}(\omega', t') \tau + \mathcal{O}(\tau^2), \quad \text{with } D^{(n)}(\omega', t') = \lim_{\tau' \rightarrow 0} \mathbb{E}((X_{t'+\tau'} - X_{t'})^n) |_{X_{t'} = \omega'}$$

yields the differential equation

$$\frac{\partial \mathbb{P}(X_t = \omega | X_{t'} = \omega')}{\partial t'} = -\mathcal{L}_{KM}^\dagger(\omega', t') \mathbb{P}(X_t = \omega | X_{t'} = \omega') \quad (3.3)$$

with the *Kramers–Moyal operator*

$$\begin{aligned} \mathcal{L}_{KM}(\omega', t') &= \sum_{n=0}^{\infty} \left( -\frac{\partial}{\partial \omega'} \right)^n D^{(n)}(\omega', t') , \\ \mathcal{L}_{KM}^\dagger(\omega', t') &= \sum_{n=0}^{\infty} D^{(n)}(\omega', t') \left( \frac{\partial}{\partial \omega'} \right)^n . \end{aligned}$$

Similar to Eq. (3.3) with find the analog deterministic differential equation

$$\frac{\partial P(\omega, t)}{\partial t} = \mathcal{L}_{KM}(\omega, t) P(\omega, t) \quad (3.4)$$

describing the time evolution of the probability density  $P(\omega, t)$ . The mayor question arising at this point is under which circumstances this expansion terminates. The answer is given by Pawulas theorem[15]:

*For positive transition probabilities  $\mathbb{P}(\omega, t | \omega', t')$  the expansion in Eq. (3.4) terminates after one or two terms. If it does not terminate after the second term it never terminates.*

If the expansion in Eq. (3.4) terminates after the second term, the resulting differential equation is called a Fokker-Planck equation:

$$\frac{\partial P(\omega, t)}{\partial t} = -\frac{\partial}{\partial \omega} D^{(1)}(\omega, t) P(\omega, t) + \frac{\partial^2}{\partial \omega^2} D^{(2)}(\omega, t) P(\omega, t) = \mathcal{L}_{FP} P(\omega, t) ,$$

with the Fokker-Planck operator

$$\mathcal{L}_{FP} = -\frac{\partial}{\partial \omega} D^{(1)}(\omega, t) + \frac{\partial^2}{\partial (\omega)^2} D^{(2)}(\omega, t) .$$

The term  $D^{(1)}$  is called *drift coefficient* and is responsible for the time-evolution of the random variables mean value. The term  $D^{(2)}$  is called *diffusion coefficient* and is responsible for the time-evolution of the random variables standard deviation. In this work we consider a constant diffusion coefficient, in atomic unties,  $D^{(2)} = 1/2$ . Consequently, the convergence to a stationary probability density, i.e.,  $\partial P(\omega, t)/\partial t = 0$ , then corresponds to the solution of

$$\frac{\partial^2}{\partial (\omega)^2} P(\omega, t) = D^{(1)}(\omega, t) \frac{\partial}{\partial \omega} P(\omega, t) + P(\omega, t) \frac{\partial}{\partial \omega} D^{(1)}(\omega, t) .$$

We assume the drift coefficient to be of the form  $D^{(1)}(\omega, t) = g(\omega, t) \partial P(\omega, t)/\partial \omega$ . Together with *Random–Nikodym theorem* this yields

$$\frac{\partial^2 P(\omega, t)}{\partial (\omega)^2} = g(\omega, t) \left( \frac{\partial P(\omega, t)}{\partial \omega} \right)^2 + P(\omega, t) g(\omega, t) \frac{\partial^2 P(\omega, t)}{\partial \omega^2} + P(\omega, t) \frac{\partial g(\omega, t)}{\partial P(\omega, t)} \left( \frac{\partial P(\omega, t)}{\partial \omega} \right)^2 .$$

For  $g(\omega, t) = 1/P(\omega, t)$  this equality is fulfilled. Indeed,

$$\frac{\partial^2 P(\omega, t)}{\partial(\omega)^2} = \frac{1}{P(\omega, t)} \left( \frac{\partial P(\omega, t)}{\partial \omega} \right)^2 + \frac{\partial^2 P(\omega, t)}{\partial \omega^2} + P(\omega, t) \left( -\frac{1}{P^2(\omega, t)} \right) \left( \frac{\partial P(\omega, t)}{\partial \omega} \right)^2 .$$

As in this work  $P = |\psi_T|^2$  this yields

$$D^{(1)}(\omega, t) = \frac{1}{P} \frac{\partial P}{\partial \omega} = 2 \frac{1}{\psi_T} \nabla \psi_T = F ,$$

which is also known as *quantum force*. As the drift coefficient describes the movement of the mean value, it is this term that forces the taken samples, also called walkers, into regions of the configuration space where the trial wave function  $\psi_t$  is large. This increases the simulation's efficiency in contrast to the here considered Metropolis–Hastings sampling where the samples are normally distributed, i.e, they have the same probability of moving in every direction. Using the Green's functions theory, we obtain a solution of above described Fokker-Planck equation, implying the following description of transition probabilities

$$G(x, y, \Delta t) = \frac{1}{(2\pi\Delta t)} \exp \left( -\frac{(y - x - \frac{\Delta t F(x)}{2})^2}{2\Delta t} \right) ,$$

where we used  $D^{(2)} = 1/2$ .

Using the Langevin equation we can describe the random variable of the transition probability using an *explicit Euler procedure* with time step size  $\Delta t$ , i.e.,

$$Y_t = X_t + D^{(2)} F(X_t) \Delta t + \xi \sqrt{\Delta t} .$$

Using this random variable and the transition probability described by the Fokker–Planck's Greens function in the Metropolis–Hastings algorithm we obtain the importance sampling, see Alg. 2.

### 3.2 Statistical Analysis

In this section, we describe the statistical tools that will be used for a proper evaluation of our data's statistical errors. The computational results are obtained by using Monte Carlo simulations. These simulations can be treated as computational experiments and its results can therefore be analyzed with the same statistical tools used for laboratory experiments. We hereby focus on two *resampling method*, i.e. methods that estimate the precision of sample statistics (medians, variances, percentiles), namely:

- The *bootstrap technique* also called *bootstrapping*, where the estimation is based on drawing randomly with replacement from a set of data points.
- The *blocking method* also just *blocking*, where the samples are divided in sub-samples of certain size, allowing to use estimators for uncorrelated samples.



---

**Algorithm 2** Importance Sampling

---

- 1: **procedure** CREATE IMPORTANCE SAMPLING MARKOV CHAIN
- 2:     Given  $X_t = x_t$
- 3:     Generate  $Y_t \sim X_t + D^{(2)}F(X_t)\Delta t + \xi\sqrt{\Delta t}$
- 4:     Take

$$X_{t+1} = \begin{cases} Y_t, & \text{with probability } \rho(x_t, Y_t), \\ x_t, & \text{with probability } 1 - \rho(x_t, Y_t), \end{cases}$$

where

$$\rho(x, y) = \min \left\{ 1, \frac{G(x, y, \Delta t) |\psi_T(y)|^2}{G(y, x, \Delta t) |\psi_T(x)|^2} \right\}.$$

---

The estimators used in this work are a *mean estimator* defined by

$$\hat{\Theta}(X) = \hat{\Theta}(X_1, \dots, X_n) = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

and a *variance estimator* defined by

$$\hat{\varphi}(X) = \hat{\varphi}(X_1, \dots, X_n) = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 = S^2 X,$$

where  $X = (X_1, \dots, X_n)$  describes a set of samples.

### 3.2.1 Bootstrapping

In the wider sense of statistical analysis, any test that relies on random sampling with replacement is called bootstrapping. This technique allows the estimation of the sampling distribution and provides an error analysis assigning measures of accuracy defined in terms of bias, variance, confidence intervals, prediction error. The basic idea of bootstrapping is treating inference of the true probability distribution  $J$ , given the original data, as being analogous to inference of the empirical distribution of  $\hat{J}$ , given the resampled data. The accuracy of inferences regarding  $\hat{J}$  using the resampled data is assessable because  $\hat{J}$  is known. Assuming that  $\hat{J}$  is a *reasonable approximation* to the unknown distribution  $J$ , then the quality of inference on  $J$  can in turn be inferred. In a more informal way bootstrapping can be described as follows:

Assume we are interested in the mean value of  $X_1, \dots, X_K$  independently and identically  $J$ -distributed with  $K$  being very large. If  $K$  becomes too large, we have to restrict the evaluation to a subset of size  $n$ . Without loss of generality we denote this subset by  $X_1, \dots, X_n$  with  $n \ll K$ . From this sample the mean estimator yields a single estimate  $\bar{X} = \hat{\Theta}(X_1, \dots, X_n)$ . In order to reason about  $\Theta(X_1, \dots, X_K)$ , we need a sense of the variability of the mean  $\bar{X}$ . Bootstrapping now treats the sample  $X_1, \dots, X_n$  in the same way

as  $X_1, \dots, X_K$ . A straightforward bootstrap approach therefore involves sampling from  $X_1, \dots, X_n$  to form a new sample (called a *resample* or *bootstrap sample*) of the same size. Under the assumption that  $n$  is sufficiently large, there is practically zero probability that the sample will be identical to the original real sample. This process is repeated numerous times, and for each of these bootstrap samples the mean is computed, the so-called *bootstrap estimates*. This yields a histogram of bootstrap means providing an estimate of the unknown distribution's mean. How much the mean varies can then be deduced from this histogram and used to measure the quality of the mean computed from the original sample.

The above described method is also called *non-parametric Bootstrap*, and is used when the distribution  $J$  is unknown. We summarize this algorithm in the following pseudo-code Alg. (3):

---

**Algorithm 3** Non-parametric Bootstrap

---

- 1: **procedure** CREAT BOOTSTRAP SAMPLE HISTOGRAM
  - 2:   Draw histogram of  $X(P^*(X))$
  - 3:   **procedure** BOOTSTRAP SAMPLE
  - 4:     Draw from  $P^*(X)$ ,  $n$  random numbers  $\{X_i\} = X^*$
  - 5:     Compute  $\hat{\Theta}(X^*) = t(X^*)$
  - 6:     write  $t$  to memory
  - 7:   Use Bootstrap Samples to obtain the final histogram  $P^*(\hat{\Theta})$
- 

In the case the histogram  $P(X)$  is known, we resample from this histogram and do the Bootstrap sample steps from Alg. (3). This is then known as *parametric Bootstrap*.

### 3.2.2 Blocking

The Blocking method is used for correlated data samples, where we can no longer use the estimators for uncorrelated sample sets. In blocking methods, the sample set  $X$  is grouped into sets of blocks  $\{X^{(i)}\}$ , where  $X^{(i)}$  describes a blocked sample set. The idea is now to use a block size large enough such that the elements of the different blocks are independent, which allows us to apply the estimators for uncorrelated sample sets to the blocked sample sets. Note that various possibilities exist to block the original sample set. The blocking procedure used in this work is introduced in the sequel.

We start this section by describing the approach of blocking methods in a more descriptive way. Consider experimental data from a Monte Carlo simulation of step size  $\Delta t$ . In case of uncorrelated samples the mean's standard deviation is given by

$$\sigma = \sqrt{\hat{\varphi}(X)} . \quad (3.5)$$

The Monte Carlo simulation however, will most likely produce a correlated sample set. In this case the standard deviation needs to be corrected as follows

$$\sigma = \sqrt{\frac{1 + 2\tau/\Delta t}{n - 1} (\overline{X^2} - \bar{X}^2)} ,$$

where  $\tau$  is the so called *correlation time*, i.e., the time between a sample and the next uncorrelated sample. In case  $\Delta t \gg \tau$  the Monte Carlo samples will be uncorrelated and Eq. (3.5) can be used to compute the standard deviation. As mentioned before, it is most likely that  $\Delta t < \tau$ , in which case Eq. (3.5) is not valid anymore. Furthermore, we are facing the difficulty of not knowing the correlation time  $\tau$ . However, with blocking we are still able to determine a valid standard deviation of the mean  $\overline{X}$ . Using the uncorrelated expression Eq. (3.5), we compute the mean of each block  $\overline{X^{(i)}}$  with  $i = 1, \dots, n_b$  where  $n_b$  is the number of blocks, and calculate then the total mean and variance. The crucial part for this method is the block size, i.e., it must be so large that sample  $j$  of block  $i$  is not correlated with sample  $j$  of block  $i + 1$ . An optimal choice would therefore be the unknown correlation time. A quick and dirty solution to this problem is a manual fix of the block size:

A graphical representation of the standard deviation as a function of the block size will show a plateau region, i.e., a connected set of block sizes for which the standard deviation is constant. As soon as the standard deviation stops increasing, we know that the blocks are uncorrelated. This number can then be manually put into the blocking algorithm.

An alternative to this brute force fix of the blocking method, is the *automatic blocking*. We here proceed as follows:

For a given set of samples  $(X_1, \dots, X_n)$  with  $n = 2^d$  we create the block set

$$\begin{aligned} X^{(1)} &= \left( \frac{1}{2}(X_1 + X_2), \dots, \frac{1}{2}(X_{n-1} + X_n) \right) = (X_1^{(1)}, \dots, X_{n_1}^{(1)}) , \\ &\vdots \\ X^{(d)} &= \left( \frac{1}{2}(X_1^{(d-1)} + X_2^{(d-1)}), \dots, \frac{1}{2}(X_{n_{d-1}-1}^{(d-1)} + X_{n_{d-1}}^{(d-1)}) \right) = (X_1^{(d)}, \dots, X_{n_d}^{(d)}) . \end{aligned}$$

From these block we can compute various quantities like the mean and the variance, and many which will be used to derive an automatized blocking method. Before going into further detail of these quantities, however, we introduce some more statistical concepts. We define the so-called auto-variance  $f_d$  for  $m$  experiments of  $n$  samples by

$$f_d = \frac{1}{nm} \sum_{\alpha=1}^m \sum_{k=1}^{n-d} (x_{\alpha,k} - \langle x_m \rangle)(x_{\alpha,k+d} - \langle x_m \rangle) ,$$

where  $\langle x_m \rangle$  denotes the means over all samples, i.e.,

$$\langle x_m \rangle := \frac{1}{nm} \sum_{\alpha=1}^m \sum_{k=1}^n x_{\alpha,k} .$$

The total variance

$$\sigma_m = \frac{1}{mn^2} \sum_{\alpha=1}^m \sum_{k,l=1}^n (x_{\alpha,k} - \langle x_m \rangle)(x_{\alpha,l} - \langle x_m \rangle)$$

can then be written as

$$\sigma_m = \frac{\sigma^2}{n} + \frac{2}{n} \sum_{d=1}^{n-1} f_d ,$$

where

$$\sigma^2 = \frac{1}{mn} \sum_{\alpha=1}^m \sum_{k=1}^n (x_{\alpha,k} - \langle x_m \rangle)^2 .$$

For the sake of completeness note that the expression

$$\mathcal{E} := \frac{2}{n} \sum_{d=1}^{n-1} f_d$$

describes the covariance. For each blocked sample set  $X^{(i)}$  we can then compute the total variance

$$\mathbb{V}(\overline{X^{(i)}}) = \frac{\mathbb{V}(X^{(i)})}{n_i} + \mathcal{E}_i .$$

We then state the following proposition [CITE MARIUS...]

*If  $X_1, \dots, X_n$  is stationary, i.e., it exists a real number  $c$  such that  $\langle X_i \rangle = c$  for all  $i$ , and  $\mathbb{V}(X_i) < \infty$  for one  $i$ , then  $\mathcal{E}_i - \mathcal{E}_{i+1} = \text{cov}(X_{j+1}^{(i)}, X_j^{(i)})/n_i =: \gamma^i(1)/n_i$ .*

Furthermore, it is straightforward that  $\mathcal{E}_i$  is a monotone function in  $i$  if we assume that  $\gamma^k(1) > 0$  for all  $1 \leq k \leq d_1$ . Hence, if  $X^{(i)}$  has independent elements, then  $\mathcal{E}_k = 0$  and  $\gamma^k(1) = 0$  for all  $k \geq i$ . The inverse of this statement follows immediately as the covariance of independent samples is equal to zero. This yields the condition used to automatize the blocking method, namely, we stop blocking if  $\gamma^k(1) = 0$ . Before presenting the pseudo-code for the automatic blocking method, we give the following theorem [CITE MARIUS...] on which the procedure is based on:

*Let  $X_1, \dots, X_n$  with  $n = 2^d$  be stationary, with  $\mathbb{V}(X_i) < \infty$  for one  $i$  and  $\gamma(h) \rightarrow \infty$ . Then the following implication holds:  
If  $\gamma^k(1) = 0$  then*

$$M_j := \sum_{k=1}^{d-1} \frac{n_k((n_k - 1)(S^{(k)})^2/nk + \gamma^k(1))2}{(S^{(k)})^4}$$

*is chi-squared distributed ( $M_j \sim \chi^2$ ).* We finalize this section by presenting a pseudo-code for the automatic blocking method, see Alg. (4).

## 4 Implementation

In this section we present the implementation we used to derive next section's results. We start by deriving an analytic expression for the local energy.

---

**Algorithm 4** Automatic Blocking

---

```
1: procedure ESTIMATE  $\mathbb{V}(\overline{X})$ 
2:   Set  $i = 1$ 
3:   Compute  $(S^{(i)})^2, \hat{\gamma}^i(1)$ 
4:   Transform  $Q^{(i)} \rightarrow Q^{(i+1)}$ 
5:   if  $|Q^{(i)}| \leq 2$  then
6:     Compute  $M_j$  using  $\hat{\gamma}^j(1)$  and  $(S^{(j)})^2$ 
7:     Find the first  $k$  such that  $M_k \leq q_k$ 
8:     Return  $(S^{(k)})^2/n_k$ 
9:   else
10:    Go back to 2
```

---

## 4.1 Analytical Solutions

We observe that the considered Hamiltonian contains the Laplace operator, which can be approximated by the method of finite differences. However, we note that the used quantities are the local energy  $E_L$  and the quantum force  $F$ , which can be described analytically and therewith save computational time. We start by deriving the local energy in the harmonic oscillator potential, i.e.,  $a = 0$  implying  $f(a, |r_i - r_j|) = 1$  and  $V_{\text{int}}(|r_i - r_j|) = 0$ .

### 4.1.1 Harmonic Oscillator Potential

Following, we start with  $\beta = 1$ , i.e.,  $g(\alpha, \beta, r_i) = \exp(-\alpha r_i^2)$  and find the relevant local energies in one, two and three dimensions for one and  $N$  particles with the same mass. The trial function of a one dimensional single particle is given by

$$\psi_T(r_1, \alpha, 1) = \exp(-\alpha r_1^2) .$$

We then compute

$$\begin{aligned} H\psi_T(r_1, \alpha, 1) &= \left( \frac{-\hbar^2}{2m} \Delta + \frac{1}{2} m \omega_{ho}^2 r_1^2 \right) \exp(-\alpha r_1^2) \\ &= -\frac{\hbar^2}{2m} (4\alpha^2 r_1^2 - 2\alpha) \exp(-\alpha r_1^2) + V_{\text{ext}}(r_1) \exp(-\alpha r_1^2) , \end{aligned}$$

which yields

$$E_L(r_1, \alpha) = -\frac{\hbar^2}{2m} (4\alpha^2 r_1^2 - 2\alpha) + V_{\text{ext}}(r_1) .$$

Note that in atomic unites with  $\alpha = 1/2$  the local energy is given by  $E_L = 1/2$ , which is the ground-state energy of the one dimensional harmonic oscillator. This can be straightforwardly generalized to the  $n$ -dimensional case, where

$$H\psi_T(r_1, \alpha, 1) = -\frac{\hbar^2}{2m} (4\alpha^2 r_1^2 - 2\alpha n) \exp(-\alpha r_i^2) + V_{\text{ext}}(r_1) \exp(-\alpha r_1^2)$$

and therewith

$$E_L(r_1, \alpha) = -\frac{\hbar^2}{2m} (4\alpha^2 r_1^2 - 2\alpha n) + V_{\text{ext}}(r_1) .$$

For  $N$  particles in  $n$  dimensions the trial function reads

$$\psi_T(r_1, \dots, r_N, \alpha, 1) = \prod_{i=1}^N \exp(-\alpha r_i^2) = \exp\left(-\alpha \sum_{i=1}^N r_i^2\right) .$$

Similar to the one dimensional case we then compute

$$H\psi_T(r_1, \dots, r_N, \alpha, 1) = \left( \sum_{i=1}^N -\frac{\hbar^2}{2m} (4\alpha^2 r_i^2 - 2\alpha n) + V_{\text{ext}}(r_i) \right) \psi_T(r_1, \dots, r_N, \alpha, 1)$$

and therewith

$$E_L(r_1, \dots, r_N, \alpha) = \sum_{i=1}^N -\frac{\hbar^2}{2m} (4\alpha^2 r_i^2 - 2\alpha n) + V_{\text{ext}}(r_i) .$$

Assuming  $\beta \neq 1$  only influences the three dimensional case. We compute for one particle

$$H\psi_T(r_1, \alpha, \beta) = -\frac{\hbar^2}{2m} (4\alpha^2(x^2 + y^2 + \beta^2 z^2) - 2\alpha(2 + \beta)) \psi_T(r_1, \alpha, \beta) + V_{\text{ext}}(r_1) \psi_T(r_1, \alpha, \beta) ,$$

with the trial function

$$\psi_T(r_1, \alpha, 1) = \exp(-\alpha(x^2 + y^2 + \beta^2 z^2)) .$$

Hence,

$$E_L(r_1, \alpha, \beta) = -\frac{\hbar^2}{2m} (4\alpha^2(x^2 + y^2 + \beta^2 z^2) - 2\alpha(2 + \beta)) + V_{\text{ext}}(r_1) .$$

We conclude for the  $N$ - particle case for the trial function

$$\psi_T(r_1, \dots, r_N, \alpha, \beta) = \exp\left(-\alpha \sum_{i=1}^N (x_i^2 + y_i^2 + \beta^2 z_i^2)\right)$$

that

$$\begin{aligned} H\psi_T(r_1, \dots, r_n, \alpha, \beta) &= \sum_{i=1}^N -\frac{\hbar^2}{2m} (4\alpha^2(x_i^2 + y_i^2 + \beta^2 z_i^2) - 2\alpha(2 + \beta)) \psi_T(r_1, \dots, r_n, \alpha, \beta) \\ &\quad + V_{\text{ext}}(r_i) \psi_T(r_1, \dots, r_n, \alpha, \beta) , \end{aligned}$$

and therewith

$$E_L(r_1, \dots, r_n, \alpha, \beta) = \sum_{i=1}^N -\frac{\hbar^2}{2m} (4\alpha^2(x_i^2 + y_i^2 + \beta^2 z_i^2) - 2\alpha(2 + \beta)) + V_{\text{ext}}(r_i) .$$

#### 4.1.2 Quantum Force for the Harmonic Oscillator Potential

Another often used quantity is the drift coefficient of the Fokker–Planck equation, i.e., the quantum force  $F$ . As we do not want to approximate the gradient of the trial function for every sampling, we derive the expression analytically for the harmonic oscillator potential with  $\beta = 1$ . Then the trial function is again given by

$$\psi_T(r_1, \alpha, 1) = \exp(-\alpha r_1^2) ,$$

for which the quantum force is

$$F(r_1, \alpha, 1) = \frac{1}{\psi_T(r_1, \alpha, 1)} \nabla \psi_T(r_1, \alpha, 1) = -2\alpha r_1 .$$

As the gradient is a map  $\nabla : L^2(\mathbb{R}^{3N}) \rightarrow (L^2(\mathbb{R}^{3N}))^{3N}$  (using its weak formulation), we obtain the following quantum force for the  $N$ -particle case

$$F(r_1, \dots, r_N, \alpha, 1) = \frac{1}{\psi_T(r_1, \dots, r_N, \alpha, 1)} \nabla \psi_T(r_1, \dots, r_N, \alpha, 1) = -2\alpha (r_i)_{i=1}^N ,$$

where  $(r_i)_{i=1}^N = (x_1, y_1, z_1, x_2, \dots, z_N)^T$ .

#### 4.1.3 The Local Energy in the full Potential

Next we consider the full problem in three dimensions. The aim is to find an analytic expression for the derivative of the trial wave function

$$\frac{1}{\Psi_T(r)} \sum_i^N \nabla_i^2 \Psi_T(r) ,$$

with the trial function given by Eq. (2.4). Rewriting this trial function yields

$$\psi_T(r_1, \dots, r_N, \alpha, \beta) = \prod_{i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i<j} u(r_{ij}) \right) ,$$

where we have defined  $r_{ij} = |r_i - r_j|$  and

$$f(r_{ij}) = \exp \left( \sum_{i<j} u(r_{ij}) \right) ,$$

where  $u : \mathbb{R} \rightarrow \mathbb{R}$  with  $x \mapsto \ln \left( 1 - \frac{a}{x} \right)$ . Using the product rule we find

$$\begin{aligned} \nabla_k \psi_T(r_1, \dots, r_N, \alpha, \beta) &= (\nabla_k g(\alpha, \beta, r_k)) \prod_{i \neq k; i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i<j} u(r_{ij}) \right) \\ &+ \prod_{i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i<j} u(r_{ij}) \right) \sum_{i<j}^N \nabla_k u(r_{ij}) , \end{aligned} \tag{4.1}$$

where

$$\sum_{i < j}^N \nabla_k u(r_{ij}) = \frac{1}{2} \sum_{j=1}^N \sum_{i \neq j; i=1}^N \nabla_k u(|r_i - r_j|) .$$

Note that we slightly abuse the notation here, namely  $u(|r_i - r_j|)$  describes the map  $u(|\cdot - r_j|) : \mathbb{R}^3 \rightarrow \mathbb{R}$  evaluated in  $r_i$  as well as the map  $u(|r_i - \cdot|) : \mathbb{R}^3 \rightarrow \mathbb{R}$  evaluated in  $r_j$ . Consequently holds

$$\nabla_i u(|r_i - r_j|) = u'(|r_i - r_j|) \nabla_i |r_i - r_j| = \frac{a(r_i - r_j)}{(|r_i - r_j| - a)|r_i - r_j|^2} = \nabla_i u(|r_j - r_i|) , \quad (4.2)$$

and therewith

$$\sum_{i < j}^N \nabla_k u(r_{ij}) = \sum_{j \neq k; j=1}^N \nabla_k u(r_{kj}) .$$

From Eq. (4.1) we can then derive that the quantum force in the full potential is given by

$$F(r_1, \dots, r_N, \alpha, \beta) = 2 \left( \left( \frac{\nabla_i g(\alpha, \beta, r_i)}{g(\alpha, \beta, r_i)} \right)_{i=1}^N + \left( \sum_{j \neq i} \nabla_i u(|r_i - r_j|) \right)_{i=1}^N \right) .$$

Deriving another time, we find

$$\begin{aligned} \Delta_k \psi_T(r_1, \dots, r_N, \alpha, \beta) &= (\Delta_k g(\alpha, \beta, r_k)) \prod_{i \neq k; i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i < j} u(r_{ij}) \right) \\ &+ (\nabla_k g(\alpha, \beta, r_k)) \prod_{i \neq k; i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i < j} u(r_{ij}) \right) \sum_{j \neq k; j=1}^N \nabla_k u(r_{kj}) \\ &+ \prod_{i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i < j} u(r_{ij}) \right) \left( \sum_{j \neq k; j=1}^N \nabla_k u(r_{kj}) \right)^2 \\ &+ \prod_{i=1}^N g(\alpha, \beta, r_i) \exp \left( \sum_{i < j} u(r_{ij}) \right) \sum_{j \neq k; j=1}^N \Delta_k u(r_{kj}) . \end{aligned} \quad (4.3)$$

As for the local energy this needs to be divided by the trial function, we find the following expression for the second derivative

$$\begin{aligned} \frac{\Delta_k \psi_T(r_1, \dots, r_N, \alpha, \beta)}{\psi_T(r_1, \dots, r_N, \alpha, \beta)} &= \frac{\Delta_k g(\alpha, \beta, r_k)}{g(\alpha, \beta, r_k)} + \frac{\nabla_k g(\alpha, \beta, r_k)}{g(\alpha, \beta, r_k)} \sum_{j \neq k; j=1}^N \nabla_k u(r_{kj}) \\ &+ \sum_{i, j \neq k} \nabla_k u(r_{kj}) \nabla_k u(r_{kj}) + \sum_{j \neq k; j=1}^N \Delta_k u(r_{kj}) , \end{aligned} \quad (4.4)$$



with

$$\begin{aligned}
\Delta_k u(r_{kj}) &= \nabla_k \cdot \frac{r_k - r_j}{|r_k - r_j|} u'(|r_k - r_j|) \\
&= u''(|r_k - r_j|) + u'(|r_k - r_j|) \left( \frac{3}{|r_k - r_j|} - (r_k - r_j) \cdot \frac{(r_k - r_j)}{|r_k - r_j|^3} \right) \\
&= u''(|r_k - r_j|) + u'(|r_k - r_j|) \frac{2}{|r_k - r_j|} .
\end{aligned}$$

It remains to compute  $u'$  and  $u''$ . We already computed  $u'$  in Eq. (4.2), however, for the sake of completeness we list it again subsequently. We find

$$\begin{aligned}
u'(r_{ij}) &= \frac{a}{(r_{i,j} - a)r_{i,j}} , \\
u''(r_{ij}) &= -\frac{a}{((r_{i,j} - a)r_{i,j})^2} (2r_{ij} - a) .
\end{aligned}$$

This together with the analytic expression for the harmonic oscillator yields an analytic expression for the local energy.

## 4.2 The Repulsive Interaction

### 4.2.1 Change units of Length

We here consider a different unit of length, namely,  $r \rightarrow r/a_{ho}$ . This corresponds to introducing the energy in units of  $\hbar\omega_{ho}$ , i.e., the Hamiltonian transforms as  $H \rightarrow H/\hbar\omega_{ho}$ . The transformed Hamiltonian reads

$$\frac{1}{\hbar\omega_{ho}} H = \frac{1}{\hbar\omega_{ho}} \sum_{i=1}^N \frac{1}{2} \left( -\frac{\hbar^2}{m} \nabla_i^2 + m\omega_{ho}^2 (x_i^2 + y_i^2) + m\omega_z^2 z_i^2 \right) + \frac{1}{\hbar\omega_{ho}} V_{\text{int}}(|r_i - r_j|) . \quad (4.5)$$

As  $V_{\text{int}}$  is either zero or infinity it follows that

$$\frac{1}{\hbar\omega_{ho}} V_{\text{int}}(|r_i - r_j|) = V_{\text{int}}(|r_i - r_j|) .$$

We recall that

$$a_{ho} = \left( \frac{\hbar}{m\omega_{ho}} \right)^{1/2}$$

and simplify Eq. (4.5) as follows

$$\begin{aligned}
\frac{1}{\hbar\omega_{ho}} H &= \sum_{i=1}^N \frac{1}{2} \left( -\frac{\hbar}{m\omega_{ho}} \nabla_i^2 + \frac{m\omega_{ho}}{\hbar} (x_i^2 + y_i^2) + \frac{m}{\hbar\omega_{ho}} \frac{\omega_z^2}{\omega_{ho}} z_i^2 \right) + V_{\text{int}}(|r_i - r_j|) \\
&= \sum_{i=1}^N \frac{1}{2} \left( -a_{ho}^2 \nabla_i^2 + a_{ho}^{-2} (x_i^2 + y_i^2) + a_{ho}^{-2} \frac{\omega_z^2}{\omega_{ho}^2} z_i^2 \right) + V_{\text{int}}(|r_i - r_j|) .
\end{aligned}$$

Indeed, this corresponds to the change of length unit  $r \rightarrow r/a_{ho}$ . The Hamiltonian then reads

$$H = \sum_{i=1}^N \frac{1}{2} \left( -\nabla_i^2 + x_i^2 + y_i^2 + \gamma^2 z_i^2 \right) + V_{\text{int}}(|r_i - r_j|) ,$$

where we introduced the constant  $\gamma = \omega_z/\omega_{ho}$ .

## 5 Numerical Results

## 6 Conclusion and Perspectives

## References

- [1] Paul AM Dirac. Quantum mechanics of many-electron systems. *Proc. R. Soc. Lond. A*, 123(792):714–733, 1929.
- [2] John C Slater and Arnold Russek. Quantum theory of molecules and solids, vol. 1: Electronic structure of molecules. *American Journal of Physics*, 32(1):65–66, 1964.
- [3] Max Born and Robert Oppenheimer. Zur quantentheorie der molekeln. *Annalen der Physik*, 389(20):457–484, 1927.
- [4] Werner Heisenberg. *Die Kopenhager Deutung der Quantentheorie*. Stuttgart. Battenberg, 1963.
- [5] Harry Yserentant. A short theory of the rayleigh–ritz method. *Computational Methods in Applied Mathematics*, 13(4):495–502, 2013.
- [6] Nasrollah Etemadi. An elementary proof of the strong law of large numbers. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 55(1):119–122, 1981.
- [7] Andreï Kolmogorov. Sur la loi forte des grands nombres.
- [8] Nicholas Metropolis, Arianna W Rosenbluth, Marshall N Rosenbluth, Augusta H Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6):1087–1092, 1953.
- [9] W Keith Hastings. Monte carlo sampling methods using markov chains and their applications. *Biometrika*, 57(1):97–109, 1970.
- [10] James R Norris. *Markov chains*. Number 2. Cambridge university press, 1998.
- [11] Richard Durrett and R Durrett. *Essentials of stochastic processes*. Springer, 2016.
- [12] Christian P Robert, George Casella, and George Casella. *Introducing monte carlo methods with r*, volume 18. Springer, 2010.
- [13] Christian P Robert and George Casella. The metropolis—hastings algorithm. In *Monte Carlo Statistical Methods*, pages 231–283. Springer, 1999.
- [14] James P Hobert and Christian P Robert. A mixture representation of  $\pi$  with applications in markov chain monte carlo and perfect sampling. *Annals of Applied Probability*, pages 1295–1305, 2004.
- [15] Hannes Risken. The fokker-planck equation. methods of solution and applications, vol. 18 of. *Springer series in synergetics*, 301, 1989.

- [16] Ronald R Coifman, Ioannis G Kevrekidis, Stéphane Lafon, Mauro Maggioni, and Boaz Nadler. Diffusion maps, reduction coordinates, and low dimensional representation of stochastic systems. *Multiscale Modeling & Simulation*, 7(2):842–864, 2008.
- [17] Setsuo Ichimaru. Basic principles of plasma physics: a statistical approach. 1973.