

Topology - X400416

Yoav Eshel

March 9, 2021

Contents

1	Summary	2
2	Topological Spaces	2
3	Basis for a Topology	3
4	The Product Topology on $X \times Y$	4
5	The Subspace Topology	4
6	Closed Sets and Limit Points	5
7	Continuous Functions	6
8	The Product Topology	8
9	The Metric Topology	9
10	The Quotient Topology	9
11	Exercises	10
	Basis for a topology	10
	The Subspace Topology	12
	Closed Sets and Limit Points	13
	Continuous Functions	17
	Extra Exercises	18
	The Product Topology	19
	Metric Spaces	21
	The Quotient Topology	24
	Connected Spaces	26
	Connected Subspaces of the Real Line	27
	Components and Local Connectedness	28

These notes are based on Topology (2nd edition) by James R. Munkres.

1 Summary

2 Topological Spaces

The motivation behind defining a topological space is to generalize the notion of a metric space. Recall that a metric on a set X is a map $d : X \times X \rightarrow [0, \infty)$ that satisfies

- a. $d(x, y) = d(y, x)$
- b. $d(x, x) = 0$
- c. $d(x, y) > 0, x \neq y$
- d. $d(x, y) \leq d(x, z) + d(z, y)$

Then we say that a set $U \subset X$ is open if for all $x \in U$ and some $r > 0$

$$B(x, r) := \{y \in X \mid d(x, y) < r\} \subset U.$$

In other words, around every point in U there is a "ball" that is contained in U . In Analysis I one learns about continuity using the classic ε - δ definition which requires a metric. As it turns out, we don't really need a metric to define a continuous function, only open sets:

Definition. *A function between metric spaces is continuous if and only if the preimage of an open set is open.*

Using this definition, different seeming metric can yield the same notions of which functions are continuous! We call the collection of open subsets of X defined by some metric $d : X \times X \rightarrow [0, \infty)$ a **topology**. This open sets satisfy some important properties. Namely: (1) X and \emptyset are open, (2) arbitrary unions of open sets are open and (3) finite intersections of open sets is open. It turns out that a metric is not required to define a topology, only these three properties:

Definition. *Let X be a set. Then a topology on X is a set $\mathcal{T} \subset \mathcal{P}(X)$ such that*

- a. $\emptyset \in \mathcal{T}, X \in \mathcal{T}$
- b. *If $\{U_\alpha\} \subset \mathcal{T}$ then $\bigcup_\alpha U_\alpha \in \mathcal{T}$*
- c. *If $\{U_i\}_{i=0}^n \subset \mathcal{T}$ then $\bigcap_{i=0}^n U_i \in \mathcal{T}$*

A topological space is the pair (X, \mathcal{T})

Then we say that $U \subset X$ is open if $U \in \mathcal{T}$. Note that which sets are open depends on the topology, which might conflict with your notion of open set as defined above. For example, in the topology $\mathcal{P}(\mathbb{R})$ every subset of the real line is open, while in the topology $\{\emptyset, \mathbb{R}\}$ only the empty set and \mathbb{R} are open. We often say that U is open in X without giving a specific topology, which simply means that the statement that follows will hold for any topology we define on X and any element in that topology.

If \mathcal{T} and \mathcal{T}' are two topologies on X such that $\mathcal{T} \subseteq \mathcal{T}'$ then we say that \mathcal{T}' is **finer** (or **strictly finer** if the containment is proper) than \mathcal{T} . We similarly say that \mathcal{T} is **coarser** (or **strictly coarser**) than \mathcal{T}' . It might also be that case that two topologies are not **comparable**.

3 Basis for a Topology

Specifying topologies directly is often not possible, due the enormous size of many topologies. So we often define a topology using a smaller subset called a **basis**.

Definition. *If X is a set, then a **basis** of a topology is a collection \mathcal{B} of subsets of X such that*

- a. $\forall x \in X, \exists B \in \mathcal{B}$ such that $x \in B$
- b. *If $x \in B_1 \cap B_2$ with $B_1, B_2 \in \mathcal{B}$ then there exists $B_3 \in \mathcal{B}$ with $x \in B_3 \subset B_1 \cap B_2$.*

If \mathcal{T} is a topology generated by a basis \mathcal{B} then U is open if for all $x \in U$ there exists $B \in \mathcal{B}$ such that $x \in B \subset U$. Also $B \in \mathcal{T}$ for all $B \in \mathcal{B}$. The proof that \mathcal{T} is indeed a topology is not included. An alternative construction of a topology from a basis is given by the following lemma

Lemma 3.1. *Let X be a set and \mathcal{B} a basis for a topology \mathcal{T} . Then \mathcal{T} equals the collection of all unions of elements in \mathcal{B} .*

Using this lemma is sometime easier in practice; given a basis \mathcal{B} and $U \subset X$, if one can write U as a union of elements in \mathcal{B} then U is open.

We can also go in the reverse direction: from topology to a basis.

Lemma 3.2. *Let X be a topological space. If \mathcal{C} is a collection of open sets such that for each open set U and each $x \in U$ there is $C \in \mathcal{C}$ such that $x \in C \subset U$, then \mathcal{C} is a basis.*

When topologies are given in terms of basis, we can already determine which one is finer using the following criterion

Lemma 3.3. *Let \mathcal{B} and \mathcal{B}' be bases for the topologies \mathcal{T} and \mathcal{T}' respectively. Then $\mathcal{T} \subset \mathcal{T}'$ if and only if for each $x \in X$ and $B \in \mathcal{B}$ containing x there exists $B' \in \mathcal{B}'$ such that $x \in B' \subset B$.*

It might be tricky to remember the direction of the inclusion. One way to think about it is since \mathcal{T}' has more subsets of X it needs to have smaller basis elements.

Lastly, we define the notion of a **subbasis**.

Definition. A subbasis \mathcal{S} for a topology on X is a collection of subsets of X whose union equals X . The topology generated by \mathcal{S} is the collection of all unions of finite intersection of elements of \mathcal{S} .

To conclude this section we define 3 topologies on the real line using the notion of a basis:

- a. The **standard topology** generated by the collection of all open intervals (a, b) with $a < b$ (it is not a recursive definition. Here we use open in the familiar metric sense).
- b. The **lower limit topology** is generated by half-open intervals $[a, b)$. When \mathbb{R} is given in the lower limit topology we denote it \mathbb{R}_l .
- c. The **K-topology** is generated by open intervals and sets of the form $(a, b) - K$ where $K = \{1/n \mid n \in \mathbb{N}\}$. When \mathbb{R} is given in this topology we denote it \mathbb{R}_k .

One maybe surprising property is that both \mathbb{R}_l and \mathbb{R}_k are finer than the standard topology, but are not comparable with one another.

4 The Product Topology on $X \times Y$

Let X and Y be topological spaces. The product topology on $X \times Y$ is generated by the collection \mathcal{B} of all sets of the form $U \times V$ with U open in X and V open in Y . Alternatively, if \mathcal{B} is a basis for X and \mathcal{C} is a basis for Y , then

$$\mathcal{D} = \{B \times C \mid B \in \mathcal{B}, C \in \mathcal{C}\}$$

is a basis for $X \times Y$.

5 The Subspace Topology

If X is a topological space with topology \mathcal{T}_X and $Y \subset X$ then the subspace topology on Y is defined as

$$\mathcal{T}_Y = \{Y \cap U \mid U \in \mathcal{T}_X\}.$$

The fact that the collection \mathcal{T}_Y has all the properties of a topology follows from the \mathcal{T}_X being a topology. Then if \mathcal{B}_X is a basis for a topology on X , the basis of the subspace topology on Y is given by

$$\mathcal{B}_Y = \{Y \cap B \mid B \in \mathcal{B}_X\}.$$

Open sets in the subspace topology on Y are not necessarily open in X . If A is an open set in Y , then A is open in X if Y is open in X .

Another fun property of subspace topology is that it "commutes" with the product topology: If A is a subspace of X and B is a subspace of Y then the product topology on $A \times B$ is equal to the subspace topology on $A \times B$ as a subspace of $X \times Y$.

6 Closed Sets and Limit Points

If X is a topological space and $A \subset X$ then A is closed if $X - A$ is open. Closed sets have similar properties to open sets:

Theorem 6.1. *Let X be a topological space. Then*

- a. X and \emptyset are closed*
- b. Arbitrary intersections of closed sets are closed*
- c. Finite unions of closed sets are closed.*

One can just as well define a topology in terms of closed sets, but the definition using open sets is much more common. Of course, mathematics wouldn't be fun if there wasn't any space for confusion. In a topology, a set can be open, closed, neither or both. So don't think of sets as doors.

If Y is a subspace of a topological space X , then a closed set in X is not necessarily closed in Y . A set A is closed in Y if and only if it equals the intersection of a closed set of X with Y . This is easy to verify since if A is closed in Y then $Y - A$ is open in Y and so it equals $U \cap Y$ for some open set U in X . Then $X - U$ is closed and $A = Y \cap (X - U)$. The other direction is similarly proved.

Let $A \subset X$. Then the smallest closed set that contain A is called the closure of A and is denoted

$$\bar{A} = \bigcap_{\substack{C \text{ closed} \\ A \subset C}} C.$$

The smallest open set containing A is called the interior of A and is denoted

$$\text{Int}A = \bigcup_{\substack{U \text{ open} \\ U \subset A}} U.$$

Then clearly

$$\text{Int}A \subset A \subset \bar{A}.$$

A point $x \in X$ is in the closure of A if and only if for any basis element B containing x the intersection $A \cap B$ is non empty.

To add to the soup of metaphors, we define the **neighborhood** of $x \in X$ is an open set U containing x . Then a **limit point** of $A \subset X$ is defined as

Definition. A limit point of $A \subset X$ is a point $x \in A$ such that every neighborhood of x intersects A in a point different than x

In other words, x is a limit point of A if for every open set U containing x the intersection $U \cap (A - x)$ is non empty. Now denote the set of limit points of A by A' . Then $\bar{A} = A' \cup A$. This gives us an alternative condition for a set being closed. Namely, if $A' \subset A$ then $\bar{A} = A$ and so A is closed. Alternatively, if A is closed, then $\bar{A} = A$ and so $A' \subset A$. Hence a set is closed if and only if it contains all of its limit points.

We conclude this section with a definition of convergence in a topological space.

Definition. Let (X, \mathcal{T}) be a topological space. If x_n is a sequence in X then $x_n \rightarrow x$ as $n \rightarrow \infty$ if

$$\forall U \in \mathcal{T}, x \in U \exists N \in \mathbb{N} : \{x_n\}_{n \geq N} \subset U$$

Note that this definition does not require any metric. The downside is that limit are not necessarily unique and sequences can converge to any number of points. To avoid this horrific phenomenon, we add an extra condition to rid our definition of topological spaces of such situations.

Definition. A topological space X is called **Hausdorff space** if for every $x_1, x_2 \in X, x_1 \neq x_2$ there exists disjoint neighborhoods U_1 and U_2 that contain x_1 and x_2 respectively.

In essence, this means that we can distinguish between every two points in our topology. Limits of sequences in Hausdorff spaces are unique, which is quite pleasant.

7 Continuous Functions

Recall from the introduction that we defined a function $f : X \rightarrow Y$ to be continuous if for each open set V in Y the set

$$f^{-1}(V) = \{x \in X \mid f(x) \in V\}$$

is open in X . To prove continuity it suffices to show that for any basis element (or even any subbasis element!) $B \subset Y$, $f^{-1}(B)$ is open in X . Using this definition, a given function may or may not be continuous depending on the topologies specified for its domain and range. If we wish to emphasize it we can say that f is continuous relative to specific topologies. To illustrate this point, let \mathbb{R} denote the standard topology on the real line and let

$$f : \mathbb{R} \rightarrow \mathbb{R}_l \quad \text{and} \quad g : \mathbb{R}_l \rightarrow \mathbb{R}$$

be the identity maps, i.e. $x = f(x) = g(x)$ for all $x \in \mathbb{R}$. Then f is not continuous since $f^{-1}([a, b)) = [a, b)$ which is not open in the standard topology. However g is continuous since $g^{-1}((a, b)) = (a, b)$ which is open in \mathbb{R}_l .

There are other equivalent conditions for a function to be continuous

Theorem 7.1. *Let X and Y be topological spaces and $f : X \rightarrow Y$. The following are equivalent*

- a. f is continuous
- b. For every subset A of X it holds that $f(\overline{A}) \subset \overline{f(A)}$
- c. For every closed subset B of Y , the set $f^{-1}(B)$ is closed in X .
- d. For each $x \in X$ and each neighborhood V of $f(x)$ there is a neighborhood U of x such that $f(U) \subset V$.

A bijection $f : X \rightarrow Y$ is called a **homeomorphism** (the e is not a typo. It is indeed a different word to homomorphism!) if both f and f^{-1} are continuous. An isomorphism in algebra is a bijective map that preserve algebraic structure (between groups, rings, fields, etc.). In topology, a homeomorphism is a bijective map that preserve topological structure. However in algebra any homomorphism (without an e) that is one-to-one and onto is an isomorphism. This is, in general, not true in topology. There are continuous bijective maps whose inverses are not continuous. If $f : X \rightarrow Y$ is injective continuous map, and $g : X \rightarrow f(X)$ obtained by restricting the range of f is a homeomorphism, then we say that f is an **imbedding**.

There are a few important rules for constructing continuous functions

- a. Constant mappings are continuous
- b. Inclusion mappings are continuous (i.e. if A is a subspace of X and $i : A \rightarrow X$ is the identity then i is continuous)
- c. Composition of continuous maps are continuous
- d. Continuous maps restricted to a subspace of their domain are continuous
- e. Restricting or expanding the range of continuous maps gives continuous maps
- f. The map $f : X \rightarrow Y$ is continuous if X can be written as the union of open sets U_α such that $f|_{U_\alpha}$ is continuous for each α

Lemma 7.2. *(The pasting lemma) Let $X = A \cup B$ be a topological space and A, B are closed. Consider*

$$f : A \rightarrow Y \quad \text{and} \quad g : B \rightarrow Y$$

continuous such that $f(x) = g(x)$ for all $x \in A \cap B$. Then $h : X \rightarrow Y$ defined by

$$h(x) = \begin{cases} f(x), & x \in A \\ g(x), & x \in B \end{cases}$$

is continuous.

The lemma also holds if A and B are both open.

Lemma 7.3. (*Maps into products*) Let $f : A \rightarrow X \times Y$ be given by

$$f(a) = (f_1(a), f_2(a)).$$

Then f is continuous if and only if f_1 and f_2 are continuous.

Note that there is no useful criterion for maps out of products.

8 The Product Topology

In Section 4 we defined the topology on $X \times Y$ as the topology generated by products of open sets. However, when considering arbitrary products there are two ways to proceed. Let J be an arbitrary index set and X_α a topological space for each $\alpha \in J$. Then

Definition. The box topology on $\prod_{\alpha \in J} X_\alpha$ is generated by sets of the form

$$\prod_{\alpha \in J} U_\alpha$$

where each U_α is open in X_α .

Definition. The product topology on $\prod_{\alpha \in J} X_\alpha$ is generated by the subbasis of all subsets of the form $\pi_\beta^{-1}(U_\beta)$, $\beta \in J$ where $\pi_\beta : \prod_{\alpha \in J} X_\alpha \rightarrow X_\beta$ is the projection mapping. Therefore a basis for the product topology is the collection of subsets of the form

$$\prod_{\alpha \in J} V_\alpha$$

where

$$V_\alpha = \begin{cases} U_\alpha \text{ open in } X_\alpha, & \alpha \in I \\ X_\alpha, & \alpha \notin I \end{cases}$$

for some finite $I \subset J$.

It follows that these two topologies agree for finite cartesian products but differ for the infinite one. The product topology is quite important, while the box topology main use is as a counter example.

However there are some properties that hold for both topologies

- If A_α is a subspace of X_α for each α then $\prod A_\alpha$ is a subspace of $\prod X_\alpha$ when both products are given in the same topology.
- If each X_α is Hausdorff, then $\prod X_\alpha$ is Hausdorff in either topology.
- If $A_\alpha \subset X_\alpha$ for each α then

$$\prod \overline{A_\alpha} = \overline{\prod A_\alpha}$$

in either topology.

A property that does not hold for the box topology is this

Theorem 8.1. *Let $\prod X_\alpha$ have the product topology. Let $f : A \rightarrow \prod X_\alpha$ be given by*

$$f(a) = (f_\alpha(a))_{\alpha \in J}.$$

Then f is continuous if and only if f_α is continuous $\forall \alpha \in J$.

9 The Metric Topology

10 The Quotient Topology

Definition. *Let X and Y be topological space and $p : X \rightarrow Y$ be a surjective map. The map p is called a **quotient map** if a subset U of Y is open if and only if $p^{-1}(U)$ is open in X .*

We say that a subset C of X is **saturated** if every set $p^{-1}(y)$ that intersects C is contained in C . So we can also say the p is a quotient map if p is continuous and p maps saturated open sets of X to open sets of Y . We can also go the other way and use a quotient map to define a topology.

Definition. *If X is a space and A is and if $p : X \rightarrow A$ is a surjective map, then there exists exactly one topology on A relative to which p is a quotient map. We call this topology the **quotient topology** induced by p .*

Definition. *Let X be a space and X^* a partition of X . Let $p : X \rightarrow X^*$ be a surjective map that carries each point of X to the element of X^* containing it. In the quotient topology induced by p , the space X^* is called the **quotient space** X .*

Since X^* defines an equivalence relation on X , one can think of p as the map that send each element of X to its equivalence class. Hence a subset U of X^* is a collection of equivalence classes, and so U is open if the union of the equivalence classes contained in U is open in X .

11 Exercises

Basis for a topology

Exercise 13.1

Let X be a topological space; let $A \subset X$. Suppose that for each $x \in A$ there is an open set U such that $x \in U \subset A$. Show that A is open.

Proof. For every $x \in A$, let U_x denote the open set containing x such that $U_x \subset A$. Then $U = \bigcap_{x \in A} U_x \subset A$ since each U_x is contained in A . For the other inclusion, take $x \in A$. Then $x \in U$ since x is in U_x by definition. Hence $A \subset U$ and it follows that $A = U$. Since each U_x and arbitrary unions of open sets are open it follows that A is open. \square

Exercise 13.4

- If $\{\mathcal{T}_\alpha\}$ is a family of topologies on X , how that $\bigcap \mathcal{T}_\alpha$ is a topology on X . Is $\bigcup \mathcal{T}_\alpha$ a topology on X ?
- Let $\{\mathcal{T}_\alpha\}$ be a family of topologies on X . Show that there is a unique smallest topology containing all the all the collection of \mathcal{T}_α and a unique largest topology contained in all \mathcal{T}_α
- If $X = \{a, b, c\}$ let

$$\mathcal{T}_1 = \{\emptyset, X, \{a\}, \{a, b\}\} \quad \text{and} \quad \mathcal{T}_2 = \{\emptyset, X, \{a\}, \{b, c\}\}.$$

Find the smallest topology containing \mathcal{T}_1 and \mathcal{T}_2 and the largest topology contained in \mathcal{T}_1 and \mathcal{T}_2 .

Solution.

- (a) Since $\emptyset, X \in \mathcal{T}_\alpha$ for all α it follows that $\emptyset, X \in \bigcap \mathcal{T}_\alpha$. (b) If $U_\beta \in \bigcap \mathcal{T}_\alpha$, then $U_\beta \in \mathcal{T}_\alpha$ for all α and so $\bigcup U_\beta \in \mathcal{T}_\alpha$ for all α since \mathcal{T}_α is a topology. Hence $\bigcup U_\beta \in \bigcap \mathcal{T}_\alpha$. (c) If $U_1, U_2 \in \bigcap \mathcal{T}_\alpha$ then $U_1, U_2 \in \mathcal{T}_\alpha$ for all α and so $U_1 \cap U_2 \in \mathcal{T}_\alpha$ for all α . Therefore $U_1 \cap U_2 \in \bigcap \mathcal{T}_\alpha$. It follows by induction that $\bigcap \mathcal{T}_\alpha$ is closed under countable intersections. Hence an intersections of topologies is a topology.

Let $X = \{a, b, c\}$. Then $\mathcal{T}_1 = \{\emptyset, X, \{a\}\}$ and $\mathcal{T}_2 = \{\emptyset, X, \{b\}\}$ are topologies on X . But $\mathcal{T}_1 \cup \mathcal{T}_2 = \{\emptyset, X, \{a\}, \{b\}\}$ is not a topology since $\{a\}, \{b\} \in \mathcal{T}_1 \cup \mathcal{T}_2$ but $\{a\} \cup \{b\} = \{a, b\} \notin \mathcal{T}_1 \cup \mathcal{T}_2$. Hence a union of topologies is, in general, not a topology.

- Let $\mathcal{S} = \bigcup_\alpha \mathcal{T}_\alpha$. Then $X \in \mathcal{S}$ since X is in each individual \mathcal{T}_α as they are all topologies. It follows that $X = \bigcup_{S \in \mathcal{S}} S$ and so \mathcal{S} is a sub-basis. Let \mathcal{B} be the basis generated by \mathcal{S} and \mathcal{T}_s be the topology generated by \mathcal{B} . Fix some \mathcal{T}_α and take $U \in \mathcal{T}_\alpha$. Then $U \in \mathcal{S} \subset \mathcal{B} \subset \mathcal{T}_s$ by construction. Hence

$U \in \mathcal{T}_S$ and it follows that $\mathcal{T}_\alpha \subset \mathcal{T}_S$ for all α . Is it the smallest topology with such property? Let \mathcal{T}' be a topology on X such that $\mathcal{T}_\alpha \subset \mathcal{T}'$ for all α and take $U \in \mathcal{T}_S$. Then U is an arbitrary union of finite intersections of elements of $\mathcal{S} = \bigcup \mathcal{T}_\alpha \subset \mathcal{T}'$. Since \mathcal{T}' is a topology it is closed under arbitrary unions and finite intersections and so $U \in \mathcal{T}'$. Hence $\mathcal{T}_S \subset \mathcal{T}'$ and it follows that \mathcal{T}_S is the smallest topology containing all \mathcal{T}_α .

From part one we know that $\bigcap \mathcal{T}_\alpha$ is a topology, and by definition it is contained in \mathcal{T}_α for all α . If $\mathcal{T}' \subset \mathcal{T}_\alpha, \forall \alpha$ is a topology, then for every $U \in \mathcal{T}'$, $U \in \bigcap \mathcal{T}_\alpha$ and so $\mathcal{T}' \subset \bigcap \mathcal{T}_\alpha$. Therefore $\bigcap \mathcal{T}_\alpha$ is the largest topology that is contained in all \mathcal{T}_α .

c. Apply part (2).

Exercise 13.5

Show that that topology \mathcal{T} on X generated by a basis \mathcal{B} is equal to the intersections of all the topologies on X that contain \mathcal{B} .

Proof. Let $T = \{\mathcal{T}_\beta \mid \mathcal{B} \subset \mathcal{T}_\beta\}$ be the collection of all topologies on X that contain \mathcal{B} . Let $u \in \mathcal{T}$. Then U can be written as a union of element in \mathcal{B} , i.e.

$$U = \bigcup_{\alpha} B_{\alpha}, \quad B_{\alpha} \in \mathcal{B}$$

Since \mathcal{T}_β is a topology and $\mathcal{B} \subset \mathcal{T}_\beta$ for all $\mathcal{T}_\beta \in T$ it follows that $U = \bigcup_{\alpha} B_{\alpha} \in \mathcal{T}_\beta$ for all $\mathcal{T}_\beta \in T$ and so

$$\mathcal{T} \subset \bigcap_{\mathcal{T}_\beta \in T} \mathcal{T}_\beta.$$

Since $\mathcal{B} \subset \mathcal{T}$ by definition of a basis it follows that $\mathcal{T} \in T$ and so

$$\bigcap_{\mathcal{T}_\beta \in T} \mathcal{T}_\beta \subset \mathcal{T}.$$

Hence $\bigcap_{\mathcal{T}_\beta \in T} \mathcal{T}_\beta = \mathcal{T}$. □

Exercise 13.7

Solution.

Exercise 13.8

a. Show that the collection

$$\mathcal{B} = \{(a, b) \mid a < b, a, b \in \mathbb{Q}\}$$

generates the standard topology on \mathbb{R} .

b. Show that the collection

$$\mathcal{C} = \{[a, b) \mid a < b, a, b \in \mathbb{Q}\}$$

generates a topology different from the lower limit topology.

Solution.

- a. Let \mathcal{T} be the standard topology on \mathbb{R} and $\mathcal{T}_{\mathcal{B}}$ the topology generated by \mathcal{B} . Let $x \in (a, b) \in \mathcal{T}$. Since the rationals are dense in \mathbb{R} , there exist $a', b' \in \mathbb{Q}$ such that $x \in (a', b') \subset (a, b)$. Hence $\mathcal{T} \subset \mathcal{T}_{\mathcal{B}}$. The other inclusion is trivial since every basis element $(a, b) \in \mathcal{B}$ is a basis element of \mathcal{T} . We conclude that $\mathcal{T} = \mathcal{T}_{\mathcal{B}}$.
- b. Let \mathcal{T}_c be the topology generated by \mathcal{C} and let \mathcal{B} be the basis of \mathcal{T}_l , the lower limit topology on \mathbb{R} . Then for any $[a, b) \in \mathcal{C}$, $[a, b) \in \mathcal{B}$, and so $\mathcal{T}_c \subset \mathcal{T}_l$. To show that this inclusion is strict we need to prove the statement

$$\begin{aligned} & \neg (\forall B \in \mathcal{B} \forall x \in B \exists C \in \mathcal{C} : x \in C \subset B) \\ \iff & \exists B \in \mathcal{B} \exists x \in B \forall C \in \mathcal{C} : x \notin C \vee C \not\subset B \\ \iff & \exists B \in \mathcal{B} \exists x \in B \forall C \in \mathcal{C} : x \in C \implies C \not\subset B \end{aligned}$$

Let $[x, b) \in \mathcal{B}$ with $x \notin \mathbb{Q}$. Then $[a, c) \in \mathcal{C}$ can contain x only if $a < x$ since x is irrational. Therefore there is no element in \mathcal{C} that contains x and is a subset of $[x, b)$. This proves that $\mathcal{T}_c \subsetneq \mathcal{T}_l$.

The Subspace Topology

Exercise 16.1

Show that if Y is a subspace of X , and A is a subset of Y then the topology A inherits as a subspace of Y is the same as the topology it inherits as a subspace of X .

Proof. Let \mathcal{T} be a topology on X , \mathcal{T}_Y subspace topology on Y . Let \mathcal{T}'_A be the topology A inherits as a subset of Y . Then

$$\begin{aligned} \mathcal{T}'_A &= \{A \cap U \mid U \in \mathcal{T}_Y\} \\ &= \{A \cap U \mid U \in \{Y \cap V \mid V \in \mathcal{T}\}\} \\ &= \{A \cap U \mid U = Y \cap V, V \in \mathcal{T}\} \\ &= \{A \cap (Y \cap V) \mid V \in \mathcal{T}\} \\ &= \{(A \cap Y) \cap V \mid V \in \mathcal{T}\} \\ &= \{A \cap V \mid V \in \mathcal{T}\} \end{aligned}$$

which is by definition the topology A inherits as a subset of X . \square

Exercise 16.4

Show that $\pi_1 : X \times Y \rightarrow X$ is an open map.

Proof. Let U be open in $X \times Y$ and take $(x, y) \in U$. Then there exists a basis element $B_x \times B_y$ such that $(x, y) \in B_x \times B_y \subset U$. For any $b \in B_x$, $(b, y) \in B_x \times B_y \subset U$ and so $b = \pi_1(b, y) \in \pi_1(U)$. It follows that $B_x \subset \pi_1(U)$. Since the basis of a product topology is the the product of open sets, B_x is open in X which means that for every $x \in \pi_1(U)$ there is an open set $B_x \subset X$ such that $x \in B_x \subset \pi_1(U)$. From Exercise 13.1 it follows that $\pi_1(U)$ is open in X . \square

Closed Sets and Limit Points**Exercise 17.1**

Let \mathcal{C} be a collection of subsets of X . Suppose that $X, \emptyset \in \mathcal{C}$ and that \mathcal{C} is closed under finite unions and arbitrary intersections. Prove that

$$\mathcal{T} = \{X - C \mid C \in \mathcal{C}\}$$

is a topology.

Proof. Since $X - X = \emptyset$ and $X - \emptyset = X$ it follows that $X, \emptyset \in \mathcal{T}$. Let U_α be some collection of elements of \mathcal{T} . Then $U_\alpha = X - C_\alpha$ for some $C_\alpha \in \mathcal{C}$. Then

$$\bigcup_{\alpha} U_{\alpha} = \bigcup_{\alpha} (X - C_{\alpha}) = X - \bigcap_{\alpha} C_{\alpha} \in \mathcal{T}$$

Since \mathcal{C} is closed under arbitrary intersections. Lastly let $U_i = X - C_i$, $1 \leq i \leq n$ be a finite collection in \mathcal{T} . Then

$$\bigcap_{i=1}^n U_i = X - \bigcup_{i=1}^n C_i \in \mathcal{T}.$$

It follows that \mathcal{T} is a topology on X . \square

Exercise 17.2

Show that if A is closed in Y and Y is closed in X , then A is closed in X .

Proof. Suppose that A is closed in Y and Y is closed in X . Then $Y - A$ is open in Y and so $Y - A = U \cap Y$ for some $U \subset X$ open. Hence $A = (X - U) \cap Y$. Since $X - U$ and Y are closed in A and arbitrary intersections of closed sets are closed, it follows that A is closed in X . \square

Exercise 17.6

Let A, B and A_α denote subsets of a space X . Prove the following

- a. If $A \subset B$, then $\bar{A} \subset \bar{B}$

- b. $\overline{A \cup B} = \bar{A} \cup \bar{B}$
- c. $\overline{\bigcup_{\alpha} A_{\alpha}} \supset \bigcup_{\alpha} \bar{A}_{\alpha}$

Proof.

- a. Suppose that $A \subset B$. Let $x \in \bar{A}$. Then for every neighborhood U containing x intersects A . Hence $U \cap A$ is non empty, so take $y \in U \cap A$. Then $y \in U \cap B$ since $A \subset B$ and it follows that every neighborhood of x intersects B . Therefore $x \in \bar{B}$ and the result follows.
- b. Let $x \in \overline{A \cup B}$. Then every neighborhood U of x intersects $A \cup B$. Since $A \subset \bar{A}$ and $B \subset \bar{B}$ it follows that $A \cup B \subset \bar{A} \cup \bar{B}$ and so U intersects $\bar{A} \cup \bar{B}$. Hence $x \in \bar{A} \cup \bar{B}$ and so $\overline{A \cup B} \subset \bar{A} \cup \bar{B}$.
Let $x \in \bar{A} \cup \bar{B}$. Then x is in \bar{A} or \bar{B} and every neighborhood U of x intersects either A or B . Therefore U intersects $A \cup B$ and it follows that $x \in \overline{A \cup B}$. Therefore $\bar{A} \cup \bar{B} \subset \overline{A \cup B}$ and equality follows.
- c. Let $x \in \bigcup_{\alpha} \bar{A}_{\alpha}$. Then $x \in \bar{A}_{\alpha}$ for at least one α . Hence every neighborhood U of x intersects A_{α} and so U intersects $\bigcup_{\alpha} A_{\alpha}$. Hence $x \in \overline{\bigcup_{\alpha} A_{\alpha}}$ and so $\bigcup_{\alpha} \bar{A}_{\alpha} \subset \overline{\bigcup_{\alpha} A_{\alpha}}$.

To show that the other inclusion doesn't hold in general let $A_{\alpha} = (\frac{1}{\alpha}, 1)$. Then

$$\bigcup_{\alpha=1}^{\infty} \bar{A}_{\alpha} = \bigcup_{\alpha=1}^{\infty} \overline{\left(\frac{1}{\alpha}, 1\right)} = \bigcup_{\alpha=1}^{\infty} \left[\frac{1}{\alpha}, 1\right] = (0, 1]$$

and

$$\overline{\bigcup_{\alpha=1}^{\infty} A_{\alpha}} = \overline{(0, 1)} = [0, 1].$$

□

Exercise 17.11

Show that the product of two Hausdorff spaces is Hausdorff

Proof. Let X, Y be Hausdorff spaces and take $(x_1, y_1), (x_2, y_2) \in X \times Y$. Then there are open subsets U_1, U_2 in X such that $x_1 \in U_1, x_2 \in U_2$ and $U_1 \cap U_2 = \emptyset$. Similarly there exists $V_1, V_2 \subset Y$ that are open in Y with $y_1 \in V_1, y_2 \in V_2$ and $V_1 \cap V_2 = \emptyset$. Then $(x_1, y_1) \in U_1 \times V_1, (x_2, y_2) \in U_2 \times V_2$ and

$$(U_1 \times V_1) \cap (U_2 \times V_2) = (U_1 \cap U_2) \times (V_1 \cap V_2) = \emptyset \times \emptyset = \emptyset.$$

Since $U_1 \times V_1$ and $U_2 \times V_2$ are open it follows that $X \times Y$ is a Hausdorff space. □

Exercise 17.12

Show that a subspace of Hausdorff space is Hausdorff.

Proof. Let X be a Hausdorff space and $Y \subset X$. Then for every $x_1, x_2 \in Y$ there exists neighborhoods U_1 and U_2 of x_1 and x_2 (respectively) that are disjoint. Then $U_1 \cap Y$ and $U_2 \cap Y$ are open in Y , contain x_1 and x_2 (respectively) and are clearly disjoint. Hence Y is Hausdorff. \square

Exercise 17.13

Show that X is Hausdorff if and only if $\Delta = \{(x, x) \mid x \in X\}$ is closed in $X \times X$.

Proof. Suppose X is Hausdorff and consider $(x, y) \notin \Delta$. Since X is Hausdorff, there exists disjoint neighborhoods U and V that contain x and y (respectively). Then take $(x', y') \in U \times V$. Since $U \cap V = \emptyset$ it follows that $x' \neq y'$ and so $(x', y') \notin \Delta$. Since $U \times V$ is open in $X \times X$ it follows that (x, y) is not a limit point of Δ . Hence Δ contains all of its limit points and so it is closed.

Now suppose that Δ is closed in $X \times X$ and consider $(x, y) \notin \Delta$. Since Δ is closed, it contains all of its limit points and so (x, y) is not a limit point of Δ . Hence there exists a neighborhood T of (x, y) that does not intersect Δ . Then there is a basis element $U \times V$ such that $(x, y) \in U \times V \subset T$. Since $U \times V$ does not intersect Δ it follows that for every $(x', y') \in U \times V$ $x' \neq y'$. Hence $U \cap V = \emptyset$. It follows that for every x and y in X there exists disjoint neighborhoods U and V that contain x and y respectively. Therefore X is Hausdorff. \square

Exercise 17.14

In the finite complement topology on \mathbb{R} , to what point or points does the sequence $x_n = \frac{1}{n}$ converge?

Solution. This sequence converges to every point in \mathbb{R} ! To see why, suppose there exists $a \in \mathbb{R}$ that the sequence does not converge to. Then there exists an open neighborhood U of a such that for all $N \in \mathbb{N}$ there exists an $n \geq N$ for which $x_n \notin U$. There has to be an infinite number of such x_n , for otherwise we could just take N' bigger than the largest n . But then $\mathbb{R} - U$ is not finite nor empty, so it must be all of \mathbb{R} . Therefore $U = \emptyset$ which is a contradiction since we assumed that $a \in U$.

Exercise 17.19

For $A \subset X$, the *boundary* of A is

$$\text{Bd } A = \overline{A} \cap \overline{(X - A)}.$$

Show that

- $\text{Int } A \cap \text{Bd } A = \emptyset$ and $\overline{A} = \text{Int } A \cup \text{Bd } A$.
- $\text{Bd } A = \emptyset \iff A$ is both open and closed.
- U is open $\iff \text{Bd } U = \overline{U} - U$
- If U is open, is it true that $U = \text{Int } \overline{U}$? Justify your answer.

Proof.

- a. Let $x \in \text{Int } A$. Then there is a neighborhood U of x that is contained in A , and so U does not intersect $X - A$. Hence x is not a limit point of $X - A$ and so $x \notin \overline{X - A}$. Therefore x is not in $\overline{A} \cap \overline{(X - A)}$, and so $\text{Int } A \cap \text{Bd } A = \emptyset$.

Let $x \in X$. If $x \in \text{Int } A$ then clearly $x \in \text{Int } A \cup \overline{(X - A)}$. So suppose $x \notin \text{Int } A$. Then $x \in X - \text{Int } A \subset \overline{X - \text{Int } A}$. Consider any neighborhood U containing x . Since x is not in the interior of A , U is not a subset of A , hence U intersects $X - A$. It follows that x is a limit point of $X - A$ and so $x \in \overline{X - A}$. Therefore $X = \text{Int } A \cup \overline{(X - A)}$ and it follows that

$$\begin{aligned} \text{Int } A \cup \text{Bd } A &= \text{Int } A \cup \left(\overline{A} \cap \overline{(X - A)} \right) \\ &= (\text{Int } A \cup \overline{A}) \cap \left(\text{Int } A \cup \overline{(X - A)} \right) \\ &= \overline{A} \cap \left(\text{Int } A \cup \overline{(X - A)} \right) \\ &= \overline{A} \cap X \\ &= \overline{A}. \end{aligned}$$

- b. Suppose $\text{Bd } A = \emptyset$. Then

$$\overline{A} = \text{Int } A \cup \text{Bd } A = \text{Int } A.$$

Since $\text{Int } A \subset A \subset \overline{A}$ it follows that $A = \text{Int } A = \overline{A}$ and so A is both closed and open.

Now suppose that A is both closed and open. Then $A = \text{Int } A$ and $A = \overline{A}$ so $\text{Int } A = \overline{A}$. From $\overline{A} = \text{Int } A \cup \text{Bd } A$ it follows that $\text{Bd } A \subset \overline{A}$ but

$$\emptyset = \text{Int } A \cap \text{Bd } A = \overline{A} \cap \text{Bd } A.$$

Hence $\text{Bd } A = \emptyset$.

- c. Suppose U is open. Then $U = \text{Int } U$ and $\overline{U} = U \cup \text{Bd } U$. Since $U \cap \text{Bd } U = \emptyset$ it follows that $\text{Bd } U = \overline{U} - U$.

Now suppose that $\text{Bd } U = \overline{U} - U$ and take $x \in U$. Then $x \in \overline{U}$ and

$$x \notin \overline{U} - U = \text{Bd } U = \overline{U} \cap \overline{(X - U)}.$$

So $x \notin \overline{(X - U)}$. Then there exists a neighborhood V of x that does not intersect $X - U$. Since for every $y \in V \implies y \notin X - U \implies y \in U$ it follows that $V \subset U$. We showed that for every $x \in U$ there exists an open neighborhood $V \subset U$ that contains x . Therefore U is open.

- d. No. Consider $U = (0, 1) \cup (1, 2)$ which is open in the standard topology on \mathbb{R} . Then $\overline{U} = [0, 2]$ and $\text{Int } \overline{U} = (0, 2) \neq U$.

□

Continuous Functions

Exercise 18.3

Let X and X' denote a single set in two topologies, \mathcal{T} and \mathcal{T}' respectively. Let $i : X' \rightarrow X$ be the identity map.

- Show that i is continuous $\iff \mathcal{T} \subset \mathcal{T}'$.
- Show that i is a homeomorphism $\iff \mathcal{T} = \mathcal{T}'$.

Proof.

- Suppose i is continuous. Let B be a basis element in X and take $x \in X$. Then $B = i^{-1}(B)$ is open in X' . So there exists a basis element $B' \subset X$ such that $x \in B' \subset B$. It follows that \mathcal{T}' is finer than \mathcal{T} .

Now suppose that $\mathcal{T} \subset \mathcal{T}'$. Then for any $U \in \mathcal{T}$, $U \in \mathcal{T}'$. This is equivalent to saying that for any open set U in X , $i^{-1}(U) = U$ is open in X' . Therefore i is continuous.

- i is a homeomorphism if and only if i and i^{-1} are continuous. By the result above, i is continuous if and only if $\mathcal{T} \subset \mathcal{T}'$. Similarly, i^{-1} is continuous if and only if $\mathcal{T}' \subset \mathcal{T}$. Therefore i is a homeomorphism if and only if $\mathcal{T}' = \mathcal{T}$.

□

Exercise 18.10

Let $f : A \rightarrow B$ and $g : C \rightarrow D$ be continuous functions. Let us define a map $f \times g : A \times C \rightarrow B \times D$ by

$$(f \times g)(a \times c) = f(a) \times g(c).$$

Show that $f \times g$ is continuous.

Proof. Let $U \times V \subset B \times D$ be a basis element and suppose $a \times c \in (f \times g)^{-1}(U \times V)$. Then $a \in f^{-1}(U)$ and $c \in g^{-1}(V)$ and so $a \times c \in f^{-1}(U) \times g^{-1}(V)$. Now suppose that $a \times c \in f^{-1}(U) \times g^{-1}(V)$. Then $f(a) \in U$ and $g(c) \in V$ and so

$$\begin{aligned} (f \times g)(a \times c) &= f(a) \times g(c) \\ &\in U \times V. \end{aligned}$$

Therefore $a \times c \in (f \times g)^{-1}(U \times V)$ and it follows that $(f \times g)^{-1}(U \times V) = f^{-1}(U) \times g^{-1}(V)$. Since $f^{-1}(U)$ is open in A and $g^{-1}(V)$ is open in C it follows that $f^{-1}(U) \times g^{-1}(V)$ is open in $A \times C$ and so $f \times g$ is continuous. □

Exercise 18.13

Let Y be Hausdorff, $A \subset X$ and $f : A \rightarrow Y$ continuous. Show that if f may be extended to a continuous function $g : \overline{A} \rightarrow Y$, then g is uniquely determined by f .

Proof. Let g and \tilde{g} be continuous maps from \overline{A} to Y s.t. $g(a) = f(a) = \tilde{g}(a)$ for all $a \in A$. Suppose there exists $x \in \overline{A}$ such that $g(x) \neq \tilde{g}(x)$. Then there exists disjoint open neighborhoods U and V in Y such that $g(x) \in U$ and $\tilde{g}(x) \in V$. Then $U' = g^{-1}(U)$ and $V' = \tilde{g}^{-1}(V)$ are open in \overline{A} and contain x . Thus $U' \cap V'$ is open in \overline{A} and contains x , so it intersects A . Take $y \in A$ such that $y \in U' \cap V'$. Then $f(y) = g(y) = \tilde{g}(y)$, $g(y) \in U$ and $\tilde{g}(y) \in V$. But then $f(y) \in U \cap V$ and so U and V cannot be disjoint. By contradiction, $g(x) = \tilde{g}(x)$ for all $x \in \overline{A}$. \square

Extra Exercises

Exercise 2.18.1

Let X be a Hausdorff space and Y a non Hausdorff space.

- Can there be a continuous bijective map $X \rightarrow Y$? Give an example or prove that this is not possible
- Can there be a continuous bijective map $Y \rightarrow X$? Give an example or prove that this is not possible
- Show that X and Y are not homeomorphic.

Proof.

- Yes. Suppose $X = Y$, $\mathcal{T}_X = \mathcal{P}(X)$, $\mathcal{T}_Y = \{\emptyset, X\}$ and $f : X \rightarrow Y$ is the identity map. Then X is Hausdorff, Y is not and f is clearly bijective and continuous.
- Suppose it is possible. Then there exists a continuous bijective map $f : Y \rightarrow X$. Let $x, y \in Y$ such that there are no disjoint open neighborhoods containing x and y . Then $f(x) \neq f(y)$ since f is injective and so there exists open sets U and V in X such that $f(x) \in U$, $f(y) \in V$ and $U \cap V = \emptyset$. Then $x \in f^{-1}(U)$ and $y \in f^{-1}(V)$ and $f^{-1}(U) \cap f^{-1}(V)$ is nonempty. Let $z \in f^{-1}(U) \cap f^{-1}(V)$. Then $f(z) \in U$ and $f(z) \in V$ which is a contradiction. Therefore f does not exist :)
- Follows immediately from (1), (2) and the definition of Homeomorphism.

\square

Exercise 2.18.2

Give an explicit homeomorphism $f : \mathbb{R} \rightarrow (0, \infty)$.

Solution. Let $f(x) = e^x$. Then f is injective since $e^x = e^y \implies x = y$ and surjective since $\forall x \in (0, \infty)$, $e^{\ln x} = x$. So f is bijective. Moreover, $f((a, b)) = (e^a, e^b)$ and $f^{-1}((a, b)) = (\ln a, \ln b)$ and so f is a homeomorphism.

The Product Topology

Exercise 19.3

If each space X_α is Hausdorff, then $X = \prod_\alpha X_\alpha$ is Hausdorff in both the box and product topology.

Proof. Let $\mathbf{x}, \mathbf{y} \in X$ such that $\mathbf{x} \neq \mathbf{y}$. Then $x_\beta \neq y_\beta$ for at least one $\beta \in J$. Then there exists disjoint open neighborhoods W_x and W_y around x_β and y_β in X_β . Let

$$U_\alpha = \begin{cases} W_x, & \alpha = \beta \\ X_\alpha, & \alpha \neq \beta \end{cases} \quad \text{and} \quad V_\alpha = \begin{cases} W_y, & \alpha = \beta \\ X_\alpha, & \alpha \neq \beta \end{cases}.$$

Then $\prod_\alpha U_\alpha$ and $\prod_\alpha V_\alpha$ are disjoint and open in both the product and box topologies (open in the box topology since they are a product of open sets and in the product topology since they have finitely many components that are not all of X_α). To prove that they are disjoint, suppose there exists $\mathbf{z} \in (\prod_\alpha U_\alpha) \cap (\prod_\alpha V_\alpha)$. Then $z_\beta \in U_\beta \cap V_\beta = W_x \cap W_y = \emptyset$. It follows that X is Hausdorff. \square

Exercise 19.7

Let \mathbb{R}^∞ be a subset of \mathbb{R}^ω consisting of all sequences that are "eventually zero", that is, all sequences (x_1, x_2, \dots) such that $x_i = 0$ for all $i \geq n$ for some $n \in \mathbb{N}$. What is the closure of \mathbb{R}^∞ in \mathbb{R}^ω in the box and product topologies?

Solution. We start with the box topology. Let $\mathbf{x} \in \mathbb{R}^\omega \setminus \mathbb{R}^\infty$. Then let J be an index set such that $x_j \neq 0$ for all $j \in J$. Note that J is infinite since $\mathbf{x} \notin \mathbb{R}^\infty$. Then there exists an open neighborhood U_j around x_j that does not contain 0. Let $U = \prod_{i \in \mathbb{N}} V_i$ where

$$V_i = \begin{cases} U_i, & i \in J \\ A_i \text{ open in } \mathbb{R} \text{ and } x_i \in A_i, & i \notin J \end{cases}.$$

Since for any $(y_1, y_2, \dots) \in U$, $y_i \neq 0$ for infinitely many i , it follows that $(y_1, y_2, \dots) \notin \mathbb{R}^\infty$ and so $U \cap \mathbb{R}^\infty = \emptyset$. Therefore for any $\mathbf{x} \in \mathbb{R}^\omega \setminus \mathbb{R}^\infty$ there exists a neighborhood U containing \mathbf{x} such that $U \subset \mathbb{R}^\omega \setminus \mathbb{R}^\infty$ and it follows that $\mathbb{R}^\omega \setminus \mathbb{R}^\infty$ is open and so \mathbb{R}^∞ is closed. Therefore $\overline{\mathbb{R}^\infty} = \mathbb{R}^\infty$ in the box topology.

We now consider the product topology. We will show that $\text{Int } \mathbb{R}^\omega \setminus \mathbb{R}^\infty = \emptyset$ and so \mathbb{R}^∞ is dense in \mathbb{R}^ω . Let U be a basis element in \mathbb{R}^ω and J an index set such that $U_j \subsetneq \mathbb{R}$ for all $j \in J$. Then J is finite by definition of the product topology. So there exists a sequence $\mathbf{y} = (y_1, y_2, \dots)$ in \mathbb{R}^∞ such that $y_i \in U_i$ for $i \in J$ and $y_i = 0$ otherwise. Then $\mathbf{y} \in U$. Therefore there is no open set contained in $\mathbb{R}^\omega \setminus \mathbb{R}^\infty$ and it follows that $\text{Int } \mathbb{R}^\omega \setminus \mathbb{R}^\infty = \emptyset$. Hence \mathbb{R}^∞ is dense in \mathbb{R}^ω and so $\overline{\mathbb{R}^\infty} = \mathbb{R}^\omega$.

Exercise 19.10

Let A be a set; let $\{X_\alpha\}_{\alpha \in J}$ be an indexed family of spaces; and let $\{f_\alpha\}$ be an indexed family of functions $f_\alpha : A \rightarrow X_\alpha$.

- a. Show that there is a unique coarsest topology \mathcal{T} on A relative to which each of the functions f_α is continuous.

- b. Let

$$\mathcal{S}_\beta = \left\{ f_\beta^{-1}(U_\beta) \mid U_\beta \text{ is open in } X_\beta \right\}$$

and let $\mathcal{S} = \bigcup_\alpha \mathcal{S}_\alpha$. Show that \mathcal{S} is a subbasis for \mathcal{T} .

- c. Show that a map $g : Y \rightarrow A$ is continuous relative to \mathcal{T} if and only if each map $f_\alpha \circ g$ is continuous.

- d. Let $f : A \rightarrow \prod X_\alpha$ be defined by

$$f(a) = (f_\alpha(a))_{\alpha \in J}.$$

Let Z denote the subspace $f(A)$ of the product space $\prod X_\alpha$. Show that the image under f of each element of \mathcal{T} is an open set of Z .

Solution.

- a. Let \mathcal{C} be the collection of all topologies on A such that each f_α is continuous. This collection is not empty since every function is continuous relative to $\mathcal{P}(A)$. Let $\mathcal{T} = \bigcap_{\mathcal{T}' \in \mathcal{C}} \mathcal{T}'$. By Exercise 13.4 we know that \mathcal{T} is the unique coarsest topology in the collection. Since $\mathcal{T} \in \mathcal{C}$ each f_α is continuous relative to \mathcal{T} .

- b. We show that \mathcal{S} is a subbasis by proving that $\mathcal{T}' \in \mathcal{C}$ if and only if $\mathcal{S} \subset \mathcal{T}'$. Since \mathcal{T} is the coarsest topology in \mathcal{C} , it is the smallest topology containing \mathcal{S} and so \mathcal{S} is a basis.

Suppose $\mathcal{T}' \in \mathcal{C}$ and let $U \in \mathcal{S}$. Then $U = f_\beta^{-1}(U_\beta)$ for some U_β open in X_β . Since $\mathcal{T}' \in \mathcal{C}$ it follows that U is open in \mathcal{T}' and so $\mathcal{S} \subset \mathcal{T}'$.

Conversely suppose that \mathcal{T}' is a topology on A such that $\mathcal{S} \subset \mathcal{T}'$. Then for any U_β open in X_β , $f_\beta^{-1}(U_\beta) \in \mathcal{S}$ and so it is open in \mathcal{T}' . Hence every f_α is continuous relative to \mathcal{T}' and so $\mathcal{T}' \in \mathcal{C}$.

- c. Suppose $g : Y \rightarrow A$ is continuous and let U_β be an open subset of X_β . Then $U = f_\beta^{-1}(U_\beta)$ is open in A and so $g^{-1}(U)$ is open in Y . Since β was arbitrary, it follows that $f_\alpha \circ g : Y \rightarrow X_\alpha$ is continuous for all α .

Conversely suppose that $f_\alpha \circ g : Y \rightarrow X_\alpha$ is continuous for all α and let U be open in A . By part (b) we know that U is an arbitrary union of finite intersections of elements $f_\beta^{-1}(U_\beta) \in \mathcal{S}$. There $g^{-1}(U)$ is an arbitrary union of finite intersections of elements of the form $g^{-1}\left(f_\beta^{-1}(U_\beta)\right) = (f_\beta \circ g)^{-1}(U_\beta)$ which are all open in Y . Hence $g^{-1}(U)$ is open in Y and so g is continuous.

- d. Let U be an open subset of A and consider $f(U) = (f_\alpha(U))_{\alpha \in J}$. Let $\mathbf{y} \in f(U)$ such that $\mathbf{y} = f(a)$ for some $a \in A$. Then there exists a basis element B_A such that $a \in B_A$. By part (b) B_A is a finite intersection of preimages of open sets. Hence

$$B_A = \prod_{\alpha \in I} f_\alpha^{-1}(U_\alpha)$$

for some finite $I \subset J$. Then the element

$$V = \prod_{\alpha \in I} U_\alpha \times \prod_{\alpha \in J \setminus I} X_\alpha$$

is open in $\prod X_\alpha$ since I is finite. Then $B_Z = V \cap Z$ is a basis element of Z . Furthermore, since $\mathbf{y} = f(a) \in f(A) = Z$ and $y_\alpha = f_\alpha(a) \in U_\alpha$ for all $\alpha \in I$ and $y_\alpha \in X_\alpha$ for all $\alpha \in J \setminus I$ it follows that $\mathbf{y} \in B_Z$. Since \mathbf{y} was arbitrary, $f(U)$ is an arbitrary union of basis elements in Z and so it is open in Z .

Metric Spaces

Exercise 20.1a

In \mathbb{R}^n let

$$d'(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^n |x_i - y_i|.$$

Show that d' is a metric that induces the usual topology on \mathbb{R}^n .

Proof. Clearly d' define a metric. Let $\varepsilon > 0$. We will show that

$$B_d\left(\mathbf{x}, \frac{\varepsilon}{\sqrt{n}}\right) \subset B_{d'}(\mathbf{x}, \varepsilon) \subset B_d(\mathbf{x}, \mathbf{y}) \quad (1)$$

from which the result would follow. Recall the inequality

$$\sqrt{a_1 + \cdots + a_n} \leq \sqrt{a_1} + \cdots + \sqrt{a_n} \leq \sqrt{n(a_1 + \cdots + a_n)}, \quad (2)$$

for $a_i \geq 1$. Let $\mathbf{y} \in B_{d'}(\mathbf{x}, \varepsilon)$. Then

$$\begin{aligned} d(\mathbf{x}, \mathbf{y}) &= \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}} \\ &\leq \sum_{i=1}^n \sqrt{(x_i - y_i)^2} && \text{(By (2))} \\ &= \sum_{i=1}^n |x_i - y_i| \\ &= d'(\mathbf{x}, \mathbf{y}) \\ &< \varepsilon \end{aligned}$$

which proves the right inequality of (1). Now let $\mathbf{y} \in B_d\left(\mathbf{x}, \frac{\varepsilon}{\sqrt{n}}\right)$. Then using the left right inequality of (2) we get

$$\begin{aligned} d'(\mathbf{x}, \mathbf{y}) &= \sum_{i=1}^n |x_i - y_i| \\ &= \sum_{i=1}^n \sqrt{(x_i - y_i)^2} \\ &\leq \sqrt{n} \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}} \\ &= \sqrt{n} d(\mathbf{x}, \mathbf{y}) \\ &< \sqrt{n} \frac{\varepsilon}{\sqrt{n}} = \varepsilon \end{aligned}$$

which proves the left inequality of (1) and completes the proof. \square

Exercise 20.3

Let X be a metric space with a metric d .

- Show that $d : X \times X \rightarrow \mathbb{R}$ is continuous.
- Let X' denote a space having the same underlying set as X . Show that if $d : X' \times X' \rightarrow \mathbb{R}$ is continuous, then the topology of X' is finer than the topology of X .

Proof.

- Let $\varepsilon > 0$ and $(x, y) \in X \times X$. Then for $\delta = \varepsilon/2$ the open set $U = B(x, \delta) \times B(y, \delta)$ is a neighborhood of (x, y) . Let $d = d(x, y)$ and consider $(d - \varepsilon, d + \varepsilon)$ in \mathbb{R} . Then for $(x', y') \in U$

$$\begin{aligned} d(x', y') &\leq d(x', x) + d(x, y) + d(y, y') \\ &< 2\delta + d \\ &< \varepsilon + d \end{aligned}$$

and

$$\begin{aligned} d(x, y) &\leq d(x', x) + d(x', y') + d(y, y') \\ &< d(x', y') + \varepsilon. \end{aligned}$$

Therefore $d(x', y') \in (d - \varepsilon, d + \varepsilon)$ for all $(x', y') \in U$ and so $d(U) \subset (d - \varepsilon, d + \varepsilon)$. Since for any open set V in \mathbb{R} containing $d(x, y)$ there exists a neighborhood U in $X \times X$ such that $d(U) \subset V$ it follows that d is continuous.

b. Suppose $d : X' \times X' \rightarrow \mathbb{R}$ is continuous. Let $\varepsilon > 0$ and consider

$$U = d^{-1}((-\infty, \varepsilon)) = \{(x, y) \mid d(x, y) < \varepsilon\}$$

which is open in $X' \times X'$ since d is continuous. Let $x \in X$ and $y \in B(x, \varepsilon)$. Since $d(x, y) < \varepsilon$ it follows that $(x, y) \in U$ and so there exists a basis $V_1 \times V_2$ element of $X' \times X'$ such

$$(x, y) \in V_1 \times V_2 \subset U.$$

Where V_2 is an open neighborhood in X' containing y . Since for any $z \in V_2$, $(x, z) \in U$ it follows that $d(x, z) < \varepsilon$ and so $z \in B(x, \varepsilon)$. Therefore $V_2 \subset B(x, \varepsilon)$. Since for all y there exists an open neighborhood $V \subset B(x, \varepsilon)$ in X' containing y and it follows that $X \subset X'$.

□

Exercise 20.11

Show that if d is a metric for X , then

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}$$

is a bounded metric that gives the topology of X .

Proof. We start by proving the d' is a metric. Let

$$f(x) = \frac{x}{1 + x}.$$

Then $d'(x, y) = (f \circ d)(x, y)$. Since $f : \mathbb{R}_+ \rightarrow [0, 1)$ is one-to-one, increasing and $f(x) = 0$ iff $x = 0$ the first two properties of a metric follow immediately. For the triangle inequality, let $a, b \geq 0$. Then

$$f(a + b) - f(b) = \frac{a}{(1 + a + b)(1 + b)} \leq \frac{a}{1 + a} = f(a)$$

and so

$$\begin{aligned} d'(x, y) &= f(d(x, y)) \\ &\leq f(d(x, z) + d(z, y)) \\ &\leq f(d(x, z)) + f(d(z, y)) \\ &\leq d'(x, z) + d'(z, y). \end{aligned}$$

Therefore d' is a metric.

Let X' be the space generated by the metric d' . Since $d : X \times X \rightarrow [0, \infty)$ and $f : [0, \infty) \rightarrow [0, \infty)$ are continuous and $d' = f \circ d$ it follows that $d' : X \times X \rightarrow [0, \infty)$ is continuous and so by Exercise 3, X is finer than X' .

To show that X' is finer than X , let $\varepsilon > 0$ and $\delta = \frac{\varepsilon}{1+\varepsilon}$. Then for $y \in B_{d'}(x, \delta)$, we have

$$\frac{d(x, y)}{1 + d(x, y)} < \delta$$

and so $d(x, y) < \frac{\delta}{1-\delta} = \varepsilon$. Hence $B_{d'}(x, \delta) \subset B_d(x, \varepsilon)$ and so X' is finer than X . It follows that d' generates the same topology as d . \square

The Quotient Topology

Exercise 22.2

- Let $p : X \rightarrow Y$ be a continuous map. Show that if there is a continuous map $f : Y \rightarrow X$ such that $p \circ f$ is the identity map on Y , then p is a quotient map.
- If $A \subset X$, a retraction of X onto A is a continuous map $r : X \rightarrow A$ such that $r(a) = a$ for all $a \in A$. Show that a retraction is a quotient map.

Proof.

- Let $y \in Y$. Then $(p \circ f)(y) = p(f(y)) = y$ and so p is surjective. Let U be an open subset of Y . Then $p^{-1}(U)$ is open in X since p is continuous. Now suppose that U is a subset of Y such that $p^{-1}(U)$ is open in X . Then $f^{-1}(p^{-1}(U)) = (p \circ f)^{-1}(U) = U$ since f is continuous. It follows that p is a quotient map.
- Let $i : A \rightarrow X$ be given by $i(a) = a$ for each $a \in A$. Let U be open in X . Then $i^{-1}(U) = U \cap A$ is open in A by definition of subspace topology. Therefore i is continuous. Since $r \circ i$ is the identity on A it follows by part (a) the r is a quotient map. \square

Exercise 22.3

Let $\pi_1 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be projection on the first coordinate. Let A be the a subspace of $\mathbb{R} \times \mathbb{R}$ consisting of all points (x, y) for which either $x \geq 0$ or $y = 0$. Let $q : A \rightarrow \mathbb{R}$ be obtained by restricting π_1 . Show that q is a quotient map that is neither closed not open.

Proof. Let $x \in \mathbb{R}$. Then $(x, 0) \in A$ and $q((x, 0)) = x$. Since for U open in \mathbb{R} , $\pi^{-1}(U)$ is open in $\mathbb{R} \times \mathbb{R}$ it follows that $q^{-1}(U) = \pi^{-1}(U) \cap A$ is open in A . Therefore q is continuous. Let $i : \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ be given by $x \mapsto (x, 0)$. Then for $U \times V$ open in $\mathbb{R} \times \mathbb{R}$, we have that

$$i^{-1}(U \times V) = \begin{cases} U, & \text{if } 0 \in V \\ \emptyset, & \text{otherwise} \end{cases}$$

is open in \mathbb{R} and so i is continuous. Since $i(\mathbb{R}) \subset A$ it follows that $i' : \mathbb{R} \rightarrow A$ is continuous. Since $(q \circ i')(x) = q((x, 0)) = x$ is the identity on \mathbb{R} it follows by Exercise 2 that q is a quotient map.

Consider the set

$$U = (-1, 1) \times (1, 2) \cap A = [0, 1) \times (1, 2)$$

which is open in A since $(-1, 1) \times (1, 2)$ is open in $\mathbb{R} \times \mathbb{R}$. Then $q(U) = [0, 1)$ which is not open in \mathbb{R} since any neighborhood containing 0 is not contained in $[0, 1)$. Next consider $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x, y) = xy$. Since f is continuous, $f^{-1}(\{1\})$ is closed and so

$$C = f^{-1}(\{1\}) \cap A = \{(x, 1/x) \mid x > 0\}$$

is closed in X . But $q(C) = (0, \infty)$ is not closed in \mathbb{R} . Therefore q is a quotient map that is neither open nor closed. \square

Exercise 22.4

- a. Define an equivalence relation on the plane $X = \mathbb{R}^2$ as follows:

$$x_0 \times y_0 \sim x_1 \times y_1 \quad \text{if} \quad x_0 + y_0^2 = x_1 + y_1^2$$

Let X^* be the corresponding quotient space. It is homeomorphic to a familiar space, what is it?

- b. Repeat (a) for the equivalence relation

$$x_0 \times y_0 \sim x_1 \times y_1 \quad \text{if} \quad x_0^2 + y_0^2 = x_1^2 + y_1^2.$$

Solution.

- a. Let

$$\begin{aligned} g : \mathbb{R}^2 &\rightarrow \mathbb{R} \\ (x, y) &\mapsto x + y^2. \end{aligned}$$

If g is continuous and surjective quotient map, then it would induce a homeomorphism from

$$X^* = \{g^{-1}(\{a\}) \mid a \in \mathbb{R}\} = \{(x, y) \mid x + y^2 = a, a \in \mathbb{R}\}$$

to \mathbb{R} . Let $(a, b) \subset \mathbb{R}$. Then

$$U = g^{-1}((a, b)) = \{(x, y) \mid x + y^2 \in (a, b)\}.$$

Then U is the region between the parabolas $x = a - y^2$ and $x = b - y^2$. Then for any point $(x, y) \in U$ we can find the minimum distance to each parabola and take an ε -neighborhood smaller than that. Therefore U is open and so g is continuous. Let $h : \mathbb{R} \rightarrow \mathbb{R}^2$ be given by $h(x) = (x, 0)$. Then h is continuous and $(g \circ h)(x) = g((x, 0)) = x$ is the identity on \mathbb{R} . Therefore g is a quotient map and so X^* is homeomorphic to \mathbb{R} .

b.

Exercise 22.5

Let $p : X \rightarrow Y$ be an open map. Show that if A is open in X , then the map $q : A \rightarrow p(A)$ obtained by restricting p is an open map.

Proof. Let U be open in A . Then U is open in X since A is open in X and so $p(U)$ is open in Y . Then $p(U) \cap p(A) = p(U)$ is open in $p(A)$ and so q is an open map. \square

Connected Spaces

Exercise 23.1

Let \mathcal{T} and \mathcal{T}' be two topologies on X . If $\mathcal{T} \subset \mathcal{T}'$, what does connectedness of X in one topology imply about connectedness in the other?

Solution. Suppose $X = U \cup V$ is a separation of X in \mathcal{T} . Then U, V are open in \mathcal{T}' and so \mathcal{T}' is not connected. Therefore

$$\mathcal{T}' \text{ connected} \implies \mathcal{T} \text{ connected.}$$

Exercise 23.3

Let $\{A_\alpha\}$ be a collection of connected subspaces of X . Let A be a connected subspace of X . Show that if $A \cap A_\alpha \neq \emptyset$ for all α , then $A \cup (\bigcup_\alpha A_\alpha)$ is connected.

Proof. Each subspace $A \cup A_\alpha$ is connected because A and A_α share a point. Then $\bigcap_\alpha (A \cup A_\alpha)$ is not empty and so $\bigcup_\alpha (A \cup A_\alpha) = A \cup (\bigcup_\alpha A_\alpha)$ is connected. \square

Exercise 23.4

Show that if X is an infinite set, it is connected in the finite complement topology.

Proof. Suppose $X = U \cup V$ is a separation of X in the finite complement topology. Then $U = X \setminus \{x_1, \dots, x_n\}$ and $V = X \setminus \{y_1, \dots, y_m\}$. But

$$\begin{aligned} U \cap V &= (X \setminus \{x_1, \dots, x_n\}) \cap (X \setminus \{y_1, \dots, y_m\}) \\ &= X \setminus (\{x_1, \dots, x_n\} \cup \{y_1, \dots, y_m\}) \end{aligned}$$

cannot be empty. Hence X is connected in the finite complement topology. \square

Exercise 23.9

Let A be a proper subset of X and B a proper subset of Y . If X and Y are connected, show that

$$(X \times Y) - (A \times B)$$

is connected.

Proof. For every $x \notin A$, $U_x = \bigcup_{y \in Y} x \times y$ is homeomorphic to Y and so it is connected. Similarly for every $y \notin B$, $V_y = \bigcup_{x \in X} x \times y$ is homeomorphic to X . Let $a \notin A$ and $b \notin B$. Then $U_a \cup V_b$ is connected since $U_a \cap V_b = (a, b)$ is nonempty. Note that for any $x \notin A$ and $y \notin B$,

$$(U_x \cup V_y) \cap (U_a \cup V_b) = \{(a, y), (x, b)\}.$$

and so $(U_x \cup V_y) \cup (U_a \cup V_b)$ is connected as well. Therefore by Exercise 23.3 it follows that

$$\bigcup_{(x,y) \notin A \times B} U_x \cup V_y = (X \times Y) - (A \times B)$$

is connected. □

Exercise 23.11

Let $p : X \rightarrow Y$ be a quotient map. Show that if Y is connected and each set $p^{-1}(\{y\})$ is connected then X is connected

Proof. Let $U \cup V$ be a separation of X and take $y \in Y$. Then $p^{-1}(\{y\})$ is connected and so it is contained in either U or V (but not both). Thus U and V are saturated with respect to p . Therefore $p(U)$ and $p(V)$ are disjoint open sets in Y and since p is surjective $Y = p(U) \cup p(V)$, a contradiction. □

Connected Subspaces of the Real Line

Exercise 24.1

- a. Is a product of path-connected spaces necessarily path connected?
- b. If $A \subset X$ and A is path connected, is \bar{A} necessarily path connected?
- c. If $f : X \rightarrow Y$ is continuous and X is path connected, is $f(X)$ necessarily path connected?
- d. If $\{A_\alpha\}$ is a collection of path connected subspaces of X and $\bigcap_\alpha A_\alpha \neq \emptyset$, is $\bigcup_\alpha A_\alpha$ necessarily path connected?

Solution.

- a. Let $\prod_\alpha X_\alpha$ be given in the product topology and let $\mathbf{x}, \mathbf{y} \in \prod_\alpha X_\alpha$. Then for each α there exists a continuous map $f_\alpha : [0, 1] \rightarrow X_\alpha$ such that $f_\alpha(0) = x_\alpha$ and $f_\alpha(1) = y_\alpha$. Then $f : [0, 1] \rightarrow \prod_\alpha X_\alpha$ mapping $f(t) = (f_\alpha(t))_\alpha$ is continuous and so $\prod_\alpha X_\alpha$ is path connected.
- b. No. The topologist's sine curve is a counter example.

- c. let $a, b \in f(X)$. Then there exists $x, y \in X$ such that $f(x) = a$ and $f(y) = b$. Moreover, there exists a continuous map $g : [0, 1] \rightarrow X$ such that $g(0) = x$ and $g(1) = y$. Then $f \circ g : [0, 1] \rightarrow f(X)$ is continuous, $(f \circ g)(0) = f(x) = a$ and $(f \circ g)(1) = f(y) = b$. Hence $f(X)$ is path connected.
- d. Let $p \in \bigcap_{\alpha} A_{\alpha}$ and $x, y \in \bigcup_{\alpha} A_{\alpha}$. If $x, y \in A_{\alpha}$ then there exists a path between them since A_{α} is path connected. So suppose $x \in A_{\alpha}$ and $y \in A_{\beta}$. Then there exists $f : [0, 1] \rightarrow A_{\alpha}$, $g : [0, 1] \rightarrow A_{\beta}$ such that $f(0) = x$, $f(1) = p$, $g(0) = p$ and $g(1) = y$. Then the map $h : [0, 2] \rightarrow A_{\alpha} \cup A_{\beta}$ given by

$$h(a) = \begin{cases} f(a), & \text{if } a \in [0, 1] \\ g(a - 1), & \text{if } a \in [1, 2] \end{cases}$$

is continuous by the pasting lemma. It follows that $\bigcup_{\alpha} A_{\alpha}$ is path connected.

Components and Local Connectedness

Exercise 25.1

What are the components and path components of \mathbb{R}_l ? What are the continuous maps $f : \mathbb{R} \rightarrow \mathbb{R}_l$?

Solution. Let $A \subset \mathbb{R}_l$ be a connected subspace that is not empty and take $a \in A$. Then $a \in A \cap [a, \infty)$ so $A \cap (-\infty, a) = \emptyset$ since A is connected. Choose any $b \in A$. Then $a \leq b$ but also $b \in A \cap [b, \infty)$ and $A \cap (-\infty, b) = \emptyset$ so $b \leq a$. It follows that $b = a$ and $A = \{a\}$. Therefore the only connected sets of \mathbb{R}_l are singletons. Since any path component is a component, it follows that path components are singletons as well.

Since the image of a connected space under a continuous map must be connected, the only continuous maps $f : \mathbb{R} \rightarrow \mathbb{R}_l$ are constant functions.