

Dual Descent

February 20, 2025

**Chapter 2 in Shai Shalev Shwartz / Online Learning and Online
convex Optimization**

Follow-the-Regularized-Leader (FTRL)

$$\forall t, \quad w_t = \arg \min_{w \in S} \sum_{i=1}^{t-1} f_i(w) + R(w)$$

- ▶ Regularizer controls length of weight vector - ζ changes from iteration to iteration.

Follow-the-Regularized-Leader (FTRL)

$$\forall t, \quad w_t = \arg \min_{w \in S} \sum_{i=1}^{t-1} f_i(w) + R(w)$$

- ▶ Regularizer controls length of weight vector - ζ changes from iteration to iteration.
- ▶ Each step requires solving a constrained minimization problem.

Review: Property of FoRel Algorithm

Lemma 2.3:

Let w_1, w_2, \dots be the sequence of vectors produced by the FoRel algorithm. Then, for all $u \in S$, we have:

$$\sum_{t=1}^T (f_t(w_t) - f_t(u)) \leq R(u) - R(w_1) + \sum_{t=1}^T (f_t(w_t) - f_t(w_{t+1}))$$

Review: One step of Gradient Descent using strongly convex regularizer

Lemma 2.10:

Let $R : S \rightarrow \mathbb{R}$ be a σ -strongly-convex function over S with respect to a norm $\|\cdot\|$. Let w_1, w_2, \dots be the predictions of the FoReL algorithm.

Then, for all t , if f_t is L_t -Lipschitz with respect to $\|\cdot\|$, we have:

$$f_t(w_t) - f_t(w_{t+1}) \leq L_t \|w_t - w_{t+1}\| \leq \frac{L_t^2}{\sigma}$$

Main Theorem regarding FoReL using strongly convex regularizer

Let f_1, \dots, f_T be a sequence of convex functions with the following conditions:

- ▶ f_t is L_t -Lipschitz with respect to some norm $\|\cdot\|$.

Then, for all $u \in S$,

$$\text{Regret}_T(u) \leq R(u) - \min_{v \in S} R(v) + \frac{TL^2}{\sigma}$$

Main Theorem regarding FoReL using strongly convex regularizer

Let f_1, \dots, f_T be a sequence of convex functions with the following conditions:

- ▶ f_t is L_t -Lipschitz with respect to some norm $\|\cdot\|$.
- ▶ L satisfies $\frac{1}{T} \sum_{t=1}^T L_t^2 \leq L^2$.

Then, for all $u \in S$,

$$\text{Regret}_T(u) \leq R(u) - \min_{v \in S} R(v) + \frac{TL^2}{\sigma}$$

Main Theorem regarding FoReL using strongly convex regularizer

Let f_1, \dots, f_T be a sequence of convex functions with the following conditions:

- ▶ f_t is L_t -Lipschitz with respect to some norm $\|\cdot\|$.
- ▶ L satisfies $\frac{1}{T} \sum_{t=1}^T L_t^2 \leq L^2$.
- ▶ FoReL is run on the sequence with a regularization function that is σ -strongly-convex with respect to the same norm.

Then, for all $u \in S$,

$$\text{Regret}_T(u) \leq R(u) - \min_{v \in S} R(v) + \frac{TL^2}{\sigma}$$

Potential based gradient Descent

- ▶ $\text{Regret}_t = \text{Regret vector}$ $\text{Regret}_t(w) = L_{A,t} - L_t(w)$

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies badness of the state.

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies **badness** of the state.
- ▶ A state is bad if adversary can force high regret in the future.

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies badness of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies **badness** of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t
- ▶ $w_t = \nabla R(\text{Regret}_t)$ is a good choice for the interior of S .

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies **badness** of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t
- ▶ $w_t = \nabla R(\text{Regret}_t)$ is a good choice for the interior of S .
- ▶ For finite number of experts, Regret_t is finite dimensional and we can compute w_t explicitly.

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies badness of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t
- ▶ $w_t = \nabla R(\text{Regret}_t)$ is a good choice for the interior of S .
- ▶ For finite number of experts, Regret_t is finite dimensional and we can compute w_t explicitly.
- ▶ Here, $\text{Regret} = \{R(w)\}_{w \in \mathbb{R}^d}$ is uncountably infinite.

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies badness of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t
- ▶ $w_t = \nabla R(\text{Regret}_t)$ is a good choice for the interior of S .
- ▶ For finite number of experts, Regret_t is finite dimensional and we can compute w_t explicitly.
- ▶ Here, $\text{Regret} = \{R(w)\}_{w \in \mathbb{R}^d}$ is uncountably infinite.
- ▶ If Experts correspond to exponential distributions and loss is log loss- we can use conjugate priors. (recall: biased coins).

Potential based gradient Descent

- ▶ Regret_t = Regret vector $\text{Regret}_t(w) = L_{A,t} - L_t(w)$
- ▶ Regret_t = State of prediction algorithm at time t
- ▶ Potential/Regularizer: $R(\text{Regret})$ Quantifies **badness** of the state.
- ▶ A state is bad if adversary can force high regret in the future.
- ▶ Choose prediction so that $R(\text{Regret}_{t+1}) - R(\text{Regret}_t) + w_t \cdot \ell_t$ is small for all ℓ_t
- ▶ $w_t = \nabla R(\text{Regret}_t)$ is a good choice for the interior of S .
- ▶ For finite number of experts, Regret_t is finite dimensional and we can compute w_t explicitly.
- ▶ Here, $\text{Regret} = \{R(w)\}_{w \in \mathbb{R}^d}$ is uncountably infinite.
- ▶ If Experts correspond to exponential distributions and loss is log loss- we can use conjugate priors. (recall: biased coins).
- ▶ We need a new trick to compute $w_t = \nabla R(\text{Regret}_t)$ efficiently.

FoReL Update Rule for linear cost function

Define $\mathbf{z}_{1:t} = \sum_{i=1}^t \mathbf{z}_i$, the FoReL update rule can be written as

$$\mathbf{w}_{t+1} = \arg \min_{\mathbf{w} \in S} R(\mathbf{w}) + \sum_{i=1}^t \langle \mathbf{w}, \mathbf{z}_i \rangle$$

$$= \arg \min_{\mathbf{w} \in S} R(\mathbf{w}) + \langle \mathbf{w}, \mathbf{z}_{1:t} \rangle$$

$$= \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, -\mathbf{z}_{1:t} \rangle - R(\mathbf{w}).$$

Mirror Descent Update for linear functions

Update rule

$$\mathbf{w}_{t+1} = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, -\mathbf{z}_{1:t} \rangle - R(\mathbf{w}).$$

Link Function:

$$g(\theta) = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, \theta \rangle - R(\mathbf{w}),$$

Update rule can be re-written as

1. $\theta_0 = 0$

Mirror Descent Update for linear functions

Update rule

$$\mathbf{w}_{t+1} = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, -\mathbf{z}_{1:t} \rangle - R(\mathbf{w}).$$

Link Function:

$$g(\theta) = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, \theta \rangle - R(\mathbf{w}),$$

Update rule can be re-written as

1. $\theta_0 = 0$
2. $\theta_{t+1} = \theta_t - \mathbf{z}_t$

Mirror Descent Update for linear functions

Update rule

$$\mathbf{w}_{t+1} = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, -\mathbf{z}_{1:t} \rangle - R(\mathbf{w}).$$

Link Function:

$$g(\theta) = \arg \max_{\mathbf{w} \in S} \langle \mathbf{w}, \theta \rangle - R(\mathbf{w}),$$

Update rule can be re-written as

1. $\theta_0 = 0$
2. $\theta_{t+1} = \theta_t - \mathbf{z}_t$
3. $\mathbf{w}_{t+1} = g(\theta_{t+1})$

Identical update to FTRL for linear loss functions.
What about general convex loss functions?

Sub-Gradients

- ▶ we can reduce general convex to linear using the gradient.

Sub-Gradients



- ▶ we can reduce general convex to linear using the gradient.
- ▶ What can we do if $f(x)$ is convex but not differentiable at x ?

Sub-Gradients

- ▶ we can reduce general convex to linear using the gradient.
- ▶ What can we do if $f(x)$ is convex but not differentiable at x ?
- ▶ Use the sub-gradients at $x \doteq \partial f(x)$: the set of linear functions such that $l(x) = \langle w, x \rangle + o$ such that $\forall y, l(y) \leq f(y)$ and $l(x) = f(x)$

Sub-Gradients

- ▶ we can reduce general convex to linear using the gradient.
- ▶ What can we do if $f(x)$ is convex but not differentiable at x ?
- ▶ Use the sub-gradients at $x \doteq \partial f(x)$: the set of linear functions such that $l(x) = \langle w, x \rangle + o$ such that $\forall y, l(y) \leq f(y)$ and $l(x) = f(x)$
- ▶ if gradient $\nabla f(x)$ exists, then $\partial f(x) = \{\nabla f(x)\}$



Example Generalized Online Gradient Descent

Consider the ℓ_2 setup where the functions f_1, f_2, \dots are convex (but not necessarily differentiable). Let η be the learning rate.

$$w_{t+1} = w_t - \eta z_t, \quad z_t \in \partial f_t(w_t)$$

Identical to FTRL with regularization: $R(w) = \frac{1}{2\eta} \|w\|_2^2$

Regret bound on OGD: From FTRL theorem:

$$\begin{aligned} \text{Regret} &\leq \frac{\|u\|^2}{2\eta} + \eta \sum_{t=1}^T \|z_t\|^2 \\ &\leq \frac{B^2}{2\eta} + \eta T L^2 \end{aligned}$$

Gradient based Online Mirror Descent (OMD)

parameter: a link function $g : \mathbb{R}^d \rightarrow S$

initialize: $\theta_1 = 0$

for $t = 1, 2, \dots$

▶ **project** $w_t = g(\theta_t)$

Dual Decent: Instead of minimizing f , minimize ∇f .

Convexity implies equivalence of goals.

Gradient based Online Mirror Descent (OMD)

parameter: a link function $g : \mathbb{R}^d \rightarrow S$

initialize: $\theta_1 = 0$

for $t = 1, 2, \dots$

▶ **project** $w_t = g(\theta_t)$

▶ **update** $\theta_{t+1} = \theta_t - z_t$ where $z_t \in \partial f_t(w_t)$

Dual Decent: Instead of minimizing f , minimize ∇f .

Convexity implies equivalence of goals.

Duality

- ▶ OMD can be analyzed using elementary tools from duality.

Duality

- ▶ OMD can be analyzed using elementary tools from duality.
- ▶ Using Duality Gives better intuition, more general analysis, tighter bounds.

Dual Vector Spaces

- ▶ V is a vector space, with a norm $\|v\|$

Dual Vector Spaces

- ▶ V is a vector space, with a norm $\|v\|$
- ▶ U is the set of all linear mappings from V to V

Dual Vector Spaces

- ▶ V is a vector space, with a norm $\|v\|$
- ▶ U is the set of all linear mappings from V to V
- ▶ The norm of $u \in U$ is defined as

$$\|u\|_* \doteq \max_{v \in V} \frac{\|u(v)\|}{\|v\|}$$

Dual Vector Spaces

- ▶ V is a vector space, with a norm $\|v\|$
- ▶ U is the set of all linear mappings from V to V
- ▶ The norm of $u \in U$ is defined as

$$\|u\|_* \doteq \max_{v \in V} \frac{\|u(v)\|}{\|v\|}$$

- ▶ V is equivalent to the set of all linear mappings from U to U .

Dual Vector Spaces

- ▶ V is a vector space, with a norm $\|v\|$
- ▶ U is the set of all linear mappings from V to V
- ▶ The norm of $u \in U$ is defined as

$$\|u\|_* \doteq \max_{v \in V} \frac{\|u(v)\|}{\|v\|}$$

- ▶ V is equivalent to the set of all linear mappings from U to U .
- ▶ U and V are dual vector spaces, with dual norms.

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$
- ▶ L_∞ norm: $\max_i |x_i|$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$
- ▶ L_∞ norm: $\max_i |x_i|$
- ▶ L_p norm: $(\sum_{i=1}^n x_i^p)^{\frac{1}{p}}$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$
- ▶ L_∞ norm: $\max_i |x_i|$
- ▶ L_p norm: $(\sum_{i=1}^n x_i^p)^{\frac{1}{p}}$
- ▶ L_p, L_q are dual norms if $p, q \geq 1$, and $\frac{1}{p} + \frac{1}{q} = 1$

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$
- ▶ L_∞ norm: $\max_i |x_i|$
- ▶ L_p norm: $(\sum_{i=1}^n x_i^p)^{\frac{1}{p}}$
- ▶ L_p, L_q are dual norms if $p, q \geq 1$, and $\frac{1}{p} + \frac{1}{q} = 1$
- ▶ L_1, L_∞ are dual.

Dual Norms

- ▶ The space is always $U, V = \mathbb{R}^n$
- ▶ The linear operation is the dot product $u \cdot v$
- ▶ $\|u\|_*$ is the dual norm to $\|v\|$,
- ▶ L_2 norm: $\sqrt{\sum_{i=1}^n x_i^2}$
- ▶ L_1 norm: $\sum_{i=1}^n |x_i|$
- ▶ L_∞ norm: $\max_i |x_i|$
- ▶ L_p norm: $(\sum_{i=1}^n x_i^p)^{\frac{1}{p}}$
- ▶ L_p, L_q are dual norms if $p, q \geq 1$, and $\frac{1}{p} + \frac{1}{q} = 1$
- ▶ L_1, L_∞ are dual.
- ▶ L_2 is self-dual.

Lipschitz condition and the dual norm

Lemma 2.6:

Let $f : S \rightarrow \mathbb{R}$ be a convex function. Then, f is L -Lipschitz over S with respect to a norm $\|\cdot\|$ if and only if for all $w \in S$ and $z \in \partial f(w)$ we have:

$$\|z\|_* \leq L$$

where $\|\cdot\|_*$ denotes the dual norm.

Proof of Lemma 2.6

Proof:

Assume that f is L -Lipschitz. For any $w \in S$ and $z \in \partial f(w)$, choose u such that $u - w = \arg \max_{\|v\|=1} \langle v, z \rangle$. Then,

$$\langle z, u - w \rangle = \|z\|_*$$

By the sub-gradient definition,

$$f(u) - f(w) \geq \langle z, u - w \rangle = \|z\|_*$$

Since f is L -Lipschitz,

$$f(u) - f(w) \leq L\|u - w\| = L$$

Combining the inequalities:

$$\|z\|_* \leq L$$

For the converse, assume $\|z\|_* \leq L$ for all $z \in \partial f(w)$. Then,

$$f(u) - f(w) \leq \langle z, u - w \rangle \leq \|z\|_* \|u - w\| \leq L\|u - w\|$$

Hence, f is L -Lipschitz.

Fenchel Duality

- ▶ Suppose $F : A \rightarrow \mathbb{R}$ is a convex function over a convex set $A \subseteq \mathbb{R}^n$.

Fenchel Duality

- ▶ Suppose $F : A \rightarrow \mathbb{R}$ is a convex function over a convex set $A \subseteq \mathbb{R}^n$.
- ▶ The dual function to F is

$$F^*(u) = \sup_{v \in A} (u \cdot v - F(v))$$

Fenchel Duality

- ▶ Suppose $F : A \rightarrow \mathbb{R}$ is a convex function over a convex set $A \subseteq \mathbb{R}^n$.
- ▶ The dual function to F is

$$F^*(u) = \sup_{v \in A} (u \cdot v - F(v))$$

- ▶ Fenchel duality Equivalent to Legendre duality for differentiable functions.

Visualization of the Legendre Dual

Visualization of the Legendre Dual

Visualization of the Legendre Dual

► $x, y \in \mathbb{R}$

Visualization of the Legendre Dual


- ▶ $x, y \in \mathbb{R}$
- ▶ $f^*(y) = \sup_{x \in \mathbb{R}} (xy - f(x))$

Visualization of the Legendre Dual

- ▶ $x, y \in \mathbb{R}$
- ▶ $f^*(y) = \sup_{x \in \mathbb{R}} (xy - f(x))$
- ▶ $-f^*(y) = \inf_{x \in \mathbb{R}} (f(x) - xy)$


Visualization of the Legendre Dual

- ▶ $x, y \in \mathbb{R}$
- ▶ $f^*(y) = \sup_{x \in \mathbb{R}} (xy - f(x))$
- ▶ $-f^*(y) = \inf_{x \in \mathbb{R}} (f(x) - xy)$




Dual of Dual

- The dual of any function is convex.





Dual of Dual

- ▶ The dual of any function is convex.
- ▶ if F is convex then $F^{**} = F$



Gradient Duality (legendre only)



- If the gradient of f at x is k then the gradient of f^* at k is x



Gradient Duality (legendre only)

- ▶ If the gradient of f at x is k then the gradient of f^* at k is x
- ▶ In general:

$$\nabla F^* = (\nabla F)^{-1}$$



Example: Exponential Potential

- Potential: $F(u) = \sum_{i=1}^d e^{u_i}$

Example: Exponential Potential

- ▶ Potential: $F(u) = \sum_{i=1}^d e^{u_i}$
- ▶ Gradient: $\nabla F(u)_i = e^{u_i}$ or $\nabla F(u) = F(u)$.

Example: Exponential Potential

- ▶ Potential: $F(u) = \sum_{i=1}^d e^{u_i}$
- ▶ Gradient: $\nabla F(u)_i = e^{u_i}$ or $\nabla F(u) = F(u)$.
- ▶ Dual: $F^*(v) = \sum_{i=1}^d v_i (\ln v_i - 1)$

Example: Exponential Potential

- ▶ Potential: $F(u) = \sum_{i=1}^d e^{u_i}$
- ▶ Gradient: $\nabla F(u)_i = e^{u_i}$ or $\nabla F(u) = F(u)$.
- ▶ Dual: $F^*(v) = \sum_{i=1}^d v_i (\ln v_i - 1)$
- ▶ Gradient of dual: $\nabla F^*(v)_i = \ln v_i$

Example: Exponential Potential

- ▶ Potential: $F(u) = \sum_{i=1}^d e^{u_i}$
- ▶ Gradient: $\nabla F(u)_i = e^{u_i}$ or $\nabla F(u) = F(u)$.
- ▶ Dual: $F^*(v) = \sum_{i=1}^d v_i (\ln v_i - 1)$
- ▶ Gradient of dual: $\nabla F^*(v)_i = \ln v_i$
- ▶ Note $(\nabla F)^{-1} = \nabla F^*$

Bregman Divergence

Bregman Divergence

Bregman Divergence

- ▶ $R(x)$ is convex and differentiable.

Bregman Divergence


- ▶ $R(x)$ is convex and differentiable.
- ▶ $D_R(w||u) =$
 $R(w) - (R(u) + \langle \nabla R(u), (w - u) \rangle)$

Bregman Divergence

- ▶ $R(x)$ is convex and differentiable.
- ▶ $D_R(w||u) =$
 $R(w) - (R(u) + \langle \nabla R(u), (w - u) \rangle)$
- ▶ The error term of the first order
Taylor expansion around u

Bregman Divergence

- ▶ $R(x)$ is convex and differentiable.
- ▶ $D_R(w||u) =$
 $R(w) - (R(u) + \langle \nabla R(u), (w - u) \rangle)$
- ▶ The error term of the first order Taylor expansion around u



Fenchel and Bregman

- ▶ F : strictly convex with continuous first derivative.

Fenchel and Bregman

- ▶ F : strictly convex with continuous first derivative.
- ▶ F^* is the Fenchel Dual of F

Fenchel and Bregman

- ▶ F : strictly convex with continuous first derivative.
- ▶ F^* is the Fenchel Dual of F
- ▶ D_F, D_{F^*} Bregman divergences wrt F, F^*

Fenchel and Bregman

- ▶ F : strictly convex with continuous first derivative.
- ▶ F^* is the Fenchel Dual of F
- ▶ D_F, D_{F^*} Bregman divergences wrt F, F^*
- ▶ $u' = \nabla F(u)$ and $v' = \nabla F(v)$

Fenchel and Bregman

- ▶ F : strictly convex with continuous first derivative.
- ▶ F^* is the Fenchel Dual of F
- ▶ D_F, D_{F^*} Bregman divergences wrt F, F^*
- ▶ $u' = \nabla F(u)$ and $v' = \nabla F(v)$
- ▶ $D_F(u, v) = D_{F^*}(u', v')$

Mirror Descent - Step 1

Gradient Step in Dual Space:

$$z_{t+1} = \nabla R(w_t) - \eta \nabla f_t(w_t)$$

Here, $\nabla R(w_t)$ maps the point into the dual space.

Mirror Descent - Step 2

Projection Back to Primal Space:

$$w_{t+1} = \arg \min_{w \in S} D_R(w, z_{t+1})$$

Where $D_R(w, z)$ is the Bregman divergence:

$$D_R(w, z) = R(w) - R(z) - \langle \nabla R(z), w - z \rangle$$

This projection ensures w_{t+1} stays within the feasible set S .

Mirror Descent (alternative Notation)

- ▶ Gradient descent in dual space $\theta_t = \theta_{t-1} - \lambda \nabla \ell_t(\theta_{t-1})$

Mirror Descent (alternative Notation)

- ▶ Gradient descent in dual space $\theta_t = \theta_{t-1} - \lambda \nabla \ell_t(\theta_{t-1})$
- ▶ Using duality can be rewritten as

$$\nabla R^*(w_t) = \nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1})$$

Mirror Descent (alternative Notation)

- ▶ Gradient descent in dual space $\theta_t = \theta_{t-1} - \lambda \nabla \ell_t(\theta_{t-1})$
- ▶ Using duality can be rewritten as


$$\nabla R^*(w_t) = \nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1})$$

- ▶ Projection: As ∇R is the inverse of ∇R^* we get

$$w_t = \nabla R(\nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1}))$$

A picture of mirror descent

$$w_t = \nabla R(\nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1}))$$



Regret Bound for OMD

Lemma 2.20. Suppose that OMD is run with a link function $g = \nabla R^*$. Then, its regret is upper bounded by:

$$\sum_{t=1}^T \langle w_t - u, z_t \rangle \leq R(u) - R(w_1) + \sum_{t=1}^T D_{R^*}(-z_{1:t} \| -z_{1:t-1})$$

Furthermore, equality holds for the vector u that minimizes $R(u) + \sum_t \langle u, z_t \rangle$.

Proof: Step 1 - Fenchel–Young Inequality

Using the **Fenchel–Young inequality**, we have:

$$R(u) + \sum_{t=1}^T \langle u, z_t \rangle = R(u) - \langle u, -z_{1:T} \rangle \geq -R^*(-z_{1:T}).$$

Equality holds for u that maximizes $\langle u, -z_{1:T} \rangle - R(u)$, hence minimizing $R(u) + \langle u, z_{1:T} \rangle$.

Proof: Step 2 - Bregman Divergence

Since $w_t = \nabla R^*(-z_{1:t-1})$ and using the definition of the Bregman divergence, we rewrite:

$$-R^*(-z_{1:T}) = -R^*(0) - \sum_{t=1}^T (R^*(-z_{1:t}) - R^*(-z_{1:t-1})).$$

Rearranging, we get:

$$= -R^*(0) + \sum_{t=1}^T (\langle w_t, z_t \rangle - D_{R^*}(-z_{1:t} \| -z_{1:t-1})).$$

Final Step

Note: Since

$$R^*(0) = \max_w \langle 0, w \rangle - R(w) = - \min_w R(w) = -R(w_1),$$

combining all the above, we conclude the proof. \square

OMD for ℓ_2

- Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$
- ▶ We use the dual descend algorithm for the half quadratic regularizer

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$
- ▶ We use the dual descend algorithm for the half quadratic regularizer
- ▶ Regularizer: $R(u) = \frac{1}{2}\|u\|_2^2$

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$
- ▶ We use the dual descend algorithm for the half quadratic regularizer
- ▶ Regularizer: $R(u) = \frac{1}{2}\|u\|_2^2$
- ▶ Legendre Dual Regularizer $R^*(u) = \frac{1}{2}\|u\|_2^2$

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$
- ▶ We use the dual descend algorithm for the half quadratic regularizer
- ▶ Regularizer: $R(u) = \frac{1}{2}\|u\|_2^2$
- ▶ Legendre Dual Regularizer $R^*(u) = \frac{1}{2}\|u\|_2^2$
- ▶ **Gradient Mapping:** $\nabla R(w) = w$

OMD for ℓ_2

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_2 \leq X_2$
- ▶ The dual norm for the weights is $\|w\|_2$
- ▶ We use the dual descend algorithm for the half quadratic regularizer
- ▶ Regularizer: $R(u) = \frac{1}{2}\|u\|_2^2$
- ▶ Legendre Dual Regularizer $R^*(u) = \frac{1}{2}\|u\|_2^2$
- ▶ **Gradient Mapping:** $\nabla R(w) = w$
- ▶ Gradient step: $z_{t+1} = w_t - \eta \nabla f_t(w_t)$

Projection Step for ℓ_2 Norm

Bregman Divergence:

$$D_R(w, z) = \frac{1}{2} \|w - z\|_2^2$$

Projection Back to Primal Space:

$$w_{t+1} = \Pi_S(z_{t+1}) = \arg \min_{w \in S} \frac{1}{2} \|w - z_{t+1}\|_2^2$$

Where Π_S denotes the Euclidean projection onto the feasible set S .

Final Update Rule for ℓ_2 Norm

Combining both steps, the final update rule becomes:

$$w_{t+1} = \Pi_S (w_t - \eta \nabla f_t(w_t))$$

This is equivalent to the standard **Projected Gradient Descent** for the ℓ_2 norm.

Optimal Tuning for η and Regret Bound

Regret Bound:

$$\text{Regret}_T(u) \leq \frac{\|u\|_2^2}{2\eta} + \frac{\eta}{2} \sum_{t=1}^T \|\nabla f_t(w_t)\|_2^2$$

Assuming $\|u\|_2 \leq B$ and $\|\nabla f_t(w_t)\|_2 \leq L$, this simplifies to:

$$\text{Regret}_T(u) \leq \frac{B^2}{2\eta} + \frac{\eta L^2 T}{2}$$

Optimal η :

$$\eta^* = \frac{B}{L\sqrt{T}}$$

Resulting Regret Bound:

$$\text{Regret}_T(u) \leq BL\sqrt{T}$$

ℓ_∞ OMD

- Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_\infty \leq X_\infty$

ℓ_∞ OMD

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_\infty \leq X_\infty$
- ▶ The dual norm for the weights is $\|w\|_1$

ℓ_∞ OMD

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_\infty \leq X_\infty$
- ▶ The dual norm for the weights is $\|w\|_1$
- ▶ Suppose we use the dual descend algorithm for the exponential regularizer

ℓ_∞ OMD

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_\infty \leq X_\infty$
- ▶ The dual norm for the weights is $\|w\|_1$
- ▶ Suppose we use the dual descend algorithm for the exponential regularizer
- ▶ Regularizer: $R(u) = \frac{1}{\eta} \ln \sum_{i=1}^d e^{\eta u_i}$

ℓ_∞ OMD

- ▶ Problem: OCO where $\forall z \in \partial f(x) \ \|z\|_\infty \leq X_\infty$
- ▶ The dual norm for the weights is $\|w\|_1$
- ▶ Suppose we use the dual descend algorithm for the exponential regularizer
- ▶ Regularizer: $R(u) = \frac{1}{\eta} \ln \sum_{i=1}^d e^{\eta u_i}$
- ▶ Legendre Dual Regularizer $R^*(u) = \sum_{i=1}^d u_i (\ln u_i - 1)$

- Weight update:

$$w_t = \nabla R(\nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1}))$$

- ▶ Weight update:

$$w_t = \nabla R(\nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1}))$$

- ▶ Unnormalized exponentiated gradient:

$$w_{i,t} = w_{i,t-1} e^{-\lambda \nabla \ell_t(w_{i-1})}$$

- Weight update:

$$w_t = \nabla R(\nabla R^*(w_{t-1}) - \lambda \nabla \ell_t(w_{t-1}))$$

- Unnormalized exponentiated gradient:

$$w_{i,t} = w_{i,t-1} e^{-\lambda \nabla \ell_t(w_{i-1})}$$

- Normalized exponentiated gradient:

$$w_{i,t} = \frac{w_{i,t-1} e^{-\lambda \nabla \ell_t(w_{i-1})}}{\sum_{j=1}^d w_{j,t-1} e^{-\lambda \nabla \ell_t(w_{j-1})}}$$

- ▶ Normalization corresponds to projection on the simplex using the Bregman divergence according to R^* .

- ▶ Normalization corresponds to projection on the simplex using the Bregman divergence according to R^* .
- ▶ The dual descend algorithm for the exponential regularizer function R and the learning rate $\lambda = \frac{2\epsilon}{X_\infty^2}$ for some $0 < \epsilon < 1$

- ▶ Normalization corresponds to projection on the simplex using the Bregman divergence according to R^* .
- ▶ The dual descend algorithm for the exponential regularizer function R and the learning rate $\lambda = \frac{2\epsilon}{X_\infty^2}$ for some $0 < \epsilon < 1$
- ▶ yields **Loss Bound**:

$$L_{A,T} \leq \frac{L_T(u)}{1 - \epsilon} + \frac{X_\infty^2 \ln d}{2\epsilon(1 - \epsilon)}$$