



Review

Amine degradation in CO₂ capture. I. A reviewC. Gouedard ^a, D. Picq ^{a,b}, F. Launay ^c, P.-L. Carrette ^{a,*}^a IFP Energies nouvelles, Rond-point de l'échangeur de Solaize, BP 3, 69360 Solaize, France^b LMOPS/LEPMI, chemin du canal, 69360 Solaize, France^c LRS/UPMC-Paris 6, 3 rue Galilée, 94200 Ivry-sur-Seine, France

ARTICLE INFO

Article history:

Received 7 March 2012

Received in revised form 21 June 2012

Accepted 23 June 2012

Available online 17 July 2012

Keywords:

Post-combustion CO₂ capture

Amine degradation

MEA/DEA/MDEA/PZ/AMP

SO_x and NO_x

Mechanisms

ABSTRACT

Post-combustion CO₂ capture based on CO₂ absorption by aqueous amine solutions is the most mature gas separation technology. A main problem is amine degradation due to heat, CO₂, O₂, NO_x and SO_x. This review proposes to make a critical survey of literature concerning degradation, to list degradation products and to discuss mechanisms proposed by authors. Benchmark molecule is monoethanolamine (MEA) but diethanolamine (DEA), N-methyldiethanolamine (MDEA), piperazine (PZ) and 2-amino-2-methylpropan-1-ol (AMP) are also studied. Uses of other amines and amine blends are also considered. In the case of MEA, ammonia, N-(2-hydroxyethyl)-piperazin-3-one (HEPO) and N-(2-hydroxyethyl)-2-(2-hydroxyethylamino) acetamide (HEHEAA) are the main identified degradation products in pilot plants. Among lab studies, the most cited degradation products are ammonia, carboxylic acids, N-(2-hydroxyethyl)-formamide (HEF), N-(2-hydroxyethyl)-acetamide (HEA) and N-(2-hydroxyethyl)-imidazole (HEI) for oxidative degradation, and oxazolidin-2-one (OZD), N-(2-hydroxyethyl)-ethylenediamine (HEEDA) and N-(2-hydroxyethyl)-imidazolidin-2-one (HEIA) for thermal degradation. Numerous degradation products have been identified but some are still unknown. A lot of degradation mechanisms have been proposed but some are missing or need proofs. SO_x and NO_x effects are still few examined and much work remains to be done concerning volatile degradation products potentially emitted to atmosphere: their identification and their formation mechanisms need further investigations.

© 2012 Elsevier Ltd. All rights reserved.

Contents

1. Introduction	245
2. Studied amines	245
2.1. Monoethanolamine (MEA)	245
2.1.1. Thermal degradation.....	245
2.1.2. Oxidative degradation	247
2.2. Diethanolamine (DEA).....	252
2.2.1. Thermal degradation.....	254
2.2.2. Oxidative degradation	256
2.3. Methyldiethanolamine (MDEA).....	258
2.3.1. Thermal degradation.....	258
2.3.2. Oxidative degradation	258
2.4. Piperazine (PZ).....	258
2.4.1. Thermal degradation.....	258
2.4.2. Oxidative degradation	258

* Corresponding author at: IFP Energies nouvelles – Catalysis and Separation Division, Lyon Establishment, Rond-point de l'échangeur de Solaize, BP 3, 69360 Solaize, France. Tel.: +33 4 37 70 27 23; fax: +33 4 37 70 20 66.

E-mail address: p-louis.carrette@ifpen.fr (P.-L. Carrette).