



A general review of CO₂ sequestration in underground geological formations and assessment of depleted hydrocarbon reservoirs in the Niger Delta[☆]

Patrick A. Eigbe^a, Olatunbosun O. Ajayi^a, Olabode T. Olakoyejo^a, Opeyemi L. Fadipe^b, Steven Efe^c, Adekunle O. Adelaja^{a,d,*}

^a Department of Mechanical Engineering, University of Lagos, Akoka, Lagos State 100213, Nigeria

^b Department of Industrial and Systems Engineering, Morgan State University, 1700 East Cold Spring Lane, Baltimore, Maryland, USA

^c Department of Civil Engineering, Morgan State University, 1700 East Cold Spring Lane, Baltimore, Maryland, USA

^d Faculty of Engineering and the Built Environment, Durban University of Technology, PO Box 1334, Durban 4000, South Africa

HIGHLIGHTS

- Niger Delta depleted oil/gas reservoir CO₂ capture, utilization & storage reviewed
- Potential issues relating to CO₂ sequestration and modeling in Niger Delta noted
- Appropriate CFD solution should combine static and dynamic model.
- Research gaps identified and future directions recommended.

ARTICLE INFO

Keywords:

Carbon capture
geological storage
geo-mechanical modeling
multiphase flow
CO₂ Sequestration
Niger Delta

ABSTRACT

This paper investigates the viability of CO₂ storage in geological formations, including depleted hydrocarbon reservoirs applying 3-dimensional seismic and well data of the Niger Delta region as a case study for CO₂ sequestration, which represents an essential initiative for the reduction of greenhouse gas emissions. Different theoretical and experimental studies from literature on CO₂ sequestration in geological formations across the world, including the Niger Delta, are reviewed. The Niger Delta basin has a high potential for CO₂ sequestration, and this indication is shown through the review of research papers undertaken outside and within the basin, which reveal the presence of massive hydrocarbon fields, lateral continuity of reservoir, reservoir-seal pairs, faults, and traps, developed hydrocarbon fields, appropriate reservoir depth. The gaps identified from the review of various research studies are analyzed to form the basis of a future research investigation for capturing, removing, and storing CO₂ in depleted hydrocarbon reservoirs and other geological formations. The authors deduced that accurate injection pressure, rate, and depth estimation are critical factors for CO₂ sequestration and need to be thoroughly investigated.

1. Introduction

Carbon dioxide capture and sequestration are moderately considered fundamental and practical methods of reducing the release of anthropogenic greenhouse gases (GHGs) on the earth [1]. Proposals have been made to indicate that it is an instantly realizable and scientifically realistic method of decreasing CO₂ emissions [2] and has the prospect of

making a significant decrease in carbon emissions from point sources [3]. The approach is associated with the capture of CO₂ from a stationary source, transportation through pipelines, introduction into, and storage in suitable underground geological structures in sedimentary basins (Fig. 1). The introduction of CO₂ into depleted hydrocarbon fields is an age-long integral part of the enhanced oil recovery (EOR) practice. CO₂ confinement in the geological media has been proven to be realizable. The main types of storage media are saline aquifers, depleted oil

[☆] “The short version of the paper was presented at CUE2022. This paper is a substantial extension of the short version of the conference paper.”

* Corresponding author.

E-mail address: aadelaja@unilag.edu.ng (A.O. Adelaja).

<https://doi.org/10.1016/j.apenergy.2023.121723>

Received 12 May 2023; Received in revised form 2 July 2023; Accepted 2 August 2023

Available online 18 August 2023

0306-2619/© 2023 Elsevier Ltd. All rights reserved.