

1. Derive the following asymptotic moments associated with $m_3(x)$: mean, variance, first order auto-correlation. Furthermore, compute $\nabla_b g(b_0)$. Which moments are informative for estimating b ?

Answer: We obtain the asymptotic moments as follows:

$$\begin{aligned}
 \mathbb{E}[x_t] &= \mathbb{E}[\rho_0 x_{t-1} + \epsilon_t] = \rho_0 \mathbb{E}[x_{t-1}] + \mathbb{E}[\epsilon_t] = \rho_0 \mathbb{E}[\rho_0 x_{t-2} + \epsilon_{t-1}] = \rho_0^t \mathbb{E}[x_0] = 0. \\
 \mathbb{E}[(x_t - \mathbb{E}[x_t])^2] &= \mathbb{E}[x_t^2] = \mathbb{E}[(\rho_0 x_{t-1} + \epsilon_t)^2] \\
 &= \rho_0^2 \mathbb{E}[x_{t-1}^2] + 2\rho_0 \mathbb{E}[x_{t-1}\epsilon_t] + \mathbb{E}[\epsilon_t^2] \\
 &= \rho_0^2 \mathbb{E}[(\rho_0 x_{t-2} + \epsilon_{t-1})^2] + \sigma_0^2 \\
 &= \rho_0^{2T} \mathbb{E}[x_0^2] + \sigma_0^2 \sum_{i=0}^{T-1} (\rho_0^2)^i \rightarrow \frac{\sigma_0^2}{1 - \rho_0^2} = \frac{4}{3}. \\
 \mathbb{E}[(x_t - \mathbb{E}[x_t])(x_{t-1} - \mathbb{E}[x_{t-1}])] &= \mathbb{E}[x_t x_{t-1}] \\
 &= \rho_0 \mathbb{E}[x_{t-1}^2] + \mathbb{E}[\epsilon_t x_{t-1}] \\
 &= \rho_0 \mathbb{E}[(\rho_0 x_{t-2} + \epsilon_{t-1})^2] \\
 &= \rho_0^3 \mathbb{E}[x_{t-2}^2] + \rho_0 \mathbb{E}[\epsilon_{t-1}^2] \\
 &= \rho_0^{2T-1} \mathbb{E}[x_0^2] + \rho_0 \sigma_0^2 \sum_{i=0}^{T-2} (\rho_0^2)^i \rightarrow \frac{\rho_0 \sigma_0^2}{1 - \rho_0^2} = \frac{2}{3}
 \end{aligned}$$

Both the variance and the first order correlation are informative for estimating b . ■

2. Simulate a series of “true” data of length $T = 200$ using (i). We will use this to compute $M_T(x)$.
3. Set $H = 10$ and simulate H vectors of length $T = 200$ random variables e_t from $N(0, 1)$. We will use this to compute $M_{TH}(y(b))$. Store these vectors. You will use the same vector of random variables throughout the entire exercise. Since this exercise requires you to estimate σ^2 , you want to change the variance of e_t during the estimation. You can simply use σ_{e_t} when the variance is σ^2 .
- 4.
5. Next we estimating the $l = 2$ vector b for the just identified case where m_2 uses the variance and autocorrelation. Given what you found in part (i), do you now think there will be a problem? If not, hopefully the standard error of the estimate of b as well as the J test will tell us something. Let’s see. For this case, perform steps (a)-(d) above.
6. Next, we will consider the overidentified case where m_3 uses the mean, variance and autocorrelation. Let’s see. For this case, perform steps (a)-(d) above. Furthermore, bootstrap the the finite sample distribution of the estimators using the following algorithm:
 - i. Draw ϵ_t and e_t^h from $N(0, 1)$ for $t = 1, 2, \dots, T$ and $h = 1, 2, \dots, H$. Compute $(\hat{b}_{TH}^1), \hat{b}_{TH}^2$ as described.
 - ii. Repeat (e) using another seed.