Using directed evolution to simplify the production of tyrian purple in *E.coli*




Background

Tyrian purple is a dye extracted from the murex shellfish. The glands of thousands of putrefied crushed shellfish were left to bake in the sun, which could produce only a few grams of tyrian purple powder. The resulting liquid was used to dye cloth fibres in manipulated variations of colours ranging from pink to violet.




Optimize the two-cell production system to
One-cell system.
Introduce a reversible glucose protection group to make the tyrian purple dye more convenient for industrial scale-up.

Purpose



Revo_ution



- Incorporate the glycosyltransfer -ase for a scalable production of halogenated indican and achieve a sustainable dyeing method using β-glucosidase.
- Develop a synthetic phylogeny of programmable Trp repressor with directed evolution to simplify the two-cell production system.
- · Iterative rounds of positive and negative compartmentalized partnered replication (CPR) led to the exponential amplification of variants that responded with high affinity and specificity to halogenated tryptophan analogs and novel operator sites.

Design

Collaborations





灵蛛科技 LINK SPIDER









南方科技大学生命科学学院 SUSTech·SCHOOL OF **LIFE SCIENCES** Lee, J., Kim, J., Song, J.E. et al. Production of Tyrian purple indigoid dye from tryptophan in Escherichia coli. Nat Chem Biol 17, 104-112 (2021).

 Yongfei Liu, Huiling Yuan, Dongqin Ding, Huina Dong, Qinhong Wang, and Dawei Zhang. Establishment of a Biosensor-based High-Throughput Screening Platform for Tryptophan Overproduction. ACS Synthetic Biology 10 (6), 1373-1383 (2021).
 Ellefson, J.W., Ledbetter, M.P. & Ellington, A.D. Directed evolution of a synthetic phylogeny of programmable Trp repressors. Nat Chem Biol 14, 361-367 (2018).

 Abil, Z., Ellefson, J., Gollihar, J. et al. Compartmentalized partnered replication for the directed evolution of genetic parts and circuits. Nat Protoc 12, 2493–2512 (2017).

References