

# Solution For The School Geometry Problems

Yogesh Choudhary

May 31, 2020

# Question

## Exercise 8.1(Q no.36)

Two sides AB and BC and median AM of one triangle ABC are respectively equal to sides PQ and QR and median PN of  $\triangle PQR$ . Show that

a)  $\triangle ABM \cong \triangle PQN$

b)  $\triangle ABC \cong \triangle PQR$

# Codes and Figures

The python code for the figure is


```
./code/Traingle.py
```

The latex- tikz code is


```
./figs/triangle.tex
```

The above latex code can be compiled as standalone document

```
./figs/triangle_fig.tex
```



(a) By Python



(b) By Latex-tikz

# Construction method

The tables below are the values used for constructing the triangles in both Python and Latex-Tikz.

| Initial Input Values. |   |
|-----------------------|---|
| a, p                  | 3 |
| b, q                  | 5 |
| c, r                  | 6 |

**Table:** To construct  $\triangle ACB$  and  $\triangle PQR$

The steps for constructing  $\triangle ACB$  are

$$(i)\mathbf{A} = \begin{pmatrix} 3.33 \\ 4.99 \end{pmatrix} (ii)\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} (iii)\mathbf{C} = \begin{pmatrix} 3 \\ 0 \end{pmatrix}$$

$$(i)\mathbf{P} = \begin{pmatrix} 8.33 \\ 4.99 \end{pmatrix} (ii)\mathbf{Q} = \begin{pmatrix} 5 \\ 5 \end{pmatrix} (iii)\mathbf{R} = \begin{pmatrix} 8 \\ 0 \end{pmatrix}$$

**M** and **N** are the midpoints of BC and QR respectively

$$\mathbf{M} = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix}, \mathbf{N} = \begin{pmatrix} 6.5 \\ 0 \end{pmatrix}$$

| Derived Values for <i>triangleDCB</i> . |  |
|---|--|
| <b>M</b>                                | $\begin{pmatrix} 1.5 \\ 0 \end{pmatrix}$ |
| <b>N</b>                                | $\begin{pmatrix} 6.5 \\ 0 \end{pmatrix}$ |

**Table:** To construct medians AN and PN

# Solution

given that  $\rightarrow$

$$AB = PQ = c = r$$

$$BC = QR = a = p$$

$$AM = PN = m = n$$

Therefore  $\rightarrow$

**M** and **N** are the position vector of mid-point of **BC** and **QR** respectively .

$$\frac{1}{2}BC = \frac{1}{2}QR$$

## Solution a)

from triangles  $ABM$  and  $PQR \rightarrow$

$$AB = PQ(\text{given})$$

$$AM = PN(\text{given})$$

$$MB = NQ$$

(Both  $m$  and  $N$  are the mid points)

$\Rightarrow \triangle ABM$  and  $\triangle PQN$  are congruent to each other by SSS congruency.

Hence, proved

## Solution b)

given that  $\rightarrow$

$$MC = NR$$

$$\therefore \triangle ABM \cong \triangle PQN$$

$$\implies \angle AMC = \angle PNR$$

from SAS congruency  $\rightarrow$

$$\triangle AMC \cong \triangle PNR$$

$$\implies AC = PR$$

Thus from the triangle ABC ,PQR and by SSS congruency

$$\triangle ABC \cong \triangle PQR$$

Hence proved