Semiempirical Orbital-Optimized DFT in MOUSE: Spin-Pure Excited-State Surfaces with Analytic Gradients and Hessians

Vincenzo Barone
Consorzio Interuniversitario Nazionale per la Scienza e Tecnologia dei Materiali

Working manuscript, 1 November 2026

Abstract

An alternative to a semiempirical Tamm–Dancoff description is to optimize an excited-state energy directly with respect to molecular orbitals and differentiate the resulting stationary surface. Inspired by direct orbital optimization and electronic saddle searches developed by Levi and co-workers, we introduce this route in MOUSE using a GFN2-based energy informed by projected ground-state density-functional response. Spin polarization enters a restricted open-shell Kohn–Sham construction with a spin-pure singlet configuration state function. Two complementary effective energy models are combined with fixed equal weights and relative active-orbital constraints, without an additive excitation-energy correction. An analytic constrained orbital response yields nuclear gradients and Hessians, including metric, local-frame, and ground-reference contributions. For a curated set of 19 vertical transitions in 18 molecules, the mean absolute error is 0.183 eV. Four prospective tests give 0.304 eV. A separate completed 14-state extension gives MAEs of 0.492 eV for the original OO model and 0.261 eV for the Q-parent TDA control on corrected GFN2 orbitals, exposing a limitation for multiconfigurational $\pi\pi^*$ states. A minimal-CSF development candidate reduces the extension MAE to 0.228 eV on already exposed data. A subsequent frozen six-state cohort accepts four states and gives 0.675 eV, exposing limits of transferability; the other two states lack adequate compact-space representation. Six excited-state molecular minima, including a symmetry-broken cyclopentadiene descendant, have positive vibrational curvatures; complete finite-difference Hessian checks on nitroxyl and formamide agree within 1.2×10^{-8} $E_\mathrm{h}/\mathrm{bohr}^2$. Prepared OO energies are slower than matched GFN2-orbital TDA for all six measured sizes; the method targets differentiated stationary surfaces. The results establish a semiempirical OO-DFT candidate for state-specific potential-energy surfaces and analytic second derivatives, while leaving broad chemical transferability to further validation.

1 Introduction

The cost of an excited-state calculation is set by the scientific task, not only by the cost of a single vertical energy. Conformer and solvent ensembles, excited-state geometry searches, and local vibrational analyses require repeated evaluations on related molecular structures. A compact, parameterized electronic representation can make these repeated tasks accessible while retaining explicit orbitals, charge redistribution, and a differentiable energy. The purpose of a semiempirical model is therefore to allocate a finite computational budget across the structures and derivatives needed for a molecular prediction. It is not to replace high-level reference calculations indiscriminately. Its usefulness requires a controlled balance of electronic error, nuclear sampling, and computational cost.

The simplified excited-state methods of Grimme and co-workers provide a major reference point. The original sTDA approximates the response matrix and screens the excitation space while retaining a DFT ground-state reference; sTD-DFT extends the simplified construction

beyond the Tamm–Dancoff approximation.[1, 2] These are approximations to an excited-state response problem and need not imply a semiempirical ground-state Hamiltonian. In sTDA-xTB, a dedicated tight-binding Hamiltonian and an augmented basis replace the expensive ground-state reference, enabling a predominantly semiempirical spectroscopy workflow.[3] That reference is not interchangeable with an ordinary GFN2 ground-state calculation: the description of the virtual space and the response parameterization are part of the method.

This family has continued to develop. The 2026 std2 description includes sTDA, sTD-DFT, their exact-integral simplified variants XsTDA and XsTD-DFT, and spin-flip formulations, with applications to a broad range of optical response properties.[4] These developments make simplified response methods strong alternatives for multi-root spectroscopy and realistic molecular environments. The present work addresses a complementary objective: an explicitly optimized semiempirical excited-state energy and its analytic nuclear curvature. Its matched GFN2 TDA controls isolate this choice within the MOUSE energy models. A separate direct comparison below evaluates published sTDA-xTB; XsTD-DFT is not benchmarked.

Other low-cost approaches already address excited-state structure and dynamics. TD-DFTB and its long-range-corrected and hybrid extensions combine a compact DFT-derived Hamiltonian with an excited-state response treatment. Recent hybrid-functional DFTB parameterizations target organic photovoltaic molecules, illustrating that frontier levels and charge transfer remain active areas of model development.[5, 6] TDDFT+TB instead keeps a DFT ground state and simplifies the coupling matrix; analytic excited-state gradients were reported in 2023.[7] The approximate auxiliary-function TDDFT-aas approach added an alternative efficient response construction with analytic gradients in 2024.[8] These DFT-reference approaches are relevant cost-reduction strategies, although they do not make the entire calculation semiempirical.

A particularly close precedent is ΔDFTB. Kowalczyk, Le, and Irle implemented state-specific excited-state energies and gradients within DFTB, with benchmarks of geometries, Stokes shifts, and vibrational frequencies.[9] The method remains part of the modern DFTB+ framework.[5] A direct semiempirical excited-state energy is therefore an established idea, not a first introduced here. Orthogonalization-corrected OM2/OM3 and ODM2/ODM3 Hamiltonians combined with multireference configuration interaction (MRCI) provide a different established route to excited-state energies and molecular structure, with a wavefunction expansion rather than a single targeted orbital-optimized CSF.[10, 11] Their treatment of several configurations is especially relevant when evaluating the eventual scope of a ROKS-based model.

The methodological motivation comes from the orbital-optimization studies of Levi and co-workers. Levi, Ivanov, and Jónsson formulated variational excited-state calculations as direct optimization in orbital space, including convergence to electronic saddle points rather than relying exclusively on self-consistent-field iterations.[12, 13] Their work separates the definition of a stationary electronic state from the algorithm used to locate it. Ivanov and co-workers extended this approach to grid and plane-wave representations and discussed excited-state atomic forces.[14] Generalized mode following subsequently provided a way to target a specified saddle order by reversing selected electronic curvature directions.[15] The more recent freeze-and-release strategy uses an initial relaxation with selected excitation orbitals fixed, followed by their release in a full saddle search.[16] Together, these contributions motivate an excited-state solver built around an energy, its orbital gradient, and its curvature.

This perspective is particularly attractive for a semiempirical platform. A ground-state engine already supplies much of the required infrastructure: the energy, Fock matrix, density response, overlap derivatives, and nuclear derivatives. An orbital-optimized excited-state method can reuse these quantities, provided the state-specific orbital equations and their response are added consistently. Reusing the engine therefore means reusing energy and derivative primitives; it does not make an unmodified ground-state Hessian an excited-state Hessian.

MOUSE, the Molecular Orbital Utility for Semiempirical Explorations, supplies a modular environment for such a development. Here its GFN2-based energy is augmented by frozen

operators obtained from projected ground-state DFT information. GFN2-xTB provides the semiempirical foundation,[17] while restricted open-shell Kohn–Sham (ROKS) ideas provide a spin-adapted open-shell singlet construction.[18] The resulting method is called *semiempirical orbital-optimized DFT* (OO-DFT) in the sense that a density-functional-response-informed semiempirical energy is made stationary with respect to the excited-state orbitals. It is not a full-basis Kohn–Sham calculation at each molecular geometry.

Against this background, our contribution is the combination of a GFN2-based, DFT-response-informed energy, a spin-pure ROKS CSF, direct orbital stationarity with state following, and an analytic constrained response through the nuclear Hessian. The contribution concerns this explicit formulation and implementation, rather than priority for semiempirical excited-state gradients or orbital optimization in general. The numerical examples address three distinct questions: the accuracy of vertical energies, the existence of tractable molecular minima, and the numerical correctness of the derivatives. TDA is retained as a matched control, using semiempirical rather than DFT orbitals. No general replacement of TDA for all excited-state tasks is implied.

Figure 1 summarizes the flow from frozen reference information and the native semiempirical engine to the stationary excited-state surface and its derivatives.




Figure 1: Architecture of the semiempirical OO-DFT method in MOUSE. Ground-state DFT information defines two frozen effective models, while the native GFN2 engine supplies energy and derivative primitives. Common-orbital ROKS CSFs enter a fixed 50/50 constrained stationary problem. Coupled orbital response and the ground-reference terms of Eq. (9) yield the energy, analytic gradient, and analytic nuclear Hessian.

2 Theory and implementation

2.1 Two density-functional-response-informed semiempirical energies

Let $E_{\text{native}}[P^\alpha, P^\beta; R]$ denote the native GFN2-based energy at nuclear coordinates R. The retained native terms include the underlying electronic energy and its charge dependence, anisotropy, repulsion, and dispersion contributions. Each corrected parent energy E_i, $i \in \{F, Q\}$, adds a frozen one-body operator and charge/spin response corrections in a complete symmetric

AO-pair representation. In the conventions used here,

$$E_i = E_{\text{native}} + v_i^{\mathsf{T}} q + \tfrac{1}{2}(q - q_0)^{\mathsf{T}} K_c (q - q_0) + \tfrac{1}{2} m^{\mathsf{T}} K_s m, \tag{1}$$

$$q_a = \mathrm{Tr}[B_a(P^\alpha + P^\beta)], \qquad m_a = \mathrm{Tr}[B_a(P^\alpha - P^\beta)]. \tag{2}$$

Here B_a are metric-dependent AO-pair moment operators. The correction K_c replaces the corresponding native charge response by the projected DFT response, avoiding double counting. The spin channel is retained explicitly. Spin polarization is therefore part of the energy functional and its response, rather than a correction applied to a final excitation energy.

At a reference geometry, an isometric map embeds the semiempirical valence space in the DFT AO space after occupied core orbitals have been removed. The parent F uses the complete projected one-body Fock correction. Parent Q uses a qualified spectral one-body correction: DFT orbital-energy spacings are transferred only to semiempirical orbitals with sufficiently reliable individual correspondences, and the remaining levels keep their native spacings. A single occupied-centroid gauge aligns the orbital-energy origins. Neither construction uses reference excitation energies.

The reduced validation extension adds a molecular reference-selection rule at the frozen anchor: retain the original spectral operator if its native-canonical occupied/virtual gap is positive; otherwise use the complete projected DFT ground-state Fock operator for Q. The same choice applies to every target state of that molecule. Charge/spin kernels, weights and exchange scaling are unchanged, and neither target excitation energies nor energy errors enter the choice. This operator revision is assessed separately from the original 19-state campaign; its subsequent full-frame molecular derivative qualification is reported below.

Both parents use the same complete projected charge/spin response. The underlying DFT reference is PBE0.[19] The exchange scaling inherited from the uracil protocol is fixed at $\lambda = 0.9$: it multiplies the PBE0 exact-exchange contribution in the response once, giving an effective response coefficient $0.9 \times 0.25 = 0.225$. It does not multiply orbital gaps, and the semilocal exchange-correlation response is not rescaled. Thus the parameter has the same meaning in the OO-DFT and TDA calculations.

2.2 Spin-pure singlet and orbital stationarity

For an assigned occupied-to-virtual promotion $h \to p$, the spatial orbitals are shared between the mixed-spin determinant M and its associated triplet T. The two determinants obtained by exchanging the open-shell spins define the normalized singlet configuration state function (CSF). With a consistent determinant phase convention,

$$|\Psi_S\rangle = 2^{-1/2}(|D_{\alpha\beta}\rangle - |D_{\beta\alpha}\rangle), \qquad \hat{S}^2|\Psi_S\rangle = 0. \tag{3}$$

The ROKS energy of each parent is evaluated by the multiplet-sum expression

$$E_{S,i}(C_i, R) = 2E_{M,i}(C_i, R) - E_{T,i}(C_i, R). \tag{4}$$

The common spatial orbitals in Eq. (4) are essential. Spin purity is a property of the CSF ansatz, whereas the multiplet-sum energy remains an approximate density-functional construction. This procedure is not exact spin projection of a separately optimized, unrestricted broken-symmetry determinant.

Metric-preserving orbital rotations $C_i(\kappa_i)$ are used to solve the stationary equations. An excited solution can be a saddle in orbital space, and its order need not be one. Direct optimization and mode-following ideas therefore enter the electronic solver, following the motivation of Refs. 13, 15. The number of negative electronic curvature directions is evaluated in the appropriate orbital tangent space. Its value is distinct from the vibrational stability of the nuclear geometry: an electronic saddle may define a nuclear minimum. Occupation and CSF restrictions also mean that a targeted state can have zero negative curvature within its allowed orbital manifold.

The difference between electronic and nuclear stability is illustrated schematically in Fig. 2.




Figure 2: Electronic stationarity and nuclear relaxation are different problems. (a) A schematic orbital-energy saddle contains stable and unstable directions. (b) The corresponding stationary branch can have positive nuclear curvature. The plots use a local quadratic illustration, $E = \frac{1}{2}\kappa_1^2 - \frac{1}{2}\kappa_2^2 + \frac{1}{2}R^2 + 0.45R\kappa_1$, and its relaxed branch; they are not molecular calculation data. The lower expression summarizes the analytic constrained response used for the actual model. The displayed single unstable direction is illustrative; molecular electronic indices vary.

2.3 A fixed 50/50 composite and relative active-pair constraints

The two parent energies are combined with fixed weights,

$$\overline{E}_S = \tfrac{1}{2}E_{S,F} + \tfrac{1}{2}E_{S,Q}. \tag{5}$$

Averaging independent vertical energies does not by itself define a uniquely followed composite molecular surface. To retain state coherence while allowing relaxation, we constrain the relative orientations of the two assigned active orbital pairs. At the anchor geometry, the independently stationary parent states define

$$\tau_x = \left| c_{x,F}^\mathsf{T} S c_{x,Q} \right|^2, \qquad x = h, p. \tag{6}$$

At displaced geometries, the final constraints are

$$g_x(C_F, C_Q, R) = \left| c_{x,F}^\mathsf{T} S(R) c_{x,Q} \right|^2 - \tau_x = 0. \tag{7}$$

These quantities are invariant to individual orbital signs. Both absolute active-orbital orientations and the spectator orbitals remain free to relax. The two parents need not have identical active orbitals. For an overlap indistinguishable from unity within 10^{-10}, a common orbital is eliminated exactly to avoid a singular constraint Jacobian.

The stationary problem is expressed through

$$\mathcal{L}(\kappa, \mu; R) = \overline{E}_S(\kappa, R) + \mu^\mathsf{T} g(\kappa, R), \qquad \mathcal{L}_\kappa = 0, \quad g = 0. \tag{8}$$

The independently stationary anchor states satisfy Eq. (7) by construction with zero multipliers. Consequently, the anchor energy equals the original independent 50/50 mean to numerical

precision. No energy offset is introduced. Away from the anchor, it is the joint Lagrangian gradient that vanishes; the independent parent gradients generally do not.

The final persistent constraints differ from the temporary frozen orbitals in freeze-and-release initialization.[16] They define the composite surface itself. The method is accordingly a constrained composite semiempirical OO-DFT model, rather than unconstrained optimization of a single standard density functional or a formal ensemble-DFT functional.

2.4 Ground-surface referencing and geometry transport

Three restricted closed-shell minima provide the native and corrected ground-state energies. The excited-state molecular surface is

$$E_{\mathrm{exc}}(R) = E_{0,\mathrm{native}}(R) + \tfrac{1}{2} \sum_{i=F,Q} \left[E_{S,i}(R) - E_{0,i}(R)\right]. \tag{9}$$

This expression retains the native ground surface and adds corrected excitation gaps. The subtraction is geometry dependent, and its first and second derivatives are included explicitly. It is not an empirical constant shift.

In the derivative-qualified campaign, frozen projected operators are transported with local atomic covalent frames. The covalent graph is established at the anchor using the sum of covalent radii plus 0.28 Å; neighborhoods are extended until a noncollinear frame can be defined, and the graph is then held fixed. Local rotations of the orbital shells are obtained from Kabsch fits. Their exact first and second nuclear derivatives enter the operator jets, together with overlap and Pulay terms. This procedure defines a differentiable frozen-parameter local surface. It does not recalibrate the DFT model at every nuclear displacement, and transfer far from the reference geometry requires separate assessment. The reduced extension retains co-rotating AO-pair operators and currently qualifies vertical energies only; it must not inherit nuclear-derivative validation from a different reference/operator generation.

A geometry-only generalized core eigensystem supplies the working orbital chart. It is not identified with a physical ground-state solution. Excited-state energy evaluations therefore do not require an intervening ground-state SCC cycle. The three physical restricted references in Eq. (9) are optimized directly and stored separately.

2.5 Analytic nuclear gradient and Hessian

Let κ collect the nonredundant orbital coordinates of both parents, and $J = g_\kappa$. Differentiating Eq. (8) gives the bordered response system

$$\mathcal{K} \begin{pmatrix} \kappa_R \\ \mu_R \end{pmatrix} = -B, \qquad \mathcal{K} = \begin{pmatrix} \mathcal{L}_{\kappa\kappa} & J^{\mathsf{T}} \\ J & 0 \end{pmatrix}, \qquad B = \begin{pmatrix} \mathcal{L}_{\kappa R} \\ g_R \end{pmatrix}. \tag{10}$$

For a regular constrained stationary branch, its analytic gradient and Hessian are

$$\frac{d\overline{E}_S}{dR} = \mathcal{L}_R, \qquad H_S = \mathcal{L}_{RR} - B^{\mathsf{T}} \mathcal{K}^{-1} B. \tag{11}$$

The inverse denotes a solution of the bordered linear system. A minimum in orbital space is not required. Regularity of the constraint Jacobian and resolution of the relevant curvature are required.

A common metric-preserving AO chart makes the relative overlaps independent of explicit nuclear coordinates at fixed orbital coordinates. In that chart the explicit constraint nuclear and mixed jets vanish, but their orbital second derivatives remain in $\mathcal{L}_{\kappa\kappa}$. The implementation includes orbital/nuclear cross derivatives, local-frame jets, metric terms, and product-chart terms needed when an exactly common orbital is eliminated. The complete Hessian of Eq. (9) is

$$H_{\mathrm{exc}} = H_{0,\mathrm{native}} + H_S - \tfrac{1}{2}(H_{0,F} + H_{0,Q}). \tag{12}$$

An arithmetic mean of separately relaxed parent Hessians would omit the coupled constrained response. Finite differences are used only to validate this analytic implementation.

2.6 Assigned-state initialization and batched orbital response

An adiabatic interpolation between two different one-body models is not required to initialize their assigned stationary states. In the revised solver, each parent is initialized independently from the leading singular vectors of the fixed projected reference transition, placed in the occupied-hole and virtual-particle labels. These natural-transition-orbital (NTO) rotations preserve the native closed-shell density. Retained stationary and native seeds provide additional initial guesses. Analytic Newton searches are tried across these seeds before freeze-and-release generalized mode following. The final electronic index is measured from the analytic curvature rather than restricted to a list of guessed low indices. This extends the direct-optimization motivation of Levi and co-workers without changing the energy functional.[13, 15, 16]

Once the requested electronic index is resolved and no curvature is classified as unresolved, a complete Newton step is used when it lies inside the existing trust radius. This avoids artificial linear convergence when a resolved soft curvature is smaller than the regularization floor. Existing state guards and residual-growth checks still apply. The change passes 31 remote regression tests, including a soft-saddle case that previously remained strongly damped.

The orbital Jacobian is evaluated by forming the alpha and beta density perturbations for all active rotations together. Their common charge response is applied once; spin response and full AO-pair contractions are evaluated in batches with matrix products. The scalar response remains available as a fallback. This is an algebraic reorganization of the analytic derivative, with unchanged energy, orbital tolerance, and nuclear-response equations.

2.7 Tracking, certification, and restart

The APOC state-following machinery monitors each parent CSF and its original active-pair character. Accepted steps require overlap and character measures of at least 0.7, positive parent excitation gaps, joint orbital stationarity, and a resolved projected orbital index. Curvatures smaller than $10^{-7}\ E_\mathrm{h}$ in absolute value are treated as unresolved for certification. An orbital-index check complements overlap tracking, since an apparent character match alone does not certify the desired stationary branch.

The accepted restart generation contains both CSFs, constraint targets, multipliers, three ground references, and the original anchor. Model, occupation, promotion, index, and checksum checks accompany initialization and restart. Initial certified parent states are loaded automatically, and subsequent accepted checkpoints take precedence. The LINK interface explicitly selects this composite OO-DFT path; existing default methods are unchanged.

For the reduced validation extension, the assignment policy was amended after examining solver behavior on nitropyridine N-oxide. A fixed overlap of 0.90 with a DFT transition and a separate absolute active-pair anchor were too restrictive under relaxation with a different Hamiltonian. Writing X for the unit-normalized projected reference transition and σ_1 for its leading singular value, the attainable single-pair fidelity is $f = |\langle X, uv^\mathsf{T}\rangle|/\sigma_1$ for normalized occupied and virtual directions u, v. The revised criterion requires $f^2 > 1/2$, occupied and virtual projection norms of at least 0.7, and a projected-overlap margin of at least 0.05 against every available converged same-symmetry reference root. Local full-CSF step overlap remains at least 0.90. The separate absolute active-pair anchor becomes a diagnostic when a fixed reference transition is supplied. The ratio is a single-pair representability diagnostic, not a many-electron state probability. The raw-overlap prototype was discarded before predicting the remaining six states because it also penalizes transitions whose best compact-space single-pair overlap is intrinsically below unity. These are amended identity criteria, not improved excitation-energy

accuracy or relaxed stationarity tolerances. The rule is applied to the entire reduced set; the reference-root coverage remains finite, so assignment does not prove global uniqueness.

3 Computational protocol

All electronic calculations and derivative validations reported here were performed on the verified remote host `oracle.sns.it`. The MOUSE predictions and derivative checks use the archived final relative-active-pair campaign of 30 October 2026. A subsequent publication-validation campaign adds dedicated sTDA-xTB calculations on the identical archived geometries and comparisons with published structural references. The energy model, 50/50 weights, zero additive offset, and exchange-response scaling were fixed for the four prospective calculations. PBE0 ground-state reference calculations use the archived augmented correlation-consistent double-zeta basis, aug-cc-pVDZ, and numerical integration grid level 3 for projected response construction. The state-specific orbital and nuclear calculations subsequently use the semiempirical valence representation.

Vertical reference values are the archived *aug*-cc-pVTZ theoretical best estimates (TBEs) from the QUEST data collection.[20, 21] Reference geometries and state assignments were inherited from the frozen input campaign. The supporting files retain the individual TBE method labels where available. DFT transition information assists state assignment; TBEs do not enter operator construction, root selection, or the final weights. State selection uses orbital/transition identity and stationarity rather than proximity to a target energy.

The reported cohort contains 15 previously exposed regression states in 14 molecules and four additional molecular tests. These latter molecules had earlier native or shell-model predictions, so they are prospective with respect to the present projected-response composite, not a wholly blind benchmark. The 19 reported, qualified transitions form a curated development/validation cohort, not a denominator for a database-wide convergence rate. Eighteen are $n\rightarrow\pi^*$ and one is $\pi\rightarrow\pi^*$. The supporting information gives every vertical energy and signed error.

Figure 3 collects the archived anchor geometries of both molecular cohorts. The reduced extension is shown separately because its operator-selection and assignment protocol differs from the original vertical-validation cohort.

For the reduced extension, benchmark reference-response calculations use checkpointed single-vector Davidson expansion per requested root, response residual tolerance 10^{-7}, and linear-dependence threshold 10^{-18}. The PBE0 Hamiltonian, basis, grid and requested symmetry/ordinal assignments are unchanged. Completed symmetry roots and accepted Ritz vectors are reused with operator fingerprints. These reference calculations assist benchmark assignment and are accounted for as preparation, separately from the semiempirical stationary-state evaluation. Assigned NTOs and identity projections use the retained native ground-state orbital chart, never the geometry-only core eigensystem used internally for transport.

For TDA, the response equation is

$$\sum_{jb} A_{ia,jb}X_{jb} = \omega X_{ia}, \qquad A_{ia,jb} = (\epsilon_a - \epsilon_i)\delta_{ij}\delta_{ab} + K_{ia,jb}. \tag{13}$$

Each control uses the directly optimized restricted ground orbitals of its corrected GFN2 parent, with that parent's one-body operator and the same calibrated response kernel and $\lambda = 0.9$. DFT orbitals are not substituted into this eigenproblem. For a full nonlocal kernel, the real symmetrized static response is not the TDA A matrix. In the present integral convention the missing contribution is

$$A = A_{\mathrm{static}} + \tfrac{1}{2}(0.225)\big[(ib|aj) - (ij|ab)\big]. \tag{14}$$

Independent water/STO-3G comparisons with the PySCF A matrix give agreement within 3.6×10^{-15} E_{h} at exchange fractions 1 and 0.225. The older static-response values and their timing ratios are retained only as identified archival diagnostics. These are matched response




Figure 3: Molecular structures studied in this work, rendered with Jmol from the archived calculation geometries. (A) The 18 molecules of the initial 19-state vertical-validation cohort. (B) The 10 molecules of the reduced 14-state extension, with all 14 assigned vertical targets now completed. Inclusion denotes study scope, not successful validation of every target state. Ball-and-stick views use carbon in grey, hydrogen in white, nitrogen in blue, and oxygen in red. Display transformations are rigid rotations and translations only; individual panel scales vary for legibility.

controls for the present energies, not a benchmark of a separately parameterized published sTDA method.[1] A mean of the two TDA spectra can be a numerical composite control, but is not the unique linear response of the constrained composite energy.

Molecular minima were accepted at a maximum Cartesian gradient below 3×10^{-5} E_h/bohr, together with state and gap checks. Frequencies were obtained from mass-weighted analytic Hessians with translations and rotations removed. Directional Hessian checks used displacements of 2×10^{-4} and 10^{-4} bohr. Complete Hessian-column checks used 10^{-4} bohr. These checks assess derivative implementation, separately from physical frequency accuracy.

4 Results and discussion

4.1 Vertical excitation energies and the $n \to \pi^*$ problem

Table 1: Vertical-energy error statistics in eV; MSE denotes mean signed error, prediction minus TBE.

Subset	N	MAE	MSE	Max. abs.
Exposed regression	15	0.150	-0.094	0.402
Prospective	4	0.304	-0.114	0.535
All states	19	0.183	-0.099	0.535
$n \to \pi^*$ only	18	0.176	-0.121	0.535

The full cohort gives a mean absolute error (MAE) of 0.183 eV and a mean signed error of −0.099 eV (Table 1). The $n \to \pi^*$ subset alone has an MAE of 0.176 eV. The exposed regression subset is more accurate than the prospective subset, emphasizing the need to keep these groups distinct.

Uracil illustrates the original target of the development. Its $n \to \pi^*$ excitation is predicted at 4.885 eV against a TBE of 4.935 eV; the $\pi \to \pi^*$ energy is 5.686 eV against 5.381 eV. The relative constraints preserve the independently stationary 50/50 anchor values rather than improving them by a new offset. The $n \to \pi^*$ state has projected electronic saddle index three, illustrating why a universal first-order electronic saddle assumption would be inappropriate. The single $\pi \to \pi^*$ example, however, is insufficient to establish balanced accuracy across excitation families.

Table 2: Prospective OO-DFT energies (eV). The four states are valence singlets. Earlier static-response controls are withdrawn as TDA comparisons.

Molecule	TBE	OO-DFT
Cyanoformaldehyde	3.817	3.989
Cyclopropenone	4.274	4.482
Imidazole	6.715	6.180
Pyridazine	3.830	3.530
MAE	–	0.304

The four prospective OO-DFT values are retained in Table 2. Their MAE is 0.304 eV. Earlier controls labelled TDA used a symmetrized static nonlocal response and omitted the antisymmetric exchange contribution needed for the TDA A matrix. Their numerical error comparisons have therefore been withdrawn. Corrected controls retain each parent's own corrected GFN2 ground-state orbitals, rather than DFT molecular orbitals. Figure 4 shows the OO-DFT errors independently of this control correction.

The largest absolute error in the whole cohort is 0.535 eV. No state-dependent offset, orbital-extent-dependent fitted λ, or readjustment of the 50/50 weight was introduced to remove it. A universal correction of the $n \to \pi^*$ error is consequently not established. The useful result




Figure 4: (a) Composite semiempirical OO-DFT energies against archived TBEs for all 19 states, distinguishing exposed and prospective examples. The dashed line denotes equality. (b) Signed OO-DFT errors for the four prospective tests. The earlier static-response controls have been removed pending corrected TDA validation; no published sTDA-xTB calculation is represented.

is a physically explicit, stationary semiempirical candidate with a transparent residual error distribution.

4.2 Direct comparison with published sTDA-xTB

Dedicated xtb4stda (release 1.0 with the distributed stda1/stda2 parameter files) supplies the ground-state wavefunction, followed by std2 version 2.0 with `-xtb -e 10 -vectm 30`. Published method parameters are retained without fitting.[3, 4] The archived input XYZs are identical to those used for MOUSE. The recorded contracted Gaussian basis is reconstructed from the xtb4stda wavefunction, checked against its orbital orthonormality, and integrated directly against the fixed DFT reference basis. Each external transition vector is projected into the fixed occupied/virtual reference spaces. Assignments maximize normalized transition overlap, rather than agreement with the TBE. Acceptance requires overlap and projected norm of at least 0.7 and a best-versus-alternative overlap margin of at least 0.05. All 19 requested states satisfy these gates. Executable, parameter, geometry, and transition-vector hashes are retained.

Table 3: Direct comparison with dedicated published sTDA-xTB on identical archived geometries and transition-matched states. Both methods are evaluated against the same TBEs. The cohort is curated and is not the newly frozen blind set.

Subset	N	MAE (eV)		sTDA max. error (eV)
		MOUSE	sTDA-xTB	
All 19 states	19	0.183	0.454	1.506
Exposed regression	15	0.150	0.501	1.506
Previous prospective	4	0.304	0.278	0.562

On the complete curated cohort, the sTDA-xTB MAE is 0.454 eV against 0.183 eV for MOUSE. This average is influenced by sTDA-xTB underestimation for nitroxyl and nitrosomethane; the largest absolute error is 1.506 eV. The result does not establish a general ranking. For the four previous prospective examples, sTDA-xTB has a slightly lower MAE, 0.278 versus 0.304 eV, while the RMSEs are 0.359 and 0.335 eV, respectively. Figure 5 and the complete supporting

table show the distribution. These external spectra are distinct from the matched GFN2-parent TDA controls discussed above.




Figure 5: Direct comparison with published dedicated sTDA-xTB on the same geometries and transition-matched states. (a) Every signed error against the common TBE reference. Unmarked states are $n\pi^*$. (b) MAEs with exposed regression and previous prospective subsets kept separate. These are the original curated states; the newly frozen blind benchmark has no scores in this figure.

An additional read-only audit recovers TD-DFTB2/mio-1-1 spectra for all 19 original states on the same geometries. The DFTB spatial representation uses one s and three Cartesian p functions per heavy atom; group projectors classify the archived MO coefficients without introducing a radial-overlap approximation. The fixed DFT transition density defines the target irrep, and assignment uses its ordinal within that irrep. The resulting MAE is 0.545 eV versus 0.183 eV for MOUSE. Separately, 15 lowest-singlet ΔDFTB energies have the correct final determinant spatial irrep and give 0.314 eV MAE versus 0.172 eV for MOUSE on that same subset. Two incomplete SCC calculations and two higher-reference states remain explicit in Supporting Table S7. These symmetry/ordinal assignments have a weaker identity certificate than the cross-AO transition-overlap-qualified sTDA-xTB comparison: roots with the same irrep may differ in electronic character. They therefore supply a limited external comparison, not an independently validated ranking. For the three eligible previous prospective cases, ΔDFTB and MOUSE have similar MAEs, 0.234 and 0.227 eV. The ΔDFTB multiplet-sum energy does not establish an exactly spin-projected DFT energy.

Table 4: Excited-state minima: maximum Cartesian gradient in E_h/bohr, minimum projected frequency in cm^{-1}, and maximum directional Hessian error at a displacement of 10^{-4} bohr in $E_h/bohr^2$.

Molecule	max $\|g\|$	ν_{min}	HVP error
Acetaldehyde	2.23×10^{-5}	186.9	1.90×10^{-9}
Acetone	2.51×10^{-5}	148.3	3.06×10^{-10}
Formamide	5.49×10^{-6}	324.4	2.02×10^{-9}
Maleimide	1.88×10^{-5}	179.2	1.96×10^{-9}
Nitroxyl	9.14×10^{-6}	1261.4	3.33×10^{-9}

4.3 Molecular minima and analytic second derivatives

Qualified excited-state minima were obtained for nitroxyl, acetaldehyde, acetone, formamide, and maleimide. Their maximum residual gradients are between 5.5×10^{-6} and 2.5×10^{-5} E_h/bohr. All projected vibrational eigenvalues are positive; the lowest frequencies range from 148 to 1261 cm^{-1} (Table 4 and Fig. 6). Each result refers to the state-followed constrained composite surface of Eq. (9). These minima demonstrate that the energy and orbital stationarity equations can be coupled to nuclear optimization.

The Hessian-vector checks at 10^{-4} bohr differ from analytic results by at most 3.4×10^{-9} $E_h/bohr^2$ for the five minima. More demanding all-column checks give maximum matrix errors of 1.15×10^{-8} $E_h/bohr^2$ for the nine Cartesian columns of nitroxyl and 4.67×10^{-9} $E_h/bohr^2$ for the 18 columns of formamide. The corresponding scalar-energy/gradient checks give maximum discrepancies of 3.64×10^{-9} and 2.26×10^{-9} E_h/bohr. This agreement tests the assembled total Hessian, including coupled orbital response and all ground-reference contributions.

Figure 7 summarizes both levels of derivative verification.

Positive curvatures and finite-difference agreement establish local stability and numerical consistency of this model. They do not establish experimental vibrational accuracy. A separate benchmark against excited-state equilibrium structures and harmonic frequencies is needed before assigning a spectroscopic accuracy to these results. Complete final coordinates, frequency lists, and displacement-check data accompany the manuscript.

The CCSDR(3)/def2-TZVPP structural comparison is more demanding than verifying derivatives.[22] For acetaldehyde, acetone, and maleimide, the 11 selected bond lengths differ by -0.026 to $+0.007$ Å. Angular accuracy is less uniform: acetone's carbonyl pyramidalization is underestimated by 11.7°. Every selected internal coordinate and its reference is reported in the supporting information. For nitroxyl, the three harmonic frequencies ordered from low to high differ from the published (T)+EOM/aug-cc-pVTZ values by +271, +134, and +161 cm^{-1}.[23] This is a frequency-order precheck pending normal-mode vector confirmation; vibrationally averaged distances and anharmonic fundamentals are excluded. The discrepancies show that analytic correctness alone does not establish spectroscopic accuracy.

4.4 Computational cost and reuse of the ground-state engine

The previous energy-only comparison used the incomplete static-response operator and is withdrawn as a TDA benchmark. An isolated comparison at the same retained nitropyridine N-oxide checkpoint instead tests the analytic orbital Jacobian before and after batching. For 44 basis functions and 510 rotations, its time decreases from 4.1384 to 0.2416 s on Oracle with two threads, a factor of 17.13. The largest elementwise Jacobian difference is 8.88×10^{-16}. The current regression set passes 26 tests, including scalar/batched response, independent orbital-rotation and relaxed-Hessian checks. These measurements establish a kernel optimization, not an end-to-end speedup over TDA.

With reference preparation cached, the independently assigned nitropyridine $n\pi^*$ parents




Figure 6: The five optimized excited-state molecular minima of the constrained composite model. Ball-and-stick views use the archived coordinates with rigid display rotations; connections are inferred from interatomic distances and do not assign bond orders. Selected N–O or C–O distances are given in Angstrom. Each $\nu_{\min}$ is the lowest frequency after removal of translations and rotations. The panel scale varies between molecules. These are calculated structures and model frequencies, not experimental references.

require 1.01 and 4.56 s, and the coupled state plus three restricted-ground certificates requires 8.10 s. Its final joint residual is 5.22×10^{-15}. The CT parents require 1.28 and 7.90 s, followed by 14.04 s for the coupled state and restricted references; its residual is 2.56×10^{-13}. The electronic indices are six and eight for the respective constrained states. Both pass the amended assignment audit. The available DFT root list contains only the assigned root in each symmetry, so it does not establish global uniqueness. The $n\pi^*$ recovery already passes the earlier 0.90 fidelity criterion; CT uses the amended identity policy. These are vertical solver qualifications. After completion of all 14 predictions and matched TDA controls, the prediction lock was written and reference labels were opened. Nuclear derivatives of this revised reference generation are not certified by vertical solves.

The final six-size benchmark below includes total gradients, analytic Hessians, and process memory. Available reference-preparation stage costs are audited separately; missing cold preparation totals are treated as unknown. These data do not establish an end-to-end advantage over TDA.

The architectural advantage is more general than the timing result. The excited-state implementation uses the same semiempirical energy terms, Fock builds, spin-polarization response, and first/second nuclear derivative machinery as the ground-state engine. The additional




Figure 7: Numerical validation of the analytic nuclear Hessian. (a) Maximum Hessian-vector discrepancies for the five molecular minima at two finite-difference displacements. Some curves nearly coincide. (b) Maximum matrix discrepancies when every Cartesian column is checked for nitroxyl and formamide at $h = 10^{-4}$ bohr. The plotted errors come from the archived remote validation; finite differences are checks and are not used to construct the production Hessian.




Figure 8: Measured acceleration of one analytic orbital Jacobian at the identical retained nitropyridine checkpoint. Batching preserves the numerical operator while reducing repeated scalar charge and spin contractions. Reference preparation and nuclear derivatives are excluded; this is not a timing comparison with TDA.

components are the CSF energy, orbital saddle/stationarity solver, constrained response equations, and state following. A separate TDA response model is not needed to define the OO-DFT energy surface. DFT supplies the frozen reference information; the production stationary surface is evaluated through the semiempirical engine.

4.5 Position relative to simplified response and state-specific alternatives

Table 5: Methodological positioning. Derivative statements refer to the cited formulations, not an exhaustive survey of all software capabilities. No inter-method accuracy ranking is implied.

Family	Electronic reference	Excited-state description	Relation to this work
sTDA/sTD-DFT; Xs variants[1, 2, 4]	Typically DFT orbitals; optional dedicated xTB reference	Simplified multi-root linear response	Spectroscopic reference family; different target from stationary OO energy
sTDA-xTB[3]	Dedicated xTB Hamiltonian and augmented basis	Simplified TDA spectrum	Essential future external control; distinct from our GFN2 TDA controls
TD-DFTB; hybrid extensions[5, 6]	DFTB ground-state orbitals	Response with DFTB-specific interactions	Compact competing route to excited-state properties and dynamics
TDDFT+TB; TDDFT-aas[7, 8]	DFT ground state	Approximate response; analytic gradients	Demonstrate that low-cost analytic excited-state forces are established
ΔDFTB[5, 9]	State-specific DFTB configurations	Relaxed excited-state total energies and gradients	Closest state-specific semiempirical precedent
OMx/ODMx–MRCI[10, 11]	Parameterized semiempirical Hamiltonian	Multireference CI states	Established configurational alternative; broader state mixing
MOUSE OO-DFT (this work)	GFN2 plus frozen projected DFT response	Constrained spin-pure ROKS composite	Analytic gradient and coupled nuclear Hessian; local validation

Table 5 separates three methodological choices: the electronic reference, the definition of the excited state, and the derivative capability established in the cited work. These axes are important because an inexpensive response calculation may still require a full DFT ground state, and an excited-state gradient does not establish the availability of an analytic excited-state Hessian. Analytic gradients are already available in several competing approaches. In a representative TD-DFTB vibronic implementation, the excited-state Hessian was obtained by finite differences of analytic gradients.[24] This is a specific documented comparison, not a claim that analytic second derivatives are absent from every other semiempirical implementation.

Relative to sTDA-xTB and the std2 response family, the distinguishing feature here is optimization of a targeted excited-state energy together with its coupled nuclear response. We do not establish superior multi-root spectroscopy, transition intensities, or large-system throughput. Published sTDA-xTB has its own Hamiltonian, augmented basis, screening, and kernel parameters; it must be assessed separately from the matched GFN2 controls. The separate released sTDA-xTB comparison on identical geometries gives a lower aggregate MAE for MOUSE, but the four previous prospective cases slightly favor sTDA-xTB by MAE. A broader blind comparison is therefore required before a transferable accuracy ranking can be made.

Relative to ΔDFTB, the present formulation uses a different semiempirical foundation and projected charge/spin operators, common-orbital singlet CSFs, a two-parent relative-active-pair stationary problem, and an explicitly implemented analytic nuclear Hessian. The existing ΔDFTB results on structures and vibrational properties are an essential precedent.[9] Relative to OMx/ODMx–MRCI, our target is a compact stationary CSF surface rather than a multireference state expansion. The present data do not establish which approach is preferable near strong

configurational mixing or conical intersections.

The case for the semiempirical level in this implementation is correspondingly specific. The compact valence representation and the reuse of native energy and derivative primitives support repeated optimization and curvature evaluations in one consistent model. DFT information is paid for in a reference preparation step and then reused within the validated local geometry range. A useful accounting is

$$T_{\text{workflow}} = T_{\text{DFT reference}} + T_{\text{projection}} + \sum_{k=1}^{N_{\text{geom}}} T_{\text{OO/derivatives}}(R_k). \tag{15}$$

The first two terms can be amortized over subsequent evaluations, but they do not disappear. The complete AO-pair correction also introduces dense matrices whose storage can grow as the fourth power of the AO dimension; a compact basis alone does not establish linear scaling or negligible memory cost. The present small-molecule tests demonstrate consistency and initial accuracy, not an end-to-end cost advantage over DFT. Future workflow benchmarks must include reference preparation, memory, state initialization, and derivatives, alongside accuracy. This makes the intended use a hierarchy: high-level data for assessment, DFT for reference information, and the semiempirical OO surface for repeated state-specific molecular work.

4.6 Scope, limitations, and next validation targets

The present candidate is local in both parameterization and state identity. Its frozen DFT-informed operators and covalent frames are anchored to a reference structure; bond rearrangements and large departures from that anchor have not been validated. Persistent relative constraints are part of its definition, and the resulting branch need not coincide with an unconstrained adiabatic state of a single functional. State following and saddle certification reduce ambiguity but do not prove global uniqueness.

Likewise, spin purity of the CSF should be separated from the accuracy of the multiplet-sum energy. The initial 19-state sample is dominated by valence $n \to \pi^*$ transitions and contains only one $\pi \to \pi^*$ transition. Charge-transfer, Rydberg and doubly excited states have not been established as transferable target classes for this final candidate; conical intersections and transition-metal systems are untested. Successful applications of direct OO-DFT in those areas by Levi and co-workers motivate future tests; they are not evidence for their successful description by the present semiempirical energy.

The next assessment should freeze the complete model and test a broader independently assigned set, report all attempted states and assignment failures, and compare energy differences, equilibrium structures, and vibrational curvatures against suitable high-level references. Such tests can determine whether the current $n \to \pi^*$ improvements persist across orbital localization and molecular size. They should precede any further empirical adaptation of exchange scaling or weights.

An additional exposure-audited manifest was frozen before new predictions: 44 states in 30 molecules, comprising 12 $n\pi^*$, 12 $\pi\pi^*$, eight charge-transfer, and 12 Rydberg targets. The four-state CT shortfall relative to the intended quota is retained explicitly. Reference numerical labels remain segregated from the prediction inputs. No aggregate score for the full 44-state manifest is reported; the completed reduced subset is assessed below. The operational extension was subsequently reduced to 14 states in ten molecules after inspecting baseline convergence outcomes, at the requested smaller scope. Six original solver-development states, two nitropyridine repair states and six remaining states are accounted for separately. All original 44 dispositions are retained; the reduced set is not a wholly blind sample. All 14 states now satisfy stationarity and available-root assignment checks, including phthalimide after recovery of an archived stationary seed. All 14 corrected TDA assignments were also completed. The prediction lock was written before numerical labels were opened. The complete locked results are reported in Table 6 and

the Supporting Information; this adapted extension is distinct from the original cohort and is not wholly blind. An initial algebraic weight precheck on the original 19 vertical anchors gives MAEs of 0.179, 0.183, and 0.239 eV for full-parent weights 0.25, 0.50, and 0.75. The 50/50 candidate remains fixed; these values do not test off-anchor weighted surfaces or Hessians. The final campaign below completes a common-geometry full-anchor test, fresh prospective cohorts, physical stationary-point comparisons, and a six-size prepared-property benchmark. A full exchange-scaling scan and broader independently assigned external-method comparisons remain outside this finite campaign.

4.7 Completed reduced extension and the multiconfigurational limitation

Table 6: Completed reduced extension: MAE (eV) on the same geometries and assigned states. F and Q denote the two corrected GFN2 parents; TDA mean is their arithmetic spectral average. This adapted set is separate from the original 19-state cohort.

Family	N	OO 50/50	TDA-F	TDA-Q	TDA mean
All	14	0.492	1.013	0.261	0.611
$n\pi^*$	5	0.193	0.592	0.142	0.311
$\pi\pi^*$	5	0.954	1.409	0.417	0.913
CT	3	0.187	0.890	0.244	0.539
Rydberg	1	0.598	1.499	0.129	0.814

The completed 14-state extension gives an MAE of 0.492 eV for the original single-CSF 50/50 model, compared with 0.261 eV for TDA-Q, 1.013 eV for TDA-F, and 0.611 eV for their arithmetic mean. The five $\pi\pi^*$ states account for an OO MAE of 0.954 eV; cyclopentadiene, furan, and phthalimide are overestimated by 1.746, 1.444, and 1.269 eV, respectively. These results do not support a general accuracy advantage over TDA. Assignment success and predictive accuracy must be assessed separately.

The leading squared NTO singular values of these three fixed projected transitions are 0.544, 0.533, and 0.754. A single excited singlet CSF cannot represent the orthogonal transition pairs carrying the remaining weight. As a development test, an orthonormal set of singlet excitation CSFs is combined by diagonalizing a small state-interaction matrix with unchanged ROKS diagonal energies and bare Slater–Condon off-diagonal couplings,

$$H_{ii} = 2E_{M_i} - E_{T_i}, \qquad H_{ij} = 2(h_i p_i|h_j p_j) - (h_i h_j|p_i p_j), \quad i \neq j. \tag{16}$$

The use of a small configuration space with DFT-informed diagonal energies is motivated by minimal-active-space multistate DFT.[25] That approach optimizes determinant orbitals separately and uses nonorthogonal state interaction; the present common-orbital, orthonormal-CSF construction is a different approximation. This defines a new approximate multistate energy, with a spin-pure wavefunction and analytic CI-eigenvector response. The off-diagonal operator is unscaled bare exchange/Coulomb, whereas the diagonal response retains $\lambda = 0.9$; it differs from the TDA kernel. No energy offset or fitted interaction multiplier is introduced. The two parents retain equal weights. The present test chooses the smallest NTO rank containing 95% of the squared Frobenius transition norm, capped at four. All labels are already exposed at this development stage.

On the five $\pi\pi^*$ examples this trial reduces the MAE from 0.954 to 0.162 eV, and all eight newly optimized multi-CSF parent points pass the available-root identity audit. However, the same rank rule introduces three CSFs for the aza-naphthalene $n\pi^*$ state and lowers its prediction from 5.279 to 2.839 eV against 5.346 eV. The 95% prescription therefore cannot be promoted as a transferable correction; possible correlation double counting and the treatment of weak configurations remain unresolved. The Rydberg-control attempts and blocked optimization

Table 7: Exposed-label minimal-CSF development trial. Off-diagonal bare Slater–Condon couplings and a 95% NTO-norm rank rule define a new approximate functional. The $n\pi^*$ control invalidates a universal repair claim.

Molecule	Family	Rank	TBE	Original OO	Trial OO
Adenine	$\pi\pi^*$	2	5.191	5.449	5.191
Cyclopentadiene	$\pi\pi^*$	2	6.451	8.197	6.720
Furan	$\pi\pi^*$	2	6.530	7.974	6.674
Phthalimide	$\pi\pi^*$	2	4.600	5.869	4.946
Aza-naphthalene	$n\pi^*$	3	5.346	5.279	2.839
Bimane_Syn	$\pi\pi^*$	1	3.964	3.911	3.911

paths are retained in the development archive. Replacing the interaction with the native GFN2 screened charge kernel has so far only been examined at fixed orbitals and does not constitute a validated alternative.

An independent HF-limit check contracts the same CSFs with a fermionic Hamiltonian and verifies $S^2 = 0$. Finite rotations test the complete analytic orbital Hessian, including CI response. A frozen co-rotating orthonormal-AO transport defines the experimental nuclear surface explicitly. For the furan F parent, two reoptimized finite-difference columns agree with its analytic nuclear Hessian within 2.2×10^{-8} $E_\mathrm{h}/\mathrm{bohr}^2$. For the two-parent candidate, the individual-cosine constraint Jacobian is singular at the furan anchor; its second-order information cannot be discarded. A different, regular manifold preserves the full active cross-overlap matrix exactly through common metric-orthogonal rotations and independent spectator rotations. At independently stationary anchors its analytic response is a projected coupled Hessian with complete CI and orbital response. A complete 27-column finite-difference check of the furan total excited energy, including native-ground restoration and subtraction of both corrected ground references, agrees within 2.29×10^{-8} $E_\mathrm{h}/\mathrm{bohr}^2$. These calculations reuse all completed ground checkpoints. The full-frame constraint defines a stricter experimental PES than the original individual-cosine model. Off anchor, the implementation includes common/private product-chart terms and mixed spectator-gauge curvature. Two columns of the constrained parent-mean Hessian at a displaced furan geometry also agree within 2.3×10^{-8} $E_\mathrm{h}/\mathrm{bohr}^2$, despite individual parent orbital residuals around 2.3×10^{-4} E_h. The public implementation additionally passes all 33 Cartesian columns for cyclopentadiene within 1.30×10^{-7} $E_\mathrm{h}/\mathrm{bohr}^2$. These derivative checks alone do not establish broad off-anchor accuracy; the final physical-reference comparisons below assess a separate part of that question. Automatic LINK/APOC integration is qualified below.

A second development proposal defines the active rank by the effective number of transition pairs rather than a cumulative norm cutoff. With normalized NTO weights $w_i = \sigma_i^2 / \sum_j \sigma_j^2$, let

$$N_\mathrm{pair} = \frac{1}{\sum_i w_i^2}, \qquad r = \left\lfloor N_\mathrm{pair} + \tfrac{1}{2} \right\rfloor . \tag{17}$$

No excitation energy or state-family label enters this rule. It selects rank two only for cyclopentadiene ($N_\mathrm{pair} = 1.988$), furan (2.011), and phthalimide (1.595); the other 11 states have rank one and reuse the identical original single-CSF functional. Existing stationary parents suffice, with no new energy correction or interaction parameter. The resulting vertical development MAE is 0.228 eV over all 14 states and 0.214 eV over the five $\pi\pi^*$ states. All $n\pi^*$, CT, and Rydberg values remain unchanged. This is a proposed solution to weak-pair overcorrelation in the cumulative-cutoff trial, not an independently validated universal prescription: the choice was motivated after seeing that trial. Phthalimide lies only 0.095 above the nearest half-integer rank boundary, which motivates the actual reference/rank and full-energy anchor tests reported below. Rydberg accuracy remains limited.

The present rank-selection protocol uses the archived DFT reference transition amplitudes. These are reused during the semiempirical surface calculation, but are part of its reference




Figure 9: Effective-rank multi-CSF development candidate on the completed extension. (a) Every state error; (b) family MAEs. The rank criterion changes only three $\pi\pi^*$ predictions and leaves the other 11 values unchanged. Labels were already exposed, and this is not an independent validation. The adverse aza-naphthalene result of the separate 95% rule is retained in Table 7. TDA-Q is computed on the corrected GFN2 ground orbitals.

preparation cost. A stand-alone selection using ground-state information alone has not been demonstrated for this candidate.

4.8 Automatic multi-CSF integration and sensitivity

The public research selector `multicsf_50_50` loads a portable frozen-model directory containing both response operators, projected bare integrals, reference transitions, fixed active rank, initial orbitals and three certified ground references. The native ground-state energy and derivative primitives are reused; no SCC, DFT or TDA solve occurs during the qualified surface evaluation. The DFT/TDA information required by the current preparation protocol remains part of its preparation cost. The ground references at displaced geometries are optimized directly on the restricted orbital manifold.

APOC evaluates the coherent signed sum of cross-CSF determinant overlaps, including excitation-ordering phases, before taking its magnitude. LINK uses this descriptor in direct scans, geometry optimization and capped components. Only accepted points atomically publish a generation containing both orbital sets, CI coefficients, ground references, derivatives and the model fingerprint. Trial points preserve the accepted generation. Restart checks all file hashes, the model contract and electronic stationarity; prepared inputs are checked before optimizer cache access. A complete portable furan model and its accepted result accompany the qualification archive.

The furan end-to-end interface test reproduces the research energy exactly, its gradient within 1.4×10^{-17} E_h/bohr and Hessian within 1.34×10^{-14} E_h/bohr^2. Both parent overlaps at a displaced point exceed 0.99998. Cold evaluation including the analytic Hessian takes 98.99 s on two Oracle threads. A fresh session restores the accepted state and its stored Hessian in 0.613 s; this measures derivative reuse, not the cost of a new Hessian at another geometry. Relocated

model inputs, corrupt-checkpoint rejection, and preservation of accepted history pass. Forty-three core, legacy-interface and coherent-APOC tests plus three new LINK contract tests pass. The complete 27-column nuclear check is given in the Supporting Information.

Table 8: Vertical sensitivity of the multi-CSF candidate to diagonal exact-exchange response. One-body DFT anchor, semilocal response, bare off-diagonal CSF couplings and 50/50 weights remain fixed; no fitting is performed. Energies and maximum changes from $\lambda = 0.9$ are in eV.

Molecule	$\lambda = 0.85$	0.90	0.95	Max. change
Cyclopentadiene	6.783	6.720	6.657	0.063
Furan	6.743	6.674	6.604	0.070
Phthalimide	5.030	4.946	4.861	0.085

The exchange-response diagnostic changes only the diagonal exact-exchange response from 0.25λ, retaining the DFT one-body anchor, semilocal response, bare off-diagonal CSF couplings and 50/50 weights. All nine points satisfy the fixed-transition fidelity and stationary-curvature gates. Changing λ from 0.9 to 0.85 or 0.95 changes the three excitation energies by at most 0.063, 0.070 and 0.085 eV, respectively. No parameter is selected from these results. Algebraic vertical full-parent weights 0.4, 0.5 and 0.6 shift the cyclopentadiene and furan energies by up to 0.166 and 0.151 eV from the fixed mean, and phthalimide by 0.035 eV. This does not validate weighted off-anchor surfaces or Hessians.




Figure 10: Exposed-data sensitivity diagnostics. (a) Diagonal exchange-response scaling, with all other operators fixed. (b) Algebraic vertical parent weights. (c) Uniform scaling of all subleading singular values of the fixed reference transition before renormalization. The dashed line is the rank-one/rank-two boundary; the dotted line marks the phthalimide crossing. Panel (c) is an artificial reference perturbation, not a DFT calculation at a displaced anchor.

Rank sensitivity is more consequential. Reducing all subleading phthalimide NTO singular values by 9.75% lowers its effective pair number through 1.5. The corresponding rank-one and rank-two vertical predictions differ by 0.923 eV. Freezing rank at preparation protects the differentiability of each individual PES, but does not establish insensitivity to the chosen reference. The actual reference/rank recalculations below test one local geometry direction. Complete F/Q energy recalibration at displaced anchors and fresh independent validation are therefore included in the final campaign below; their outcomes constrain the transferability claim.

4.9 Second-molecule complete nuclear-response qualification

Figure 11 includes every Cartesian column for furan (27) and cyclopentadiene (33), with maximum Hessian differences 2.29×10^{-8} and 1.30×10^{-7} $E_\mathrm{h}/\mathrm{bohr}^2$, respectively. The cyclopentadiene test uses the public surface interface, with all signed displacements independently relaxed from the

same accepted anchor, including three ground references. Direct energy differences also reproduce the analytic gradients. Translational Hessian identities have residuals below 2×10^{-15} E_h/bohr^2; rotational covariance, $H(\Omega R) = \Omega g$, holds below 4×10^{-11} E_h/bohr for both stored Hessians. The rotational identity includes the nonzero nuclear gradient at these anchors and is not a claim of zero rotational eigenvalues at a nonminimum.

The cold cyclopentadiene public-surface call, including the full analytic Hessian, takes 250.27 s with two Oracle threads and a peak process RSS of 858 MiB. Model parsing precedes that timer. The 66 independently relaxed displaced energy/gradient calls used by central differences take 465.47 s in total (median 7.05 s per call), 1.86 times the analytic-Hessian cost for this one prepared model; DFT/reference preparation is excluded from both values. The furan 98.99 s figure measures the complete LINK interface call and includes its interface overhead; these two observations do not define a matched size-scaling benchmark. All checkpoints are preserved. These checks validate derivatives of the specified local frozen model, separately from independent excitation energies and physical excited-state frequency accuracy.




Figure 11: Complete numerical verification of the multi-CSF analytic nuclear derivatives. (a) Maximum error in each Hessian column against central differences of independently relaxed analytic gradients; (b) analytic gradient against independent central energy differences. The displacement is 2×10^{-4} bohr, and all native/corrected ground terms are included. Values smaller than 10^{-16} are displayed at that plotting floor. These are derivative checks at fixed model anchors, not comparisons of physical vibrational frequencies.

4.10 Actual displaced-reference rank stability

The fixed-reference singular-value perturbation above is complemented by six actual reference calculations: uniform internal Cartesian scales 0.98 and 1.02 for the three rank-two molecules. PBE0/aug-cc-pVDZ/grid-3 ground states and two singlet A_1 TDA roots are recomputed, with the original geometry, state and parameters retained as the comparison anchor. Each target must remain the first same-symmetry root, pass coherent cross-geometry transition overlap above 0.7 and exceed its available competitor by 0.05. A fresh directly optimized native GFN2 S0 defines the compact occupied/virtual projection. No excitation-energy label or parameter adjustment enters this test.

All six points retain rank two (Figure 12; Supporting Table S9). Phthalimide has $N_{\mathrm{pair}} =$ 1.5924–1.5982, compared with 1.5949 at the original anchor; furan has 2.0071–2.0135 and cyclopentadiene 1.9850–1.9927. The smallest target transition overlap is 0.9962, the smallest available-competitor margin 0.9724, and the smallest compact projection norm 0.8256. The maximum atom displacement is 0.069 Å. A separate reprojection recovers the original compact transition to better than 5×10^{-16}. Transporting the original native subspaces with the co-

rotating Löwdin map instead of relaxing S0 gives nearly identical effective ranks; the largest difference is 1.33×10^{-4}.

These observations support local rank stability along this one displacement direction. This rank diagnostic alone does not establish energy insensitivity to the anchor. The final full F/Q recalibrations below evaluate the original common target geometry, separating movement of the target nuclei from changes in reference preparation. This exposed-data reference diagnostic also does not replace a frozen independent molecular validation or remove the discrete-rank boundary identified above.




Figure 12: Actual DFT reference and compact-subspace sensitivity under uniform internal Cartesian scaling. (a) Effective compact NTO pair number with a fresh native S0 (solid circles), compared with transported original occupied/virtual subspaces (dashed). The dotted line is the rank-one/rank-two boundary. (b) Cross-geometry DFT transition overlap (circles) and compact projection norm (squares); the dotted line is the 0.7 acceptance threshold. All six recomputed points pass and retain rank two. No full F/Q energy recalibration or independent accuracy assessment is represented.

4.11 Continuation in soft orbital directions

The full relative-frame parameterization contains common rotations and independent spectator rotations. Nearly redundant combinations can make a unit-normalized energy tangent correspond to much larger matrix-exponential coordinates. We therefore bound the actual common/private rotation coordinates rather than the energy-tangent norm alone. If M maps a quotient tangent step to these coordinates, its SVD construction gives orthogonal columns; define $D_{jj} = \|M_{:j}\|$. For the local residual $g + H\delta$, a safeguarded step solves

$$\min_{\|D\delta\| \le r} \|g + H\delta\|^2. \tag{18}$$

Writing $HD^{-1} = U\Sigma V^{\mathsf{T}}$ gives

$$\delta = -D^{-1} V \operatorname{diag}\left(\frac{\sigma_j}{\sigma_j^2 + \mu}\right) U^{\mathsf{T}} g, \tag{19}$$

where $\mu \ge 0$ enforces the radius. A full Newton step is retained when it fits the radius. Backtracking must decrease the residual norm and preserve the existing fixed-transition fidelity, CI margin, coherent step overlap and full active Gram matrix. The final stationarity and response-curvature certificates are unchanged. A finite-matrix SVD failure is retried with the independent LAPACK `gesvd` driver at the same rank threshold. These are numerical changes to the solver, without changes to the energy or derivative expressions.

For a difficult independent Q parent, we also follow a stationary F parent through

$$v(\tau) = (1-\tau)v_F + \tau v_Q, \qquad 0 \le \tau \le 1. \tag{20}$$

Only the frozen one-body operator changes along this numerical path; the common charge/spin response, bare projected integrals, active rank and final Q Hamiltonian are retained. The initial increment is 1/8 and is subdivided if the state/stationarity gates fail. Accepted orbital checkpoints are reused. The effective exchange remains 0.25×0.9; τ is not an exchange parameter or a physical parent weight. Local electronic curvature is reassessed between bounded mode-following phases, while state identity remains controlled by the same projected transition. The endpoint is used only after certification with the original Q operator. Reference excitation energies do not select the continuation path.

For a geometry path, an explicit opt-in local-CSF policy uses the preceding accepted CI vector only to select the continuous eigenroot. The Hamiltonian, energy, relative Gram constraints, and full-CSF overlap guard are unchanged. The policy is serialized and incompatible restart policies are rejected. Fixed-reference transition following remains the default for vertical validation. Separate contract tests verify both restart behavior and preservation of an explicitly requested CSF rank. Nuclear descent uses bounded BFGS steps and Armijo decrease, retaining the last accepted coordinates and inverse matrix; these numerical changes do not alter the energy model. For the final difficult full-anchor recovery only, a resolved full-Newton trial with raw coordinate norm at most 1 is tested before the 0.3 residual-trust fallback. Acceptance still requires residual decrease, coherent state overlap, the CI margin and exact active-Gram preservation. This documented solver-path amendment closes the remaining phthalimide calibration without altering its Hamiltonian or selecting an excitation-energy error.

4.12 Contracted analytic nuclear response

The complete analytic response initially materialized first and second nuclear jets of every transported AO-pair operator. This is unnecessary for either the charge density $P = P_\alpha + P_\beta$ or the spin density $P = P_\alpha - P_\beta$. With real symmetric P and S, write

$$q_a = A_a^0 : Q, \qquad Q = U^{\mathsf{T}} \frac{SP + PS}{2} U, \qquad q_{a,R} = A_a^0 : Q_R. \tag{21}$$

The field $v = K(q - q_0) + v_0$ is contracted into $A_v^0 = \sum_a v_a A_a^0$ before differentiating the second-order frame contribution. Consequently,

$$E_{RR'} = q_R^{\mathsf{T}} K q_{R'} + \sum_a v_a q_{a,RR'} \tag{22}$$

can be evaluated with small density-like and rotation jets, retaining every overlap/frame mixed term. The same adjoint contraction applies to global and covalent local frames. This removes coordinate-by-feature-by-AO-pair intermediates without changing the functional, its rank or a convergence threshold. Dense frozen response storage is still required.

Charge and spin response are also applied to batches of density perturbations. The multi-CSF orbital/CI Hessian is already assembled independently; nuclear block construction therefore reuses it and computes the complete mixed and nuclear blocks without rebuilding an unused orbital block. The coupled orbital/CI response and all three ground-reference derivatives remain present. Literal operator-jet comparisons, fixed-density finite differences, whole-matrix molecular comparisons and relaxed-gradient differences qualify the reorganization. The previously censored phthalimide calculation is completed analytically; finite differences are used only for validation.

4.13 Final frozen validation and full-anchor tests

The final assessment freezes $\lambda = 0.9$ in the exchange response and the equal F/Q weights. It adds no excitation-energy offset and does not scale DFT orbital gaps by λ. Six complete

reference recalibrations, including F/Q operators and projected integrals, use anchors dilated by ±2%; every excitation is evaluated at the same original target geometry. All six converge with the state and curvature certificates retained. Changes are $-0.0696/+0.0225$ eV for furan, $-0.0113/+0.0080$ eV for cyclopentadiene, and $+0.2350/+0.1966$ eV for phthalimide. Thus local rank stability does not imply energy invariance with respect to the anchor.

The primary molecular holdout contains six states in five molecules from the archived QUEST reference generation, frozen before their predictions and scored only after a prediction lock.[20, 21] Four states satisfy the fixed representation and identity rules: their MAE is 0.6746 eV and RMSE 0.9691 eV. Both Criegee states carry partial-double-character flags in the source; they remain in the denominator. Cyclobutadiene and ammonia are outside the prescribed compact-space projection coverage. This small cohort has no CT target and does not establish transferability to that class. On the three states with qualified Q-parent TDA assignments, the OO and TDA-Q MAEs are 0.8612 and 0.9669 eV. The benzoxadiazole TDA target is unassigned; no energy-proximity replacement is made.

A subsequent new-molecule challenge contains two furofuran states and gives MAEs of 0.5370 eV for OO and 0.3616 eV for TDA-Q. Three further benzoxadiazole states, sharing a primary-cohort molecule, give 0.3998 and 0.2249 eV, respectively. One source reference is marked unsafe. These are separately frozen challenges, not five additional independent molecules or a pooled blind score. All matched TDA matrices use their own stationary corrected GFN2 orbitals and the complete singlet response, including the nonlocal exchange correction. A matrix-level certificate reproduces the previously independently checked furan operator to machine precision.




Figure 13: Final common-geometry anchor test and separately identified prospective cohorts. Crosses indicate inadequate compact-space coverage, not zero error. Both-parent operators are fully recalibrated in panel (a). Signed errors in panel (b) retain partial-double-character and unsafe-reference flags in the accompanying table.

4.14 Physical curvature and symmetry-breaking descent

For furan, the symmetric valence A1 stationary structure is compared with the corresponding EOM-CCSD stationary point of Gromov et al.[26] Five bond lengths give a mean absolute difference of 0.0237 Å, and five angles give 1.55°. Fifteen in-plane frequencies, matched by symmetry then frequency order, give 188.6 cm^{-1} MAE; a 1474 cm^{-1} B2 discrepancy is retained. Cross-method mode-vector overlaps are unavailable. The model has five negative out-of-plane modes, consistent with treating this structure as a stationary saddle rather than a minimum; the reference also identifies out-of-plane instability. Displaced-gradient checks along the most negative and highest positive modes verify the analytic curvatures. A further bounded nonsymmetric

descent preserves lower-energy accepted checkpoints but does not certify a furan minimum.

The analogous cyclopentadiene descent does reach a minimum of the frozen upper A1 surface: the maximum gradient is 2.59×10^{-5} E_h/bohr and all 27 vibrational frequencies are positive, from 204.8 to 2961.2 cm^{-1}. Its total energy is 0.01804 E_h below the symmetric stationary point. A separate central difference of total gradients along its softest vibrational mode agrees with the analytic Hessian action within 1.33×10^{-10} E_h/bohr^2. This is a model minimum, not a correlated-reference validation of that upper-state minimum. Local coherent CSF following is used only for continuous geometry paths; vertical target assignment retains the frozen reference transition. A distorted furan point independently verifies the total energy gradient within 2.46×10^{-9} E_h/bohr.




Figure 14: Furan comparison at the symmetric A1 valence stationary point. EOM-CCSD data are from Tables V and VI of Ref. 26. In-plane mode correspondence uses symmetry and frequency ordering. Imaginary model modes are shown explicitly; no unpublished reference frequencies are assigned to them.

The same analytic implementation now completes the full 48×48 phthalimide Hessian. Batched response and orbital-block reuse reproduce the first complete calculation, which still rebuilt the duplicate orbital block, within 1.32×10^{-12} E_h/bohr^2. A controlled two-step relaxed-gradient difference has quadratic step convergence and agrees after extrapolation within 4.42×10^{-7} E_h/bohr^2. Translation and nonstationary rotation identities also pass. No finite-difference Hessian enters production.

4.15 Final cost and preparation accounting

Six sizes are measured on the dual-socket AMD EPYC 7543 Oracle node (128 logical CPUs) with two threads and an exclusive campaign compute lock. Fresh processes time the prepared energy, energy plus total gradient, energy plus gradient and analytic Hessian, and both-parent all-root singlet TDA on corrected GFN2 orbitals. Three repeats are retained for nitroxyl, acetaldehyde and acetone; a closing budget fixed before larger-case measurements uses one for furan, cyclopentadiene and phthalimide. Model loading and converged-ground setup are outside the property timer; peak RSS covers the whole process. The first three cases use one CSF per parent and the last three two, so this is a practical size/rank benchmark. After exact nuclear-response acceleration, every affected gradient and Hessian timing is remeasured. Unaffected energy-only and TDA measurements are retained, with their execution hashes. External server load and single-repeat large-case variability limit fine timing comparisons.

Across all six measured sizes, the retained prepared OO energy costs more than the complete both-parent TDA evaluation: 0.286–11.478 s versus 0.014–8.731 s. Thus no energy-only speed advantage is observed, even though TDA produces more roots. The intended contribution is the stationary energy surface and its complete analytic nuclear response.

With the contracted response, furan takes 0.851 s for energy plus gradient and 5.104 s including its analytic Hessian; cyclopentadiene takes 1.042 and 7.083 s. Their previous Hessian times were 104.156 and 251.653 s, giving measured reductions by factors of 20.4 and 35.5. Phthalimide now completes energy, gradient and the full 48×48 analytic Hessian in 42.224 s, compared with the previous right-censored time of at least 600 s: a reduction exceeding 14.2. Its energy-plus-gradient call takes 9.292 s. The maximum whole-process RSS in the updated benchmark is 1.07 GiB; the earlier phthalimide gradient process reached 9.26 GiB. All six Hessians are completed. The 72-row ledger contains 48 measured completed evaluations and 24 prespecified omitted repeats; earlier measurements and the censored attempt remain archived.




Figure 15: Prepared-property wall time and whole-process peak RSS after analytic nuclear-response acceleration. Unaffected energy-only and TDA measurements are retained; gradients and Hessians are remeasured. TDA obtains all roots for both parents, whereas OO targets one state. The table records repetitions and ranks. Electronic work from this campaign does not overlap timings; external Oracle load is uncontrolled.

Preparation is audited rather than assigned zero cost. For the fresh molecular cohort, DFT ground calculations take 0.51–38.13 s, recorded reference-TDA stages 0.88–73.36 s per symmetry, and executed full-response calibrations 1.52–77.17 s. Reference TDA supplies state identity and the active-rank rule: the current preparation therefore requires excited-reference transitions as well as ground-state DFT energies. These stage measurements are from a different cohort and are not missing preparation totals for the six timing molecules. Historical complete cold-preparation totals are unavailable. Consequently, the prepared costs quantify the derivative workload but do not establish an end-to-end advantage over TDA or a fully semiempirical reference preparation.

5 Conclusions

MOUSE now supports a semiempirical OO-DFT route in which a spin-pure singlet energy is optimized directly in orbital space and differentiated analytically through second order in nuclear coordinates. The implementation combines two fixed DFT-response-informed GFN2 energy models with equal weights and relative active-pair constraints, and reuses the ground-state engine through a new stationary-orbital and constrained-response layer. It therefore provides an energy-based alternative to the TDA route for state-specific molecular surfaces.

The original 19-state MAE is 0.183 eV; the separate exposed 14-state development candidate gives 0.228 eV. The primary frozen prospective cohort accepts four of six states and gives 0.675 eV, including both source-flagged partial-double-character states. Two states are outside compact-space representability. Subsequent molecular and higher-state challenges give 0.537 and 0.400 eV, respectively, with distinct exposure histories. These results establish an implemented candidate,

not a universal accuracy improvement over GFN2-orbital TDA. Full F/Q anchor recalibrations at six points are complete, with common-geometry excitation changes up to 0.235 eV. Complete furan and cyclopentadiene Cartesian Hessian checks and independent displaced-gradient checks support analytic correctness. A symmetry-breaking cyclopentadiene descent yields 27 positive vibrational modes; the symmetric furan stationary point has a physically relevant out-of-plane instability and a direct EOM-CCSD comparison. The methodological contribution is a compact spin-adapted stationary energy surface, constrained analytic nuclear response, and reusable ground-state derivative machinery. The remaining transferability errors require improvement of the energy model, rather than a numerical offset or a claim that electronic convergence alone solves accuracy.

Data and software availability

MOUSE source code is hosted at https://github.com/yogibubu/MOUSE. The original implementation is committed as `5620e2d`, with trusted-Newton acceleration in `49e0c9d`; calculation-generation source hashes are retained separately in the accompanying protocols. The manuscript package includes machine-readable result tables, reference provenance, the validation summary and source-hash protocol, figure-generation code, and supporting information. The underlying qualification archive is `qualification/roks_relative_pair_20261030` in the associated working tree. Its transfer audit verifies 561 result files and 30 source files; the remote qualification suite reports 86 passing tests. The automatic cold-start/restart example reproduces its energy within 1.8×10^{-15} E_h. The multi-CSF development implementation is committed as `2068ee8`; its verified transfer and qualification logs are in `1eea614`. The separate `qualification/pi_pi_repair_20261031` archive records 141 transferred source/result files, 21 matching local/remote implementation files, upstream model hashes, all blocked development attempts and the effective-rank scoring protocol. The public multi-CSF integration and its full furan Hessian/sensitivity qualification are committed as `489e899`, with 110 further transferred artifacts and 12 matching local/remote modified implementation files in `qualification/multicsf_integration_20261031`. Large phthalimide tensors remain on Oracle with recorded hashes; a complete portable furan model is included. The final numerical solver and restart contracts are committed as `34965d6`; the exact nuclear-response contraction that completes the phthalimide Hessian is committed as `4fe3cd8`. Final validation evidence, prediction locks, preparation audit and transfer hashes are retained in `qualification/final_validation_20261031`; 46 final core/LINK/APOC regressions, two restart/rank contracts and two exact-response tests pass. A permanent public dataset identifier has not been assigned to this working draft.

References

[1] Stefan Grimme. A simplified Tamm-Dancoff density functional approach for the electronic excitation spectra of very large molecules. *The Journal of Chemical Physics*, 138:244104, 2013. doi: 10.1063/1.4811331.

[2] Christoph Bannwarth and Stefan Grimme. A simplified time-dependent density functional theory approach for electronic ultraviolet and circular dichroism spectra of very large molecules. *Computational and Theoretical Chemistry*, 1040-1041:45–53, 2014. doi: 10.1016/j.comptc.2014.02.023.

[3] Stefan Grimme and Christoph Bannwarth. Ultra-fast computation of electronic spectra for large systems by tight-binding based simplified Tamm-Dancoff approximation (sTDA-xTB). *The Journal of Chemical Physics*, 145:054103, 2016. doi: 10.1063/1.4959605.

[4] Pierre Beaujean, Axel Liger, Arya Satisha, Marilù G. Maraldi, Christoph Bannwarth, and Marc de Wergifosse. The std2 package: A computationally efficient program to calculate molecular response properties and excited states using simplified quantum chemistry methods. *The Journal of Chemical Physics*, 165:152501, 2026. doi: 10.1063/5.0349975.

[5] B. Hourahine, M. Berdakin, J. A. Bich, F. P. Bonafé, C. Camacho, Q. Cui, M. Y. Deshaye, G. Díaz Mirón, S. Ehlert, M. Elstner, T. Frauenheim, N. Goldman, R. A. González León, T. van der Heide, S. Irle, T. Kowalczyk, T. Kubař, I. S. Lee, C. R. Lien-Medrano, A. Maryewski, T. Melson, S. K. Min, T. Niehaus, A. M. N. Niklasson, A. Pecchia, K. Reuter, C. G. Sánchez, C. Scheurer, M. A. Sentef, P. V. Stishenko, V. Q. Vuong, and B. Aradi. Recent Developments in DFTB+, a Software Package for Efficient Atomistic Quantum Mechanical Simulations. *The Journal of Physical Chemistry A*, 129:5373–5390, 2025. doi: 10.1021/acs.jpca.5c01146.

[6] Wenbo Sun, Tammo van der Heide, Van-Quan Vuong, Thomas Frauenheim, Michael A. Sentef, Bálint Aradi, and Carlos R. Lien-Medrano. Hybrid Functional DFTB Parametrizations for Modeling Organic Photovoltaic Systems. *Journal of Chemical Theory and Computation*, 21:5103–5117, 2025. doi: 10.1021/acs.jctc.5c00232.

[7] Shana Havenridge, Robert Rüger, and Christine M. Aikens. Analytical excited state gradients for time-dependent density functional theory plus tight binding (TDDFT + TB). *The Journal of Chemical Physics*, 158:224103, 2023. doi: 10.1063/5.0142240.

[8] Yuchen Wang, Shana Havenridge, and Christine M. Aikens. Implementation of energy and gradient for the TDDFT-approximate auxiliary function (aas) method. *The Journal of Chemical Physics*, 161:024101, 2024. doi: 10.1063/5.0213587.

[9] Tim Kowalczyk, Khoa Le, and Stephan Irle. Self-Consistent Optimization of Excited States within Density-Functional Tight-Binding. *Journal of Chemical Theory and Computation*, 12:313–323, 2016. doi: 10.1021/acs.jctc.5b00734.

[10] Deniz Tuna, You Lu, Axel Koslowski, and Walter Thiel. Semiempirical Quantum-Chemical Orthogonalization-Corrected Methods: Benchmarks of Electronically Excited States. *Journal of Chemical Theory and Computation*, 12:4400–4422, 2016. doi: 10.1021/acs.jctc.6b00403.

[11] Pavlo O. Dral, Xin Wu, and Walter Thiel. Semiempirical Quantum-Chemical Methods with Orthogonalization and Dispersion Corrections. *Journal of Chemical Theory and Computation*, 15:1743–1760, 2019. doi: 10.1021/acs.jctc.8b01265.

[12] Gianluca Levi, Aleksei V. Ivanov, and Hannes Jónsson. Variational calculations of excited states via direct optimization of the orbitals in DFT. *Faraday Discussions*, 224:448–466, 2020. doi: 10.1039/D0FD00064G.

[13] Gianluca Levi, Aleksei V. Ivanov, and Hannes Jónsson. Variational Density Functional Calculations of Excited States via Direct Optimization. *Journal of Chemical Theory and Computation*, 16:6968–6982, 2020. doi: 10.1021/acs.jctc.0c00597.

[14] Aleksei V. Ivanov, Gianluca Levi, Elvar Ö. Jónsson, and Hannes Jónsson. Method for Calculating Excited Electronic States Using Density Functionals and Direct Orbital Optimization with Real Space Grid or Plane-Wave Basis Set. *Journal of Chemical Theory and Computation*, 17:5034–5049, 2021. doi: 10.1021/acs.jctc.1c00157.

[15] Yorick L. A. Schmerwitz, Gianluca Levi, and Hannes Jónsson. Calculations of Excited Electronic States by Converging on Saddle Points Using Generalized Mode Following. *Journal of Chemical Theory and Computation*, 19:3634–3651, 2023. doi: 10.1021/acs.jctc.3c00178.

[16] Yorick L. A. Schmerwitz, Elli Selenius, and Gianluca Levi. Freeze-and-Release Direct Optimization Method for Variational Calculations of Excited Electronic States. *Journal of Chemical Theory and Computation*, 22:3571–3584, 2026. doi: 10.1021/acs.jctc.5c01974.

[17] Christoph Bannwarth, Sebastian Ehlert, and Stefan Grimme. GFN2-xTB: An Accurate and Broadly Parametrized Self-Consistent Tight-Binding Quantum Chemical Method with Multipole Electrostatics and Density-Dependent Dispersion Contributions. *Journal of Chemical Theory and Computation*, 15:1652–1671, 2019. doi: 10.1021/acs.jctc.8b01176.

[18] Irmgard Frank, Jürg Hutter, Dominik Marx, and Michele Parrinello. Molecular dynamics in low-spin excited states. *The Journal of Chemical Physics*, 108:4060–4069, 1998. doi: 10.1063/1.475804.

[19] Carlo Adamo and Vincenzo Barone. Toward reliable density functional methods without adjustable parameters: The PBE0 model. *The Journal of Chemical Physics*, 110:6158–6170, 1999. doi: 10.1063/1.478522.

[20] Mickaël Véril, Anthony Scemama, Michel Caffarel, Filippo Lipparini, Martial Boggio-Pasqua, Denis Jacquemin, and Pierre-François Loos. QUESTDB: A database of highly accurate excitation energies for the electronic structure community. *WIREs Computational Molecular Science*, 11:e1517, 2021. doi: 10.1002/wcms.1517.

[21] Pierre-François Loos, Martial Boggio-Pasqua, Aymeric Blondel, Filippo Lipparini, and Denis Jacquemin. QUEST Database of Highly-Accurate Excitation Energies. *Journal of Chemical Theory and Computation*, 21:8010–8033, 2025. doi: 10.1021/acs.jctc.5c00975.

[22] Šimon Budzák, Giovanni Scalmani, and Denis Jacquemin. Accurate Excited-State Geometries: A CASPT2 and Coupled-Cluster Reference Database for Small Molecules. *Journal of Chemical Theory and Computation*, 13:6237–6252, 2017. doi: 10.1021/acs.jctc.7b00921.

[23] Megan C. Davis and Ryan C. Fortenberry. (T)+EOM Quartic Force Fields for Theoretical Vibrational Spectroscopy of Electronically Excited States. *Journal of Chemical Theory and Computation*, 17:4374–4382, 2021. doi: 10.1021/acs.jctc.1c00307.

[24] Robert Rüger, Thomas Niehaus, Erik van Lenthe, Thomas Heine, and Lucas Visscher. Vibrationally resolved UV/Vis spectroscopy with time-dependent density functional based tight binding. *The Journal of Chemical Physics*, 145:184102, 2016. doi: 10.1063/1.4966918.

[25] Ruoqi Zhao, Christian P. Hettich, Xin Chen, and Jiali Gao. Minimal-active-space multistate density functional theory for excitation energy involving local and charge transfer states. *npj Computational Materials*, 7:148, 2021. doi: 10.1038/s41524-021-00624-3.

[26] Evgeniy V. Gromov, Alexander B. Trofimov, Natalia M. Vitkovskaya, Jochen Schirmer, and Horst Köppel. Theoretical study of the low-lying excited singlet states of furan. *The Journal of Chemical Physics*, 119:737, 2003. doi: 10.1063/1.1578051.