

Cartesian Generalized-Internal-Coordinate Refinement of Semiexperimental Equilibrium Structures

Vincenzo Barone *

INSTM, via G. Giusti 9, 50121 Firenze, Italy

E-mail: vincebarone52@gmail.com

Abstract

Semiexperimental equilibrium structures combine ground-state rotational constants with computed vibrational and electronic corrections, yielding structural references that are directly comparable with equilibrium quantum-chemical geometries. Existing refinement workflows demonstrated the power of nonlinear least-squares fits of moments of inertia, predicate observations, and graphical control, but still required substantial care in the construction and maintenance of the structural parameter set. Here we introduce a Cartesian generalized-internal-coordinate (GIC) formulation implemented in Merlino4. The input geometry is Cartesian, isotopologue data are provided as structured TOML, JSON, or CSV records, and the fitted variables are generated automatically, reduced to a non-redundant set, and symmetrized after automatic point-group identification. Ring coordinates are built from endocyclic torsional coordinates and the analogous valence-angle combinations, not from Cremer-Pople variables. The solver fits moments of inertia by default, supports vibrational and electronic corrections, propagates experimental uncertainties to GIC parameters, includes

soft quantum-mechanical predicates, and allows chemically meaningful classes of parameters to be refined together or kept fixed. A Kraitchman-coordinate diagnostic is generated for single-substitution isotopologues. The resulting workflow preserves the spectroscopic logic of molecular-structure refinement while removing user-dependent coordinate construction and making the entire procedure reproducible from both command line and graphical interfaces.

1 Introduction

Accurate equilibrium geometries are central to rotational spectroscopy, force-field validation, and the assessment of electronic-structure methods. Rotational spectra provide highly precise ground-state constants, but the associated structures are vibrationally averaged. The semiexperimental strategy addresses this limitation by correcting experimental constants for vibrational and, when needed, electronic effects before fitting an equilibrium geometry.

The MSR program represented an important step in this direction by combining non-linear least-squares refinement of moments of inertia, advanced error analysis, predicate observations, automatic generation of symmetry-adapted internal coordinates, and a graphical interface.¹ That design established several principles that remain essential: the fit should be performed in a stable observable space, missing experimental information can be complemented by weighted predicates, and uncertainty propagation must be part of the standard output rather than a post-processing option.

Merlino4 keeps these principles but changes the coordinate layer. The molecular structure is read as Cartesian coordinates, while the variables optimized in the least-squares problem are generalized internal coordinates generated automatically from topology, rings, and symmetry information. As in other modern coordinate generators, the primitive coordinate list is built without manual intervention; the distinctive step is the subsequent reduction to a non-redundant active set and the symmetry adaptation obtained after automatic point-group identification. This provides a common coordinate layer for Gaussian input generation,

Wilson GF analysis, VPT2/VCI workflows, ring-puckering paths, and semiexperimental refinement.

The present article describes the new semiexperimental structure solver and its integration into the broader GIC infrastructure. Particular attention is paid to four extensions motivated by practical spectroscopy: grouped refinement of parameter classes such as CH stretches, blocking of poorly determined hydrogen-angle classes, explicit comparison with Kraitchman substitution coordinates, and complete graphical access to the same options available from the command line.

2 Theory and Method

2.1 Semiexperimental Observables

For each isotopologue s , the experimental ground-state rotational constants are converted to equilibrium targets by subtracting computed corrections,

$$B_{\alpha,e}^{(s)} = B_{\alpha,0}^{(s)} - \Delta B_{\alpha,\text{vib}}^{(s)} - \Delta B_{\alpha,\text{elec}}^{(s)}, \quad \alpha \in \{a, b, c\}. \quad (1)$$

Merlino fits principal moments of inertia by default,

$$I_{\alpha}^{(s)} = \frac{K}{B_{\alpha,e}^{(s)}}, \quad (2)$$

where K is the standard rotational conversion constant. Fitting moments avoids the reciprocal amplification that can occur when optimizing directly in rotational-constant space. Direct fits of rotational constants remain available for diagnostic comparisons, with automatic selection of the best-conditioned constant pair for planar molecules.

2.2 Automatic Non-Redundant and Symmetry-Adapted GIC Parameterization

Starting from the parent Cartesian geometry, Merlino constructs a molecular topology and a primitive internal-coordinate list automatically. This first construction step is deliberately similar in spirit to other black-box internal-coordinate generators: bonds, valence angles, linear bends, torsions, out-of-plane terms, and ring-local primitives are generated from the molecular graph. Merlino then performs two additional operations before the coordinates enter the least-squares model. First, the primitive set is reduced to a non-redundant GIC transformation,

$$\mathbf{q} = \mathbf{U}^T \mathbf{s}, \quad (3)$$

where $\mathbf{s}$ is the vector of primitive coordinates. Second, the point group is identified from the Cartesian structure and used to symmetrize the non-redundant GICs without mixing different coordinate types. Stretches, bends, torsions, ring-valence combinations, and other coordinate families are therefore reduced and symmetrized within their own homogeneous blocks.

For rings, Merlino does not use Cremer-Pople coordinates in the GIC layer. Ring puckering and ring deformation are represented by combinations of endocyclic dihedral angles and by analogous combinations of endocyclic valence angles. This choice keeps the variables in the same generalized-internal-coordinate algebra as the rest of the molecule and avoids switching to a separate ring-specific coordinate convention during structure refinement.

The same GIC machinery is used by the Merlino GF and Gaussian-input workflows, ensuring that structural refinement and vibrational analysis operate on consistent coordinates.

The Cartesian displacement associated with a GIC correction is obtained from the analytic Wilson matrix,

$$\Delta \mathbf{x} = \mathbf{B}_q^+ \Delta \mathbf{q}, \quad \mathbf{B}_q = \mathbf{U}^T \mathbf{B}_s, \quad (4)$$

with the pseudoinverse evaluated under rank control. Trial steps are accepted only if they

lower the weighted least-squares objective.

2.3 Least-Squares Model

The residual vector contains selected moments or rotational constants for all isotopologues and optional predicate observations,

$$\mathbf{r} = \mathbf{y}_{\text{obs}} - \mathbf{y}_{\text{calc}}(\mathbf{q}). \quad (5)$$

The weighted Levenberg-Marquardt step is

$$(\mathbf{J}^T \mathbf{W} \mathbf{J} + \lambda \mathbf{I}) \Delta \mathbf{p} = \mathbf{J}^T \mathbf{W} \mathbf{r}. \quad (6)$$

Experimental uncertainties in MHz are propagated to moments when the moment target is used. The final covariance is obtained from the inverse weighted normal matrix scaled by the reduced chi-square, and the Gauss-Newton Hessian is diagonalized to classify the stationary point.

2.4 Predicate Observations and Parameter Classes

Quantum-mechanical estimates of individual GIC parameters can be introduced as soft predicate observations with user-defined uncertainties. This follows the MSR philosophy while keeping all predicates in the generated GIC label space.

Merlino4 additionally supports parameter classes. A class can be refined as a single shared variable,

$$\Delta q_i = \Delta p_c \quad i \in c, \quad (7)$$

or held fixed. This is useful when deuterated isotopologues are unavailable for all hydrogen sites. Typical examples are grouped CH stretches and blocked XYH or XH₂ angle families. The Jacobian is compressed by a class transformation matrix before solving the normal

equations, and uncertainties are expanded back to all members of the class in the reported parameter table.

2.5 Kraitchman Diagnostic

For isotopologues containing a single substitution, Merlino computes substitution coordinates from the Kraitchman equations and compares their absolute values with the fitted coordinates in the parent principal-axis frame. This diagnostic is intentionally separate from the fit: Kraitchman coordinates provide valuable assignment checks, but they do not retain coordinate signs and do not exploit the full correlated least-squares model.

3 Implementation

The solver is implemented in Python and mirrored by independent Fortran77 numerical kernels for analytic **B** rows, rotational constants, weighted normal equations, and parameter-class compression. Structured isotopologue data can be provided in TOML, JSON, or CSV. The graphical dashboard exposes the same controls as the command-line interface: parent Cartesian file, isotopologue data, backend request, observable, rotational-component policy, fixed parameters, predicate observations, and parameter classes. Jobs launched from the GUI are asynchronous and write a manifest with checksums and numerical diagnostics.

4 Differences from MSR

The new implementation is directly inspired by the semiexperimental strategy and predicate-observation formalism used in MSR,¹ but differs in several operational and methodological aspects.

Table 1: Conceptual comparison between the MSR workflow and the Merlino4 GIC workflow.

Feature	MSR	Merlino4
Structural input	user molecular model	Cartesian parent XYZ
Fit coordinates	internal/symmetry coordinates	automatic GICs, reduced to non-redundant symmetry-adapted blocks
Observable target	moments of inertia	moments by default; constants optional
Predicate observations	yes	yes, in GIC label space
Parameter classes	limited/manual	shared or fixed GIC classes
Kraitchman check	external/manual	generated diagnostic CSV
GUI control	yes	same options as CLI, asynchronous jobs
Ring coordinates	internal coordinate dependent	endocyclic dihedral and valence-angle combinations, not Cremer-Pople variables
Cross-workflow reuse	structure refinement	GIC, GF/PED, VPT2/VCI, DVR interfaces

5 Planned Validation

The first validation set will include the molecules used in the MSR article where the required rotational data and vibrational corrections are available, followed by ring and fused-ring cases that stress automatic GIC generation, non-redundant pruning, symmetry adaptation, and ring coordinates built from endocyclic torsions and valence angles. For each molecule we will report residuals in moments and rotational constants, fitted GIC parameters with propagated uncertainties, correlation matrices, stationary-point diagnostics, and Kraitchman comparisons for all single-substitution isotopologues.

6 Conclusions

Merlino4 provides a semiexperimental refinement workflow in which Cartesian input, automatic GIC construction, non-redundant reduction, automatic symmetry adaptation, analytic **B** matrices, robust least squares, predicate observations, class constraints, Kraitchman di-

agnostics, and GUI execution are part of one reproducible pipeline. The approach keeps the spectroscopic strengths of MSR while extending the coordinate model to the same GIC infrastructure used by the rest of Merlino.

References

- (1) Mendolicchio, M.; Penocchio, E.; Licari, D.; Tasinato, N.; Barone, V. Development and Implementation of Advanced Fitting Methods for the Calculation of Accurate Molecular Structures. *Journal of Chemical Theory and Computation* **2017**, *13*, 3060–3075.